



IJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 14 **Issue:** III **Month of publication:** March 2026

DOI: <https://doi.org/10.22214/ijraset.2026.78300>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

A Comprehensive Review of Deep Learning Approaches for Air Quality Pollution Analysis and Prediction

Sanjana Tiwari, Vishal Sharma

Abstract: Due to the rapid urbanization and industrialization, and the escalated vehicular emissions, air pollution has resulted in a critical global environmental and public health issue. The complexity of the spatiotemporal dynamics of pollutants is usually not well represented by conventional statistical and deterministic models. This paper presents a systematic review of recent deep learning methods for air quality prediction, including LSTM, GRU, CNN, autoencoders, and hybrid CNN-LSTM models. In contrast to the previous surveys that are confined to the performance evaluation of the models, the review takes a critical look at data integration strategies, the spatiotemporal modeling processes, the uncertainty estimation, the real time monitoring applications, and the feasible issues of the deployment. The research synthesizes the recent literature (2018-2025) findings and identifies the research gaps concerning the interpretability, generalization, and infrastructure requirements. In the review, the authors also discuss the application of deep learning to the process of smart city planning, health risk identification, and environmental policy-making. It shows that the hybrid deep learning models are much more effective than the traditional ones, and present-day challenges of the explainability and scalability of the solutions are necessary to implement deep learning models in the real-world setting.

Keywords: Air quality, pollution prediction, deep learning, spatiotemporal modeling, IoT sensors, environmental monitoring, health risk assessment.

I. INTRODUCTION

In the contemporary world, air quality pollution has become one of the most topical environmental and societal health issues, which are mainly promoted by the rapid industrialization, uncontrolled urbanization, the growing number of vehicles, the growth of the population and the widespread use of fossil fuels to obtain energy. The unremitting emission of toxic waste into the atmosphere has considerably impaired the quality of the air both in the developed and developing worlds and this is a real menace to the human health, ecological equilibrium and to the socioeconomic stability [1]-[3]. The most common examples of pollutants that are considered significant sources of air pollution include particulate matter that has less than 2.5 millimeters (PM 2.5) and 10 millimeters (PM 10) aerodynamic diameters, nitrogen dioxide (NO₂), sulfur dioxide (SO₂), carbon monoxide (CO), and ground-level ozone (O₃). The long-term consequences of such pollutants have been linked to numerous negative health implications, such as respiratory infections, asthma, chronic obstructive pulmonary disease, cardiovascular diseases, neurological disorders, and premature deaths, as well as to agricultural productivity, climatic changes, and the general quality of life. These impacts are so severe that, proper analysis and forecasting of the air quality levels is now necessary to help in environmental management, planning people health and policy formulation.




Fig. 1 Overview of Air Quality Pollution [4]

The traditional methods of air quality analysis and forecasting have been based on statistical methods, regression-based models, and deterministic atmospheric dispersion models that are premised on the laws of physics and chemistry. As much as these methods have been used to elucidate the process of air pollution, they tend to be limited in situations where the relationship is too nonlinear, temporal dynamics are too complicated, and the data is heterogeneous over a large scale. Most traditional models make severe assumptions, need much parameter optimization, and need much domain expertise, which limits their flexibility and precision, especially in dynamical urban settings where sources of pollution and meteorology vary quickly. Over the last several years, the development of new high-quality air quality and meteorological data records, as well as the Internet of Things (IoT) and open platforms in the environmental domain, have led to the creation of large amounts of high-resolution data. Such a rich data environment has led to data-driven modeling methods, with deep learning being one of the most powerful and promising solutions to air quality pollution analysis and prediction. Deep learning, also known as deep neural networks (DNNs), is an advanced field of machine learning, which uses multi-layered neural network structure and can automatically learn general hierarchical feature representations to raw data, thus capturing nonlinear dependencies and hidden patterns that are hard to model with other methods. Artificial Neural Networks (ANNs), CNNs, RNNs, Long Short-Term Memory (LSTM), and Gated Recurrent Units (GRUs) have shown strong performance in a wide range of time-series forecasting and spatiotemporal analysis tasks, thus making them especially appropriate to air quality prediction applications [5]-[7]. The data of the air pollution has a strong correlation in the time, seasonal variations, sudden changes, and long-term tendencies depending on the patterns of the emissions, human activity, and climate conditions, including temperature, humidity, wind speed, wind direction, precipitations, and atmospheric pressure. Long-term temporal dynamics, particularly those found in LSTM and GRU models, are better suited to be captured by recurrent deep learning models, which store memory of previous observations and learn long-range dependencies within a sequence of observations. Likewise, CNN-based methods are useful in capturing spatial patterns and learning the spatial distribution of pollutants in various monitoring points, and therefore, analyzing the spatial correlations and further transport of pollution at regional levels. The hybrid deep learning systems based on CNNs and RNNs can further boost the predictive accuracy because spatial and temporal specifics of air quality data are learned together. The use of deep learning models to analyze air quality pollution has a few important benefits such as deeper prediction capabilities, flexibility to various types of data sets, resiliency to noises and missing data and scaling with larger data sets [8], [9]. These models have the capability to combine the data of various sources like ground monitoring stations, meteorological sensors, satellite images and traffic data to come up with a more comprehensive and accurate prediction. Precise prediction of air quality is very important in building early warning systems capable of informing the masses and government about the impending events of air pollution so that preventive measures can be taken to avert exposure and health hazards.

II. LITERATURE REVIEW

Venkata et al. (2025) proposed a hybrid deep learning model (CNN-LSTM) to analyze the effects of air pollution on the productivity of agriculture. The experiment compared the historical level of pollutants and the trends of crop yields in order to define the crops that are resistant to pollution. The suggested method was effective in forecasting CO, SO₂, NO₃, O₃, PM_{2.5}, and AQI concentrations per hour. Predictive accuracy of performance assessment based on RMSE. The results indicate the opportunities of deep learning in assisting with agricultural planning and reducing yield losses in case of different air quality levels [15].

Mohammad et al. (2025) presented a predictive air quality model that combines the Recency, Frequency, and Monetary (RFM) model and the use of deep learning to enhance pollution observation in India. The system classifies the pollution events based on the RFM metrics to bring out a better understanding and predictability. The model was implemented in Python with the help of Google Colab and evaluated the regression and classification effectiveness. Comparison with the current neural networks showed a better reliability. The research postulates that smart air quality management and sustainable city planning measures can be achieved by integrating RFM and deep learning [16].

Ishan 2023 et al. carried out a comparative research on machine learning and deep learning models to predict Air Quality Index (AQI) in major Chinese cities. The models that were assessed were RNN, Bi-GRU, BiLSTM, CNN-BiLSTM, Conv1D-BiLSTM, and XGBoost. It was found that XGBoost was the most effective with an overall performance, and Conv1D-BiLSTM and CNN-BiLSTM were the most efficient deep learning architectures. Bi-GRU and RNN were found to be less accurate. The results show that machine learning and hybrid deep learning are both effective in the ability to capture nonlinear AQI patterns [17].

Ghose 2022 et al. created a hybrid deep learning model of AQI forecasting in intelligent cities by approximating spatial and time-related pollutant correlations. The suggested system combines 1D-CNN and Bi-GRU to present spatiotemporal relationships, and it is assisted by the algorithms of the missing data and the cross-location impact of pollutants. The framework was confirmed through

actual data of New South Wales, Australia. The experimental findings showed a better stability and prediction accuracy over prevailing models and this evidence shows the importance of considering spatial correlation in predicting urban air quality [18]. Bechkit 2021 et al. presented an uncertainty-constrained ConvLSTM-based framework of dynamically predicting air pollution during emergency cases. The model has a combination of convolutional layers with the extraction of spatial features and the LSTM with the temporal modeling features. Other layers of uncertainty quantification make predictions more reliable and interpretable. The framework was tested on the Fusion Field Trial 2007 data which depicts highly dynamic dispersion conditions. The experimental findings proved the existence of better performance (as opposed to the traditional machine learning and deep learning models) as the need to estimate the uncertainty in air quality management focused on the crisis [19].

Table 1 Literature Summary

Authors/Year	Methodology	Research gap	Findings
Hssina/2021[10]	Hybrid CNN-LSTM integrates spatial-temporal pollutant and meteorological data.	Limited models capture spatial dependencies across monitoring stations.	Hybrid model achieved highest accuracy among compared algorithms.
Abdelkader/2021[11]	IMDA-VAE integrates variational autoencoder with multiple directed attention.	Existing models lack adaptive attention across pollutants and locations.	IMDA-VAE outperformed VAE, LSTM, GRU, and ConvLSTM models.
Canyang/2020[12]	Pearson clustering with RNN, LSTM, GRU adaptive ensemble.	Single deep networks overfit, lacking correlation-driven feature integration.	Ensemble with Adam achieved lowest MAE and MAPE.
Zhou/2019[13]	Deep multi-output LSTM with dropout, L2, mini-batch training.	Shallow LSTM struggles with spatio-temporal instability, lag.	DM-LSTM improved accuracy, reduced errors across horizons.
Freeman/2018[14]	LSTM-RNN forecasts ozone using imputation and feature selection.	Limited long-range forecasts with reduced yet informative inputs.	Achieved MAE below two for seventy-two-hour predictions.

A. Research Gap

Despite significant advancements in deep learning-based air quality prediction, several limitations remain. Most existing studies focus primarily on improving predictive accuracy without systematically addressing issues such as model interpretability, cross-regional generalization, data quality inconsistencies, and real-time deployment constraints. Many reviews emphasize performance comparison but lack a structured synthesis of spatiotemporal modeling strategies, multi-source data fusion techniques, and uncertainty-aware forecasting approaches. There is also limited discussion on the integration of deep learning models into practical environmental governance frameworks and smart city infrastructures. Therefore, a comprehensive review that critically evaluates methodological advancements while identifying deployment challenges and future research directions is necessary.

B. Contribution of this Review

This review makes the following distinct contributions:

- 1) Provides a structured comparison of deep learning architectures for spatiotemporal air quality modeling.
- 2) Synthesizes recent advancements including hybrid and uncertainty-aware models.
- 3) Analyzes practical implementation challenges such as data heterogeneity, computational demands, and interpretability.
- 4) Highlights the integration of deep learning with IoT, satellite systems, and smart city planning frameworks.
- 5) Identifies future research directions emphasizing explainable AI and scalable deployment strategies.

III. OVERVIEW OF AIR QUALITY POLLUTION AND MONITORING SYSTEMS

Air quality pollution is the pollution of the atmosphere by chemicals that pollute the environment and the living organisms. These substances may be gases, particulates or biological molecules whether naturally introduced or by human intervention. The impact of low air quality may be very severe health issues, as respiratory diseases and cardiovascular complications, and degradation of ecosystems. The magnitude and nature of the pollution depends on place, time and sources [20], [21]. It is imperative to understand what air pollution is in order to identify the causes, effects and the necessity to monitor and control it to safeguard the health of the people and the quality of the environment.

A. Common Air Pollutants

Air pollutants refer to a very broad category of dangerous chemicals and microscopic assemblies suspended in the atmosphere, with many of them being extremely harmful to human health and the environment. However, one of them is of even greater concern: particulate matter (PM) and, in particular, small particles, PM_{2.5} and PM₁₀, since in this way they can reach deep into the respiratory system and can even get to the lungs and even the bloodstream. This may cause diseases of the respiratory system, heart problems, and other severe health complications. Along with the particulate matter, there are a number of gaseous pollutants that cause high air pollution.



Fig. 2 Common Air Pollutants [22]

These are nitrogen oxides (NO_x), sulfur dioxide (SO₂), carbon monoxide (CO) ozone (O₃) and volatile organic compounds (VOCs), and their sources are vehicle emissions, industries, power generation and chemical reactions in the atmosphere [23]. The effects of these pollutants differ: NO_x and SO₂ are significant causes of acid rain, ozone in the ground level is a source of harmful smog. In many cases, these contaminants are chemically reactive and their negative impacts on air quality, the environment, and human health are compounded, thus making it more difficult to reduce them.

B. Sources of Air Pollution:

Pollution of the air is caused by numerous natural and artificial sources with various impacts on the quality of the atmosphere. Events and processes that are natural bear the name of wildfires, volcanic eruptions, dust storms, and the release of pollen all of which inject some particulate matter and gases into the air. Although these natural effects add to the levels of pollution in the background, human activities are the main cause of the harmful air pollution in the present day. Power plants and factories emit large quantities of pollutants through industrial emissions, and the vehicle exhaust is a major contributor of nitrogen oxides and particulate matter.




Fig. 3 Sources of Air Pollution [24]

Also, building mechanisms produce dust and agricultural processes, which involve fertilizers and pesticides, produce volatile compounds and ammonia [25]. The high rate of urbanization and industrialization across the world has only increased the level of pollution and thus has become a major issue of concern globally. Identification and awareness of these multiple sources is the key to generating effective and specific mitigation measures to minimize the emissions at the origin and eventually enhance the quality of the air and the health of people.

C. *Impact of Air Pollution*

The impacts of air pollution on the human health are extensive and far-reaching and lead to a broad spectrum of severe medical disorders. Polluted air can result in or aggravate respiratory infections like asthma and bronchitis as well as exposing people to the risk of cardiovascular illnesses, including heart attacks. Excessive exposure to high levels of pollution may result in untimely deaths in serious cases. Some of the groups such as children, the elderly and people with pre-existing health conditions are particularly susceptible to such adverse effects. In addition to human health, air pollution is also a major problem in agriculture as it leads to a decreased yield and quality of crops. It also destroys forests and wild life and disturbs fragile ecosystems and biodiversity [26]. Pollution causes acid rain including sulfur dioxide and nitrogen oxides that destroys the soil, fresh water bodies and plants.

D. *Importance of Monitoring Air Quality*

Air quality monitoring is important in environmental pollution management and understanding it as it gives the necessary information regarding the level of pollution and changes in the pollution over time. By properly and consistently collecting the data, the authorities will be able to locate the areas that have a high concentration of pollution, which are called pollution hotspots, and monitor the daily or seasonal air quality movements. The information proves crucial in the implementation of the environmental regulations and environmental standards, so that the policymakers can come up with specific strategies and action plans that will curb emissions by major sources [27]. Monitoring air quality is important to safeguard human health since this process assists in prompting alert and health warnings in case of risky air pollution to enable the community to take preventive action.

E. *Types of Air Quality Monitoring Systems*

There is a great variety of air quality monitoring systems with the focus on the technologies that they apply and the goals that they can achieve, with their pros and cons. Fixed monitoring stations, which are normally operated by government agencies, and have high precision advanced instruments, which give precise and reliable data at certain fixed locations at specific locations. They are normally limited in coverage as they are expensive and logistically difficult to install and maintain these stations. To supplement this, there are portable sensors which have been created in order to provide a high level of flexibility enabling air quality measurements to be performed in different locations or to be utilized to measure individual exposure in real time [28]. Satellite remote sensing can offer a wide perspective that is large-scale, i.e. satellite images are able to compile data on air pollution of huge regions or the planet, which comes in handy particularly in tracking transboundary pollution and long-term trends.

F. *Advances in Monitoring Technologies*

The improvement of technology has introduced much change in the air quality monitoring system and it has become more convenient, precise and effective. The emergence of cheap sensors and Internet of Things (IoT) technology has facilitated the introduction of monitoring equipment in cities and various rural settings in large numbers where data can be collected on a real-time basis and in very high detail regarding the extent of pollution. Remote sensing technologies that make use of satellite technology are used to supplement these other systems and provide a wide spatial area coverage, especially in areas that are inaccessible to standard stations, or in remote areas that need to be observed [29]. The combination of innovative data analytics and machine learning methods will increase the possibility of detecting sources of pollution, operating on intricate data trends, and developing forecasting models that predict the dynamic of pollution.

IV. TRADITIONAL AIR QUALITY PREDICTION METHODS

Deep learning is a sophisticated subdivision of machine learning, which applies the model of neural networks that are several layers deep to sort out complex data trends. Deep learning has received great interest in the context of air quality analysis due to its capacity to predict nonlinear and complex associations between environmental variables. Conventional statistical tools are not always able to adapt to the variability and complexity of air pollution data, whilst deep learning can be able to promote latent characteristics and interactions [30].

The models have the ability to handle vast amounts of data at once and the data can be collected by a weather station, sensor or a satellite which is why they are very well applicable to the air quality forecasting and evaluation.

A. Data Types Used in Deep Learning for Air Quality

Deep learning-based air quality models are based upon diverse and rich datasets to understand the entire nature of pollution dynamics. Important types of data are the real-time measurements of the pollutants such as PM_{2.5}, PM₁₀, nitrogen dioxide (NO₂), and ozone (O₃) in ground monitoring stations. Weather conditions are important context in terms of meteorological data such as temperature, humidity, speed and direction of wind which determine the distribution and formation of pollutants. The satellite imagery also provides a spatial aspect as it provides extensive details about the distribution and sources of pollution [31]. Secondly, traffic flow data, and industrial emission records would also add data on the sources of anthropogenic pollution. Combining these mixed types of data, deep learning models are able to have a full picture of air quality trends and enhance their predictive quality. The multi-source data fusion is also capable of analyzing the time variation (hourly, daily, seasonal) and space variation, which is crucial in managing and predicting air pollution.

B. Common Deep Learning Architectures Used in Air Quality

Certain deep learning structures are especially suitable to air quality analysis to which each of them is sensitive. Convolutional Neural Networks (CNNs) have a strong performance in spatial data processing, i.e. satellite images or sensor grids, as they can automatically detect spatial features and pollution hotspots. RNNs (with their more sophisticated form, Long Short-Term Memory networks also known as LSTM networks) are useful in working with time-series data such as hourly pollutant levels, as they model time-varying relationships and patterns [32]. Autoencoders are unsupervised models that are used to reduce the dimensions of data and identify anomalies in the data on air quality like the sudden spikes in pollution. The architectures can be integrated or tailored to have hybrid models to take advantage of their respective strengths. The choice of the architecture is determined by the features of the input data and the purpose, i.e., to predict the level of pollution in the future, to find anomalies, or to identify the sources of pollution.

C. Advantages of Deep Learning Over Traditional Methods

There are important advantages of deep learning over conventional statistical or machine learning in the analysis of air quality. Among the advantages, one should mention the fact that it can be applied to model nonlinear relationships and complex interactions between a number of variables in the environment without assuming or selecting features explicitly. Deep learning models have the advantage of deriving significant features of raw data automatically, eliminating human bias and pre-processing efforts. They are also the best in dealing with huge amounts of high-dimensional, noisy and incomplete data that are typical of air pollution monitoring [33]. This strength is transferred to a better predictive accuracy and extrapolation to unseen data. In deep learning, various types of data can be combined together, e.g. images, time series and categorical information, in a single framework. The strengths facilitate more active and flexible air quality models that eventually result in more informed environmental protection and population health decisions.

V. DEEP LEARNING MODELS USED IN AIR QUALITY PREDICTION

The predictive models of air quality have become potent because the deep learning models can learn intricate patterns of large volumes of data. The models can process environmental data that are nonlinear and of high-dimension, which are common in environmental monitoring. Deep learning models are able to predict pollutant concentrations more effectively than a large number of conventional approaches by taking past pollution information, weather, and other appropriate inputs [34], [35]. Their ability to be used with diverse data including time-series measurements and spatial data is what renders them suitable to the dynamic and multifaceted nature of the air pollution. This adaptability has seen popularization of deep learning methods in air quality forecasting, which has assisted governments and other environmental bodies in making sound decisions to safeguard human lives.

A. Recurrent Neural Networks (RNNs) and LSTM for Time-Series Prediction

The common type of Neural Networks known as Recurrent Neural networks (RNNs) and the sophisticated form known as Long Short-Term memory (LSTM) networks are popular in air quality prediction due to their ability to process sequential data. The level of air pollution changes with time, and the time-series forecast is, therefore, necessary. RNNs are sequential processors of data, where the past information of the data is remembered to indicate time dependencies.

Nevertheless, conventional RNNs are associated with such issues as vanishing gradients that LSTMs address with such special memory cells that help to remember significant information over time. This is why LSTMs are especially applicable when it comes to predicting hourly or daily pollutant concentrations, which are characterized by a complicated temporal pattern and patterns based on weather variations, traffic, and industrial activity. Time-dependent pollution data have been successfully modelled with them, which has rendered them one of the most popular deep learning models used in air quality forecasting [36].

B. Convolutional Neural Networks (CNNs) for Spatial Analysis

Convolutional Neural Networks (CNNs) are deep learning models that are very useful in the analysis of spatial data, thus they are of great use in predicting air quality because geographic variability is a critical factor in air quality. The mechanism of CNNs involves the use of numerous convolutional filters in input data, which can be satellite images, sensor network grids, or maps of pollution distribution, and thus help the CNNs detect meaningful spatial information and patterns. Such patterns can be pollution hotspots or places of pollutant concentration, or dispersal with regard to the local environment conditions. CNNs increase the prediction accuracy of the pollutant concentrations in various locations by capturing local spatial dependencies of the data [37]. This is more especially in an urban setting whereby sources of pollution, topography and the atmosphere of a neighborhood differ greatly across neighborhoods. CNNs are commonly combined with temporal models such as the Long Short-Term Memory networks (LSTMs) to create hybrid networks that can more accurately predict air pollution and its dynamics, since they help track the development of pollution in space and time at the same time.

C. Autoencoders for Feature Extraction and Anomaly Detection

Autoencoders are a category of unsupervised deep learning networks that are commonly applied to air quality analysis to perform such tasks as dimensionality reduction and anomaly detection. The principle of these models is to reduce the cost of high-dimensional input data to a low-dimensional latent space, which defines the key properties of the data and discards unnecessary or redundant data. Once the data has been compressed, the autoencoder is used to recreate the original information using this compressed representation, where landmark features are learned without the need to use labeled datasets. Such a capability to extract relevant patterns is useful in reducing the noise and enhancing the quality of data to be used in a further prediction or classification model more efficient and accurate. Autoencoders can be used especially in the field of air quality monitoring to identify an abnormal pollution incident or sensor malfunction by monitoring data points that do not fit the normal predictive patterns.

D. Hybrid Models Combining Multiple Architectures

Hybrid training approaches based on CNNs, RNNs, LSTMs and autoencoders have been created to exploit the advantages of various deep learning architectures in air quality forecasting. The models are capable of acquiring spatial, temporal and feature-level information simultaneously resulting in enhanced accuracy and strength. As an illustration, a CNN-LSTM hybrid takes CNN layers to retrieve the spatial characteristics of the pollution maps or sensor grid and then feeds the characteristics to LSTM layers to establish time dynamics. This is due to the fact that such integrated models can address the complexity of air pollution as compared to single architectures. The hybrid models are especially applicable in complex urban environments with changing patterns of pollution, especially in time and space, and which assist the policymakers and environmental managers to predict the pollution episodes more appropriately and set up mitigation measures to be taken.

VI. CHALLENGES IN DEEP LEARNING-BASED AIR QUALITY PREDICTION

Various complex issues that influence the performance and practical use of deep learning-based air quality prediction have interdependence. To do reliable modeling, large historical data is needed, but monitoring data often has missing values, variable format, and calibration mistakes, where intensive preprocessing is required. There are intricate interactions between meteorology, emissions, traffic activity and the geographic features that control pollution behavior which instigates nonlinear patterns that are challenging even to advanced neural networks to embrace across space and time [38].

A. Data Availability and Quality

Deep learning algorithms require huge amounts of trustworthy historical information to train on significant trends and make correct predictions. Nevertheless, in air quality, this type of data is often subject to a variety of constraints. The monitoring stations can be affected by interruptions caused by maintenance, power outage or communication error thus missing or recording incomplete

records. Monitoring infrastructure in most developing or remote areas is relatively recent and hence observation periods are brief and not adequate to record long term seasonal and annual changes. The inconsistencies are further triggered by sensor calibration issues, measurement drifts, and equipment inter-network variation. Also, the data gathered by more than one agency can frequently be presented in heterogenous form and needs a lot of preprocessing and harmonization to be analysed [39].

B. Complex Spatiotemporal Relationships

Patterns of air pollution are complex products of interaction between a web of factors which are constantly undergoing change and interact in complex and generally nonlinear ways. The meteorological variables that control the dispersion of the pollution, the accumulation or the chemical transformations of the pollutants are wind speed, wind direction, temperature, humidity, and atmospheric pressure. Meanwhile, the sources of emissions can be different based on the time of the day and place, depending on traffic concentration, industrial time, building works, and domestic fuel consumption. The geographics, topography, vegetation and proximity to highways or factories also have a more specific influence on the movement and persistence of contaminants. These influences work in concert at many scales, ranging between the street level down to the regional transport and also changing within minutes, days and seasons [40]. Even more complex neural networks such as LSTM, GRU and spatiotemporal neural networks may not be able to capture all these interactions effectively. Models can therefore simplify relationships or may not foresee sudden changes thereby restricting the level of forecasting accuracy and strength.

C. High Computational and Infrastructure Demands

The operationalization and development of deep learning systems to predict air quality require significant infrastructural requirements in computational power. Long Short-Memory Architectures like Long Short-Memory, Gated Recurrent Units (GRU), and hybrid Convolutional Neural Network (CNN) formulations are architectures made up of millions of parameters that need to be optimized by repeated training steps. Such a process involves hardware of high performance, including, but not limited to, GPUs or otherwise specialized accelerators, and much memory to hold large spatio-temporal data. More so, preprocessing steps that include data cleaning, normalization and feature extraction introduce additional computational overhead [41].

D. Limited Interpretability and Transparency

Most deep learning models produce very sensible results, but the line of thinking that resulted in such predictions is frequently non-understandable. The multi-layered designs they have process inputs with sophisticated mathematical computations which give results that are hard to attribute to particular variables or causal links. In the context of the air quality management, such absence of transparency is a critical issue since policymakers, environmental authorities, and health-related organizations should be capable of justifying their decisions in terms of traffic bans, industrial closures, or emergency warnings. Unless there are clear explanations, stakeholders might not have trust in automated systems, particularly when the decision has economic or even social implications [42]. Reduced interpretability complicates the ability of a researcher to diagnose and detect bias as well as enhance model design. The lack of evidence that can be understood can also decrease the acceptance of forecasts by the people.

E. Model Generalization and Maintenance

Air quality prediction models based on deep learning are typically trained on data and information preserved in a particular geographic place and time, making their application to other areas potentially limited. The sources of pollution, urban form, industrial operation, transportation pattern, and regulatory operation differ across regions, establishing different emission patterns. Atmospheric chemistry and dispersion processes are further changed by seasonal factors like changes in temperature, monsoon, heating needs, or farmfire. The model, which was trained in one environment, might not understand the relationships or be unable to perceive emergent patterns when transferred to another environment [43].

VII. APPLICATIONS OF DEEP LEARNING IN AIR QUALITY MANAGEMENT

Deep learning has significantly transformed air quality management by enabling accurate prediction, real-time monitoring, and intelligent decision support. Advanced neural architectures process large-scale spatiotemporal data from monitoring stations, satellites, and IoT networks.

These models support early warning systems, emission source identification, health risk estimation, and smart urban planning, thereby promoting proactive, data-driven, and sustainable environmental governance [18].

A. Air Quality Forecasting

Deep learning has now emerged as a potent forecasting device of future pollutant concentrations in the air through the patterns of past data and weather. Particularly good models in capturing the temporal dependence, seasonality, and nonlinear relationship among variables of temperature and humidity and wind behavior and the intensity of emission include the Long Short-Term Memory (LSTM), Gated Recurrent Units (GRU) and Convolutional Neural Networks (CNN) [44], [45]. These systems are also able to produce both the short term hourly predictions and the long term predictions to help in proactive environmental management. Proper forecasting enables the government to give out health warnings, control traffic movements and temporarily shut down industrial activities before the pollution becomes dangerous.

B. Real-Time Pollution Monitoring

Deep learning is an important aspect in improving the real-time monitoring capabilities because it handles continuous data feeds of ground sensors, satellites, and IoT-based devices. Neural networks are able to comprehend complicated relationships as they occur and detect minor shifts in pollutant behavior unlike conventional threshold-based systems. It allows identifying worsening air conditions faster and helps them to generate automatic alerts. Real-time analytics are also used to provide visualization on the distribution of pollution across neighborhoods, highways, and industrial areas to the environmental agencies. This sort of information is needed to provide dynamic responses, such as diversion of the traffic, controlling of the emissions, or social warning. Moreover, adaptive systems are capable of learning on the new incoming data, and as time goes by, it becomes more accurate [46].

C. Source Identification and Apportionment

Knowledge of the source of pollution is the basis of developing mitigation policies. Deep learning models help to determine the probable sources of emissions based on the spatial distribution, wind patterns, traffic volumes, industrial production and land use. Neural networks can approximate the relative contribution of vehicles, factories, construction sites, or natural events by finding hidden correlations in big data. This is more applicable in urban areas that are complicated and a combination of sources is active at any time. Proper source apportionment allows policymakers to take specific measures, instead of general restraints, which facilitates less economic havoc [47].

D. Health Risk Assessment

Deep learning can play an important role in assessing the possible health effects of air pollution exposure. On connecting the levels of pollutants with the demographic information, with hospitalization, and epidemiological data, the models will be able to calculate the levels of risks among the various groups in the population. Such forecasts assist the governments in foreseeing the rise of respiratory or heart related diseases and prepare the medical facilities against such occurrences.

VIII. CONCLUSION

To sum up, air quality pollution is a form of global challenge with grave consequences on the health of the population, environmental sustainability and socio-economic development that require sophisticated analytical and predictive solutions to this problem. Conventional air quality monitoring and forecasting methods would not be sufficient to detect the complex, nonlinear and dynamic interactions between pollutants, meteorological factors, source of emissions and geographic location. Deep learning models have proven to be a strong and efficient solution since they allow extracting features automatically, including heterogeneous data sources, and precisely predicting the level of the pollutant in space and time. Convolutional neural networks, long short term memory networks, autoencoders, and hybrid networks have proven effective in air quality prediction, real-time prediction, anomalies, and source location as well as health risk assessment.

These features help in active environmental management, through ease of early warning mechanism, mitigation plans, and formulation of policy based on evidence. However, the application of the deep learning-based air quality systems is still limited by the challenges such as scarce data availability and quality, high computational and infrastructure requirements, model interpretability, and the inability to extrapolate predictions to a different area and time. To overcome these constraints, data needs to be enhanced, and scalable computing infrastructure will need to be invested in, as well as explainable artificial intelligence methods to increase transparency and trust.

Even with these problems, the incorporation of deep learning into air quality analysis is an important step in the direction of smart and data-driven environmental regulation.

REFERENCES

- [1] K. Chan, P. Matthews, and K. Munir, "Time Series Forecasting for Air Quality with Structured and Unstructured Data Using Artificial Neural Networks," *Atmosphere (Basel)*, vol. 16, no. 3, pp. 1–17, 2025, doi: 10.3390/atmos16030320.
- [2] S. Alkayal, H. Almishahi, S. Baowidan, and E. Alkayal, "Air Pollution Trends and Predictive Modeling for Three Cities with Different Characteristics Using Sentinel-5 Satellite Data and Deep Learning," *Atmosphere (Basel)*, vol. 16, no. 2, 2025, doi: 10.3390/atmos16020211.
- [3] I. Ganguli, M. Nakum, B. Das, and N. Kshetrimayum, "Comprehensive Analysis of Air Quality Trends in India Using Machine Learning and Deep Learning Models," *ICDCN 2025 - Proc. 26th Int. Conf. Distrib. Comput. Netw.*, no. January 2025, pp. 313–318, 2025, doi: 10.1145/3700838.3703681.
- [4] T. Istiana *et al.*, "Causality Analysis of Air Quality and Meteorological Parameters for PM_{2.5} Characteristics Determination: Evidence from Jakarta," *Aerosol Air Qual. Res.*, vol. 23, no. 9, p. 230014, Sep. 2023, doi: 10.4209/aaqr.230014.
- [5] R. Zrieq, S. Kamel, F. Al-Hamazani, S. Boubaker, R. Attili, and M. J. Araúzo-Bravo, "Spatial-Temporal Forecasting of Air Pollution in Saudi Arabian Cities Based on a Deep Learning Framework Enabled by AI," *Toxics*, vol. 13, no. 8, pp. 1–31, 2025, doi: 10.3390/toxics13080682.
- [6] H. Madokoro and S. Nix, "Multimodal Particulate Matter Prediction: Enabling Scalable and High-Precision Air Quality Monitoring Using Mobile Devices and Deep Learning Models," *Sensors*, vol. 25, no. 13, pp. 1–23, 2025, doi: 10.3390/s25134053.
- [7] S. Jayaraman, N. T. A. S., and S. G., "Enhancing urban air quality prediction using time-based-spatial forecasting framework," *Sci. Rep.*, vol. 15, no. 1, pp. 1–15, 2025, doi: 10.1038/s41598-024-83248-z.
- [8] A. K. Rad, M. J. Nematollahi, A. Pak, and M. Mahmoudi, "Predictive modeling of air quality in the Tehran megacity via deep learning techniques," *Sci. Rep.*, vol. 15, no. 1, pp. 1–20, 2025, doi: 10.1038/s41598-024-84550-6.
- [9] I. Zandi, A. Jafari, and A. Lotfata, "Enhancing PM_{2.5} Air Pollution Prediction Performance by Optimizing the Echo State Network (ESN) Deep Learning Model Using New Metaheuristic Algorithms," *Urban Sci.*, vol. 9, no. 5, 2025, doi: 10.3390/urbansci9050138.
- [10] A. Bekkar, B. Hssina, S. Douzi, and K. Douzi, "Air-pollution prediction in smart city, deep learning approach," *J. Big Data*, vol. 8, no. 1, pp. 1–21, 2021, doi: 10.1186/s40537-021-00548-1.
- [11] A. Dairi, F. Harrou, S. Khadraoui, and Y. Sun, "Integrated Multiple Directed Attention-Based Deep Learning for Improved Air Pollution Forecasting," *IEEE Trans. Instrum. Meas.*, vol. 70, 2021, doi: 10.1109/TIM.2021.3091511.
- [12] C. Guo, G. Liu, and C. H. Chen, "Air Pollution Concentration Forecast Method Based on the Deep Ensemble Neural Network," *Wirel. Commun. Mob. Comput.*, vol. 2020, 2020, doi: 10.1155/2020/8854649.
- [13] Y. Zhou, F. J. Chang, L. C. Chang, I. F. Kao, and Y. S. Wang, "Explore a deep learning multi-output neural network for regional multi-step-ahead air quality forecasts," *J. Clean. Prod.*, vol. 209, pp. 134–145, 2019, doi: 10.1016/j.jclepro.2018.10.243.
- [14] B. S. Freeman, G. Taylor, B. Gharabaghi, and J. Thé, "Forecasting air quality time series using deep learning," *J. Air Waste Manag. Assoc.*, vol. 68, no. 8, pp. 866–886, 2018, doi: 10.1080/10962247.2018.1459956.
- [15] S. J. *et al.*, "Smart Agriculture: Integrating Air Quality Monitoring with Deep Learning for Process Optimization," *Scalable Comput. Pract. Exp.*, vol. 26, no. 3, pp. 1005–1016, 2025, doi: 10.12694/scpe.v26i3.4190.
- [16] J. Mohammad and M. A. Kashem, "Deep Learning Approach for Evaluating Air Pollution Using the RFM Model," *Nat. Environ. Pollut. Technol.*, vol. 24, no. 2, 2025, doi: 10.46488/NEPT.2025.v24i02.D1718.
- [17] I. Ayus, N. Natarajan, and D. Gupta, "Comparison of machine learning and deep learning techniques for the prediction of air pollution: a case study from China," *Asian J. Atmos. Environ.*, vol. 17, no. 1, 2023, doi: 10.1007/s44273-023-00005-w.
- [18] B. Ghose and L. Anthopoulos, "A Deep Learning based Air Quality Prediction Technique Using Influencing Pollutants of Neighboring Locations in Smart City," vol. 28, no. 8, pp. 799–826, 2022, doi: 10.3897/jucs.78884.
- [19] I. Mokhtari, W. Bechkit, H. Rivano, and M. R. Yaici, "Uncertainty-Aware Deep Learning Architectures for Highly Dynamic Air Quality Prediction," *IEEE Access*, vol. 9, pp. 14765–14778, 2021, doi: 10.1109/ACCESS.2021.3052429.
- [20] F. Naz *et al.*, "Comparative Analysis of Deep Learning and Statistical Models for Air Pollutants Prediction in Urban Areas," *IEEE Access*, vol. 11, no. June, pp. 64016–64025, 2023, doi: 10.1109/ACCESS.2023.3289153.
- [21] S. Hameed, A. Islam, K. Ahmad, S. B. Belhaouari, J. Qadir, and A. Al-Fuqaha, "Deep learning based multimodal urban air quality prediction and traffic analytics," *Sci. Rep.*, vol. 13, no. 1, pp. 1–19, 2023, doi: 10.1038/s41598-023-49296-7.
- [22] G. T. M. N. Anwar, G. T. M. Navi Anwar, H. P. Gunasekar, J. Krishnan, and R. Sekar, "Air quality monitoring and assessment using IoT," *IOP Conf. Ser. Mater. Sci. Eng.*, vol. 955, no. 1, Nov. 2020, doi: 10.1088/1757-899X/955/1/012006.
- [23] S. Aguilar-Gomez, B. Dwyer, J. Graff Zivin, and M. Neidell, "This is Air: The 'Non-Health' Effects of Air Pollution," *SSRN Electron. J.*, 2022, doi: 10.2139/ssrn.4062400.
- [24] A. Kurasz, G. Y. H. Lip, S. Dobrzycki, and Ł. Kuźma, "A Breath of Trouble: Unraveling the Impact of Air Pollution on Atrial Fibrillation," *J. Clin. Med.*, vol. 13, no. 23, Dec. 2024, doi: 10.3390/jcm13237400.
- [25] X. Wang, G. Wu, Z. Xiang, and J. Zhang, "Air Pollution and Media Slant: Evidence from Chinese Corporate News," *Emerg. Mark. Financ. Trade*, vol. 58, no. 10, pp. 2880–2894, 2022, doi: 10.1080/1540496X.2021.2013196.
- [26] K. Vohra *et al.*, "Rapid rise in premature mortality due to anthropogenic air pollution in fast-growing tropical cities from 2005 to 2018," *Sci. Adv.*, vol. 8, no. 14, 2022, doi: 10.1126/sciadv.abm4435.
- [27] M. J. Wolf *et al.*, "New Insights for Tracking Global and Local Trends in Exposure to Air Pollutants," 2022, doi: 10.1021/acs.est.1c08080.
- [28] A. Dun, Y. Yang, and F. Lei, "A Novel Hybrid Model Based on Spatiotemporal Correlation for Air Quality Prediction," *Mob. Inf. Syst.*, vol. 2022, 2022, doi: 10.1155/2022/9759988.
- [29] L. Montalvo, D. Fosca, D. Paredes, M. Abarca, and C. Saito, "An Air Quality Monitoring and Forecasting System for Lima City With Low-Cost Sensors and Artificial Intelligence Models," vol. 4, no. July, pp. 1–15, 2022, doi: 10.3389/frsc.2022.849762.
- [30] P. O. Czechowski *et al.*, "Financing Costs and Health Effects of Air Pollution in the Tri-City Agglomeration," *Front. Public Heal.*, vol. 10, no. March, pp. 1–10, 2022, doi: 10.3389/fpubh.2022.831312.
- [31] U. Im *et al.*, "Reviewing the links and feedbacks between climate change and air pollution in Europe," no. September, pp. 1–19, 2022, doi: 10.3389/fenvs.2022.954045.

- [32] Q. Zhang, Y. Han, V. O. K. Li, and J. C. K. Lam, "Deep-AIR: A Hybrid CNN-LSTM Framework for Fine-Grained Air Pollution Estimation and Forecast in Metropolitan Cities," *IEEE Access*, vol. 10, pp. 55818–55841, 2022, doi: 10.1109/ACCESS.2022.3174853.
- [33] A. Choudhary *et al.*, "Time Series Simulation and Forecasting of Air Quality Using In-situ and SatelliteBased Observations over an Urban Region," *Nat. Environ. Pollut. Technol.*, vol. 21, no. 3, pp. 1037–1048, 2022, doi: 10.46488/NEPT.2022.v21i03.018.
- [34] M. G. Vivanco *et al.*, "Assessment of the Effects of the Spanish National Air Pollution Control Programme on Air Quality," 2021.
- [35] A. Dutta and W. Jinsart, "Air Pollution in Indian Cities and Comparison of MLR , ANN and CART Models for Predicting PM 10 Concentrations in Guwahati , India," vol. 15, no. 1, 2021.
- [36] V. Thi, T. Minh, T. T. Tin, and T. T. Hien, "PM 2.5 Forecast System by Using Machine Learning and WRF Model , A Case Study : Ho Chi Minh City ," vol. 21, no. 12.
- [37] G. Huang *et al.*, "Large scale air pollution prediction with deep convolutional networks," *Sci. China Inf. Sci.*, vol. 64, no. 9, 2021, doi: 10.1007/s11432-020-2951-1.
- [38] M. Arsov, E. Zdravevski, R. Corizzo, N. Koteli, and K. Mitreski, "Short-term air pollution forecasting based on environmental factors and deep learning models," vol. 21, pp. 15–22, 2020, doi: 10.15439/2020F211.
- [39] L. Vitali *et al.*, "A standardized methodology for the validation of air quality forecast applications (F-MQO): lessons learnt from its application across Europe," *Geosci. Model Dev.*, vol. 16, no. 20, pp. 6029–6047, 2023, doi: 10.5194/gmd-16-6029-2023.
- [40] Y. Yarragunta, S. Srivastava, D. Mitra, and H. Chandra, "Influence of forest fire episodes on the distribution of gaseous air pollutants over," *GIScience Remote Sens.*, vol. 57, no. 2, pp. 190–206, 2020, doi: 10.1080/15481603.2020.1712100.
- [41] Y. A. Y. Okkaoğlu and E. G. M. E. Yücel, "Estimation and forecasting of - PM 10 air pollution in Ankara via time series and harmonic regressions," *Int. J. Environ. Sci. Technol.*, vol. 17, no. 8, pp. 3677–3690, 2020, doi: 10.1007/s13762-020-02705-0.
- [42] S. Lolli, Y. C. Chen, S. H. Wang, and G. Vivone, "Impact of meteorological conditions and air pollution on COVID-19 pandemic transmission in Italy," *Sci. Rep.*, vol. 10, no. 1, pp. 1–15, 2020, doi: 10.1038/s41598-020-73197-8.
- [43] Di. Madaan, R. Dua, P. Mukherjee, and B. Lall, "Vayuanukulani: Adaptive memory networks for air pollution forecasting," *Glob. 2019 - 7th IEEE Glob. Conf. Signal Inf. Process. Proc.*, 2019, doi: 10.1109/GlobalSIP45357.2019.8969343.
- [44] H. Wang, B. Zhuang, Y. Chen, N. Li, and D. Wei, "Deep inferential spatial-temporal network for forecasting air pollution concentrations," *arXiv*, 2018.
- [45] S. He, S. Tang, Y. Xiao, and R. A. Cheke, "Stochastic Modelling of Air Pollution Impacts on Respiratory Infection Risk," *Bull. Math. Biol.*, vol. 80, no. 12, pp. 3127–3153, 2018, doi: 10.1007/s11538-018-0512-5.
- [46] R. San José, J. L. Pérez, L. Pérez, and R. M. Gonzalez Barras, "Effects of climate change on the health of citizens modelling urban weather and air pollution," *Energy*, vol. 165, pp. 53–62, 2018, doi: 10.1016/j.energy.2018.09.088.
- [47] J. Bousquet *et al.*, "POLLAR: Impact of air POLLution on Asthma and Rhinitis; A European Institute of Innovation and Technology Health (EIT Health) project," *Clin. Transl. Allergy*, vol. 8, no. 1, pp. 1–13, 2018, doi: 10.1186/s13601-018-0221-z.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)