



IJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 14 **Issue:** VI **Month of publication:** June 2026

DOI: <https://doi.org/10.22214/ijraset.2026.83578>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

A Comprehensive Review on Phytochemical and Pharmacological Properties of Wood Apple (*Feronia Limonia*)

Siddhi Jadhav¹, Sonam Jadhav², Sushant Jagdale³, Mrs. Ashwini Dhole⁴, Dr. Rajendra Patil⁵

Delonix Society's Baramati College of Pharmacy, Barhanpur

Abstract: *Wood Apple (*Feronia limonia* L.), a member of a Rutaceae family, is an underutilized yet nutritionally and medicinally important fruit widely found in India and other tropical regions. Traditionally regarded as a "poor man's fruit", it has gained attention due to its rich phytochemical composition and diverse therapeutic properties. The fruit grows in semi-arid and dry climates, demonstrating high adaptability and draught tolerance. Despite its abundance, its commercial cultivation and utilization remain limited. Pharmacological studies have revealed multiple biological activities of wood apple, including antidiabetic, antimicrobial, anticancer, antioxidant, hepatoprotective, antiulcer, and anti-inflammatory effects. Experimental studies demonstrate its ability to reduce blood glucose level, improve lipid profiles, and enhance antioxidant enzyme activity. Additionally, extracts of wood apple have shown inhibitory effects against several pathogenic microorganism, supporting its antimicrobial potential.*

Keywords: *Wood Apple (*Feronia limonia*), phytochemicals, antioxidant activity, pharmacological properties, antidiabetic activities, nutraceuticals potential.*

I. INTRODUCTION

Most popular underutilized fruit crop in India is the wood apple, a prickly tree. Its botanical name is *Feronia* (*F. limonia* L.), and it is a member of the Rutaceae family [1]. Before processing the methods were established in the mid-1950s, wood apple fruit was considered a "poor man's food" in India [2]. The tree is given the genus *Feronia* in honor of the Roman goddess of forests [3]. Since elephants like it, it is sometimes known as monkey fruit or elephant apple in addition to wood apple. Other common names for wood apples include "curd apple," "golden apple," "stone apple," and so on; these names usually depend on the language, place, and culture. Rural residents believe that the animal's digestive tract has a special capacity to break down the fruit's insides without sacrificing its hard exterior. Consequently, when the animal excretes, the fruit appears whole[4,5]. Other dialect names in Thailand include ma-khwit, Kramsang in Cambodia, Ma-fi in Laos, Gelinggai or Belinggai in Malaysia, pomme d'elept, hanpomme de bois, or citron des mois in French, Tuffâhh elfil in Arabic, Kathbelin in Bangladesh, Thibin, Thanaka, and Tha nap-hkain Burmese, mu ping guo, mu ping kuo in Chinese, elefantæble in Danish, olifantsapple in Dutch, kabeet, and Vakandra in Hindu, and Elefantenapfel in German. It is referred to by various names in

India, including Kayatbael, Kavataleal, Kavita in Bengali, Askotha, Kondhu in Gujarati, Kaitha in Hindi, Kavatha in Marathi, Vilamaram, Vilangai in Tamil, and Kaith in Urdu, among others. One tropical tree is the wood apple [7,8]. With the exception of India's chilly and high-altitude regions, it is one of the hardest fruits to grow in semi-arid and dry environments. When unripe, fruit has an acidic quality, but when ripe, it has a nice flavor [9]. This deciduous tree features oddly pinnate leaves and globose, orange-sized fruits with a hard, woody rind. It belongs to the orange subfamily Citratae of the Rutaceae family, which also contains the genera *Feroniella*, *Aegle*, *Chaetospermum*, *Balsamocitrus*, and *Aeglopsis*. It is a common hard-shelled citrus fruit. Wood apple trees can be effectively grown in regions with mean annual temperatures of 20 to 35 degrees Celsius and mean annual rainfall of 25 to 60 centimeters. The wood apple is typically grown from seeds, which germinate 7–14 days after sowing. However, seedlings need at least 8–10 years to develop fruit, thus grafting and budding are used in commercial propagation to ensure precociousness. Planting trees is typically done during the rainy season, but if irrigation is available, it is best to plant them in February or March because that is when they grow the fastest. For maximum yield, grafted plants should be spaced 8 m by 5 m [10]. Wood apples can be kept year-round for human use as processed foods like jam, chutney, and jelly. They are a cheap, highly nutrient-dense, easily perishable, and seasonally available fruit [11]. It is widely valued for its medicinal qualities in Ayurveda, where it is supposed to heal liver problems, diarrhea, dysentery, respiratory problems, and piles. With its remarkable ability to satisfy thirst, wood apple juice has a lot of potential to grow into a significant crop in the beverage production sector.



Fig No. 1. Wood Apple (Feronia Limonia)

Wood apples are typically consumed raw in Indian homes because of their remarkable flavor and perfect ratio of tartness to sweetness. Herbal treatments are made from the bulk of the tree's parts. Diarrhea, dysentery, and dyspepsia can all be treated with the roots. The stem and root bark aqueous formulations are utilized topically to treat Wood fever, jaundice, eczema, urticaria, and malaria. Fruit extract is used to treat dysentery and diarrhea because it may reduce cholesterol [12]. Numerous microorganisms that cause illnesses in humans have Wood apple fruits are resistant to essential oils, which can be utilized to treat ear conditions such as earaches. Wood apple fruit pulp can significantly raise hepatic glycogen, hexokinase, and HDL levels while lowering lipid profiles and hepatic glucose-6-phosphatase levels [13]. *C. albicans*, *A. tumefaciens*, *B. subtilis*, *P. fluorescens*, and *Escherichia coli* are all inhibited by fruit extracts [14]. It exhibits anti-cancer properties as well. In pharmacological tests, the fruit and root showed hypoglycemic and antiamebic qualities. Aegline, Marmesin, Marmin, and Marmelosin (also known as Imperatorin), a significant chemical component of wood apples with anticancer, antibacterial, and anti-inflammatory properties, are among the plant's many alkaloids [15,16,17]. Two compounds that were isolated from seeds and had notable antiulcer effects were luvangetin and pyranocoumarin. The essential oil from the leaf has antifungal qualities. Cattle are fed the leaves, which have an anise-like scent [17].

II. HISTORY

In 1753, Linnaeus gave the wood apple its scientific name, *Schinus limonia*. In 1747, Linnaeus published his *Account of Hermann's Herbarium of Ceylonese Plants*, which included a fairly detailed description. Under *Limonia Acidissima*, there are four citations. The common lime *Citrus aurantifolia* (Christmas.) Swing is most likely one of the two or more species found in Burman's *Thesaurus Zeylanicus*. His own *Flora Zeylanica* is referenced in the second source, which undoubtedly relates to the wood apple. The wood apple or a nearly related species, *Rumphius Herbarium Amboinense*, is mentioned in the third citation. The fourth reference is *Hesperethus acrenulata* (Roxb.), Rheed, and *Hortus Malabaricus*. The next oldest generic name must be used as *Limonia* is therefore invalid. Published by Corrêa in 1800, this is *Feronia*, the name that is currently widely used. *Feronia limonia* (L.) n. comb is the oldest recognized name for the wood apple since Linnaeus initially described it as *Schinus limonia* in 1753. Similar to *Roktgotia* (*Haematocarpus validus*), the wood apple (*Limonia acidissima* L.) is one of Bangladesh's lesser-known fruits [18].

III. ORIGIN AND DISTRIBUTION

The wood apple originated in Sri Lanka and South India. The parched plains of Bangladesh and the Pakistan are home to the wood apple, which is also commonly grown in Ceylon and India. The wood apple is found in west Africa, Indo-China, and the Philippine Island. Additionally Penang Island, Northern Malaysia, and Southeast Asia all grow it. In Sri Lanka and India, it is planted beside roads, on field borders, and occasionally in orchards. In Malaysia and Indonesia, it is grown in parks and communities. The tree can reach a height of 450 meters above mean sea level (MSL) in the western Himalayas and the western Ghats, which are known for their high biodiversity [20]. It seems to be drought-tolerant and thrives on light soils. In India, a well-planted wood apple orchard is a rare occurrence. There is no information known about the acreage and productivity of this fruit crop in India. Typically, scattered trees can be found in homestead gardens, communal land on field bunds and road sides, and woodlands. In addition to being found in jungles, the tree is frequently cultivated as a border plant [22]. The tree may grow up to 450 meters above mean sea level (MSL) in the very biodiverse western Himalayas and western Ghats. It grows well in light soils and appears to be drought-tolerant. A well-planted wood apple orchard is uncommon in India. The productivity and acreage of this fruit crop in India are unknown. Scattered trees are typically found in woodlands, communal land on field bunds and road sides, and homestead gardens. The tree is commonly grown as a border plant in addition to being found in jungles. It is also said to exist in the woods of Chattisgarh, the Vidhyan hills of Uttar Pradesh, and some parts of Jharkhand's Hazaribagh, Palamau, and Chota Nagpur. It is dispersed throughout several districts in Uttar Pradesh, including Hardoi, Sitapur, Ayodhya, Bahraich, Gonda, Barabanki, Lucknow, Kushinagar, Etawah, and Kanpur. The wood apples in the Rajshahi and Gazipur regions are rather huge. The Indian region of Bundelkhand has a high biodiversity for wood apples, although being infrequently used. The agroclimatic conditions of Bundelkhand are quite promising for commercial farming. In Milkipur Tehsil, Faizabad, particularly in Ratapur village, fruits from wood apple trees were collected. Fresh wood apple fruit was sold in the local market in the Hardoi areas of Uttar Pradesh (U.P., India) [23,24].

IV. PHYSICO-CHEMICAL PROPERTIES

Beta carotene, a precursor to vitamin A, is abundant in wood apple pulp. It also contains substantial vitamin content, and it had little Ascorbic acid content amounts. The pulp that has been scooped out can be consumed raw with or without sugar, or it can be mixed with coconut milk and palm sugar syrup and frozen as ice cream. Wood apples are consumed for breakfast in Indonesia after being combined with honey. India uses wood apple pulp to make savory chutneys. Despite all of these advantages, this fruit is largely overlooked in India and is thought to be a minor fruit crop.

However, in the majority of the tribal areas of central India, it is readily available in large quantities and can be easily turned into chocolate. Fruits are used in India and have a strong therapeutic value. In traditional medicine, unripe fruits are used as an astringent remedy for dysentery and diarrhea, and as a liver and cardiac tonic. It works well to cure gum disease, sore throats, and hiccups. The fruit has therapeutic qualities as well, making it one of India's useful medicinal plants. Essential oil derived from wood apple fruits was found to have antimicrobial action and to be effective against twelve human pathogenic microorganisms. The brown pulp has scattered seeds and is resinous, astringent, fragrant, and acidic or sweet. The fruit could be little and acidic or big and sweet. Because wood apples are inexpensive and may cure thirst, they are used as a good supply of juice during the harvest season. The wood apple pulp is used to make a homemade beverage called "sarbat." The full fruit's flesh indigestion, diarrhea, and piles can be effectively treated using a blend of cardamom, honey, and cumin seeds. Additionally, wood apple pulp was utilized to make the powder. The product is dehydrated and has incredibly extended shelf life with no noticeable changes. The nutritive value also rose with the inclusion of ginger and aonla powder. Wood apples were found to have an average chemical composition of 10.67 to 14.33 °Brix for TSS, 1.04 to 4.50% for acidity, and 4.08 to 4.47% for total sugar. The fresh wood apple fruits' nutritional and chemical characteristics 6.3g protein, 15.6g titrable acidity, 2.6 mg/100g vitamin C, 235 mg/100g total phenol, and 1412.55µg/g total antioxidant capacity were found in it. Fruits' perishable nature and seasonality in production necessitate their preservation for year-round human consumption, given their importance as a major contributor to human well-being and as a more affordable and superior source of protective foods [25].




Fig No. 2. Physicochemical Properties

V. PHYTOCHEMICAL AND ANTIOXIDANT PROPERTIES OF WOOD APPLE

Plants are known to have naturally occurring chemical compounds that are physiologically active, protect plants, and improve human health. Since phytochemicals and their byproducts are antioxidants. They might be able to stop chronic illnesses. Sugars, amino acids, proteins, purines, pyrimidines, alkaloids, terpenes, and phenolics are among the phytochemicals. Carotenoids, flavonoids, cinnamic acids, benzoic acids, folic acid, ascorbic acid, and tocopherols are among the antioxidants found in wood apples. The wood apple's edible portions are abundant in phytochemicals and antioxidants, which are the main types of these substances [26,27,28].

Table.1. Some phytochemical and antioxidants contents of wood apple [28,29]

Phytochemical and Antioxidant	Quantity	Unit
Alkaloids	36	g.100g ⁻¹ of dry matter
saponins	0.16	g.100 ⁻¹ of dry matter
Total phenols	35.72	Ug GAE.mg ⁻¹
Flavonoid	35.51	Uggae.mg ⁻¹
Ascorbic acid	0.45	ug. g ⁻¹
Beta Carotene	0.04	ug. g ⁻¹
Riboflavin	0.23	Ug. g ⁻¹

VI. PHARMACOLOGICAL PROPERTIES

A. Anticancer Properties

Wood apple leaf essential oils have cytotoxic and antioxidant properties. shown that 89.19 μ g/ml of essential oil reduces DNA fragmentation in the human cancer cell line MCF-7 (Michigan Cancer Foundation-7). Similarly, ethanolic fruit extracts are evaluated for their antineoplastic activity on human breast cancer cell lines (SRBR3 and MDA-MBA435), with effective dose ED50 values of 56.1 and 30.6 μ g/ml, respectively. The in vitro anti-tumor activity of fruit methanolic extracts at an oral dose of 570 mg/kg body weight on a mouse model of Dalton's Ascitic Lymphoma (DAL) cell revealed that treatment with extract increased the number of nonviable cells in peritoneal exudates and decreased the number of viable cells. This could be due to the extract being absorbed by viable cells, which ultimately led to cell lysis through macrophage activation or the production of cytokines. RBC and hemoglobin content were also restored by the extract [30,31]. Flavonoids may be responsible for the fruits' anticancer properties [32].

B. Antidiabetic Properties

95% ethanolic extracts' ability to prevent diabetes When unripe wood apple fruits were administered at a dose of 250 mg/kg body weight to rats with streptozotocin-induced diabetes, the animals' blood glucose levels were significantly reduced. Alloxan-induced diabetic rats also show hypoglycemic symptoms [33]. Methanolic extract works well to prevent hyperglycemia at 1.75g/kg body weight. A 21-day study using 200 mg/kg of wood apple bark 400 mg of methanolic extract per kilogram of body weight lowers blood sugar by 54.5 and 39%, respectively [34]. Alloxan-induced rats had a comparable effect, which was measured by blood serum levels as well [35]. After drinking fruit juice for ninety days, there was a notable reduction in blood sugar levels while fasting and after meals [36].

C. Antihyperlipedimic Activity

Fruit powder administered at 2.5, 5, and 10 g/kg body weight for 28 days significantly enhances hepatic glycogen, hexokinase, and HDL while lowering the lipid profile and hepatic glucose-6phosphatase. Fibers, flavonoids, polyphenols, saponins, phytosterols, and ascorbic acid may be the cause of it [37].

D. Antioxidant Activities

Water, petroleum ether, chloroform, ethyl acetate, and methanol extracts were used to assess the antioxidant activity of wood apple pulp [38]. They found significant antioxidant activity using assays for 2, 2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging, hydroxyl radical scavenging (HRSA), ferric reducing antioxidant power (FRAP), nitric oxide radical (NO) scavenging, Trolox equivalent antioxidant capacity (TEAC), and total antioxidant activity (TAA). Antioxidant activity of petroleum ether, methanolic extract, and chloroform in vitro DPPH was used to determine the wood apple leaf the results of the radical scavenging activity, nitric oxide radical scavenging activity, and hydrogen peroxide scavenging activity technique show that every extract has adequate antioxidant qualities [39,40,41]. Additionally, methanolic extract was successful in combining antioxidant enzymes including catalase, superoxidase, and dismutase [42]. Enzymatic antioxidants such as glutathione, superoxide dismutase (SOD), catalase (CAT), and peroxidase were shown to be more active after 30 days of oral treatment of ethanolic leaf extract at 200 mg/kg body weight [43].

E. Hepatoprotective Activities

The hepatoprotective effect of pulp methanolic extracts was assessed against rats' liver damage caused by carbon tetrachloride (CCl₄) [32]. They discovered that rats treated with 100, 200, and 400 mg/kg i.p. for three days had higher levels of hepatic enzymes, particularly aspartate transaminase (AST), alkaline phosphatase (AST), total protein (TP), total bilirubin (TB), and gamma glutamyltransferase (GGT), and lower levels of superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx), and glutathione (GSH). Another study found that aqueous leaf extract increases the levels of alkaline phosphatase (ALP), aspartate aminotransferase (AST), and alanine transaminase (ALT) in the serum. Additionally, compared to the normal control, it raises the levels of urea, creatinine, potassium, and sodium in the phenyl hydrazine-induced anemic non-treated rats [44].

F. Diuretic activity

Urine production has been significantly increased by the leaves' methanolic extract. The extraction method determines the degree of urination and electrolyte excretion, specifically sodium, potassium, and chloride ions. It was discovered that Bath Sonicator extraction (BSE) is less active than Microwave Assisted Extraction (MAE) [45].

G. Antiulcer and Wound Healing Properties

Wood apple fruit pulp has been shown to be effective against rats' stomach ulcers caused by indomethacin. By lowering the concentration of stomach HCl, it prevents gastric ulcers at 500 mg/kg, by raising the pH within the stomach [46]. At 400 mg/kg of the extract, the woundbreaking strength was tightened, the epithelization period was shortened, the wound contraction was raised, and the granulation tissue weight and hydroxyproline concentration increased [36]. By reducing mucosal lesions in a dose-dependent manner, acidissima extracts considerably shield the stomach mucosa from damage caused by ethanol. At 400 mg/kg leaf extract, the fruit's phenolic chemicals provide protection against gastric wall ulcers and leucocyte infiltration in submucosal layers [47].

H. Analgesic Activities

Analgesic activity against writhing mice generated by acetic acid was reported to be 60.53% on methanol, 59.65% on fruit peel acetone extracts, and 78.07% on the common medication Diclofenac Na [48].

I. Neuroprotective Activities

An evaluation of wood apple's neuroprotective properties revealed that it reduces ischemia at 250 mg/kg and 500 mg/kg body weight. Brain damage caused by reperfusion in rats [49].

J. Spermatotoxic Activities

The wood apple fruit's antispermatogenic properties by administering ethanolic extracts at 250 and 500 mg/kg for 55 days, pulp in adult male rats was found to be the cause of the decrease in sperm count, motility, and viability. Additionally, they decreased testicular protein content by 29.86% and increased the fraction of aberrant sperm by 24.58%, respectively [50].

K. Antidiarrheal activity

Wood apple aqueous bark extract was shown to have significant antidiarrheal and gastrointestinal motility-reducing effect by lowering average feces weight and GI motility [51]. The data was similar By Thomas technique on diarrhea brought on by castor oil Peel extracts containing 500 mg/kg of methanol and acetone and found inhibitions of 47.13% and 44.83% [48].

L. Antimicrobial Activity

At 250 mg/kg, the methanol and acetone fruit peels extract exhibits moderate effectiveness, inhibiting *Klebsiella oxytoca*, *Vibrio metschnikovii*, *Escherichia coli*, *Bacillus subtilis*, and *Staphylococcus aureus* by 34.45% and 35.63%, respectively [48]. Z-anethole (22.1%), E-anethole (8.1%), Methyl chavicol (12.0%), and -pinene (28.4%) are among the essential oils found in wood apple leaves. It has antibacterial properties against eight Gram-negative bacteria (*Escherichia coli*, *Klebsiella pneumonia*, *Serratiamarcescens*, *Proteus mirabilis*, *P. vulgaris*, *Pseudomonas aeruginosa*, *Salmonella Typhimurium*, *Enterobacter aerogenes*) and five Gram-positive bacteria (*Staphylococcus aureus*, *Micrococcus flavus*, *M. luteus*, *Bacillus subtilis*, and *Candida albicans*).

Additionally, the oils have minimal inhibitory concentration values of 0.31, 0.52, 0.20, and 0.26 Mg/ml against *Micrococcus luteus*, *Proteus mirabilis*, *Penicillium chrysogenum*, and *Aspergillus niger*, respectively [52]. Agar well was also used to evaluate antibacterial activity. Diffusion technique against a gram negative bacterium (*Proteus mirabilis*) and three gram positive bacteria (*Staphylococcus aureus*, *Bacillus subtilis*, and *Staphylococcus epidermidis*). The plant's various sections contain essential oils that have antimicrobial properties [53]. MIC is the minimum inhibitory concentration on leaf extract in methanol, hexane, and chloroform and demonstrate adequate activity [54]. *Staphylococcus aureus*, *Bacillus cereus*, *Enterococcus faecalis*, *Escherichia coli*, *Salmonella Typhi*, and *Pseudomonas aeruginosa* were tested against biosynthesized silver and zinc oxide nanoparticles, and the results showed 2021 44 effectiveness [55]. Additionally, antimicrobial activity against *Salmonella typhi*, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Klebsiella pneumonia* was assessed using the well diffusion method [56]. Another study indicated that the highest zone of inhibition for *S. aureus*, *S. typhi*, and *P. aeruginosa* was 15.16, 15.5, and 13.33 mm, respectively, when silver and zinc oxide nanoparticles were applied at 400 µg/ml. Even so, zinc oxide nanoparticles have less activity than silver nanoparticles [57]. At 12.5 µg/ml, these nanoparticles are also efficient against *M. tuberculosis* [58]. In Thailand, the wood apple has long been used to cure throat and mouth infections. assesses the antibacterial activity of 500 mg/ml of dried pulp and rind against *Bacillus subtilis*, *Proteus mirabilis*, *Staphylococcus aureus*, and *Staphylococcus epidermidis*. Other researchers have also reported that wood apples have antimicrobial properties [59].

M. Antifungal Activity

Wood apple fruit shells contain antifungal substances as osthonol, xanthotoxin, psoralene, and 2, 6-dimethoxy benzoquinone [30]. Using cup plate techniques, the antifungal activity of leaf ethanolic extracts was tested on *M. gypseum*, *T. tonsurans*, *T. mentagrophytes*, *C. albicans*, and least *T. rubrum* in a dose-dependent manner [60].

N. Larvicidal Activities

With a 90% death rate at 3% aqueous extract, wood apple leaf is effective against *Culex quinquefasciatus* larvae. Additionally, the methanol and chloroform extract exhibits 95% mortality at a concentration of 100 ppm [61]. The presence of terpene in wood apple leaves, which inhibits *Aedes aegypti* eggs, larvae, and pupae, was discovered to have mosquitocidal activity. The larvicidal and pupicidal properties of the extracts are excellent. *L. acidissima* hexane extract exhibits ovicidal action against *Cx* eggs at concentrations of 79.2% and 60% at 500 ppm. *Ae.* and *Quinquefasciatus* [62,63].

VII. PRECAUTION, SIDE EFFECTS AND DISADVANTAGES

Although wood apple have many nutritional and therapeutic benefits, eating them should be done with caution as they may have negative effects. Some of these are listed i.e. Constipation, bloating flatulence, upset stomach, and other digestive disorders can result from consuming too much wood apple juice. Pregnant women and nursing mothers should generally avoid it because it may have negative health effects. Those who have digestive issue should stay away from it. Some people may develop allergies as a result, which needs to be addressed. Shouldn't be consumed on a daily basis. Without interruption since it can interfere with the intestinal peristaltic motions. Tannis found in leaves have long term carcinogenic effects and can causes abortions in pregnant women. It could significantly reduce blood sugar level. If someone on hypoglycemic medication consumers large amounts of it. A person with a thyroid condition should stay away from it [64].

VIII. MARKETING

Kaith fruit have significant market demand, especially during summer. They sell for 5-10 per fruit, and a mature, significant-yielding tree can generate 1,500-2,000 annually. Some inventive kaith growers earn more than 10,000 per plant.

REFERENCES

- [1] Thakur N, Chugh V, Dwivedi S. Wood Apple: An underutilized miracle fruit of India, *Pharm. Innov.* 2020; 9:198-202.
- [2] Chavan SS, Borse T. Study on preparation of wood apple jam enriched in Aleo Vera and Jaggery. *International Journal of Science and Research.* 2022;11(9).
- [3] Sridhar MY, Nataraja KH, Naik N, Satish D, Dileep Kumar A, Masuthi, Karadiguddi M. Identification of elite wood apple Genotypes for physical characters in Selected districts of Karnataka, India. *Int. J. Curr. Micro. Applied Sci.*, ISSN: 2319-7706. 2019;8(9):19992004.
- [4] Jayakumar A, Geetha K. Wood Apple: Uses and Benefits; 2012.
- [5] Khan MA, Singh K, Patel KK, Siddiqui M. Some physical properties of Wood Apple (*Feronia Limonia* L.) Recent Advancement In Food Science and Nutrition Research. 2019;2(1):79-86.
- [6] Nath V, Kumar D, Pandey V. Fruits for Future: Well versed arid and semi arid Fruits. 2008;1.
- [7] Lim TK. "Edible medicinal and non-Medicinal plants, Fruits, vol. 4." ZLibrary. 2012, bok.cc/book/5859961/fc985e. Accessed 23 Dec. 2021; 2021.
- [8] Anonymous. The Ayurvedic Pharmacopoeia of India. Ministry of family Welfare, Department of ISM&H, New Delhi; 1999.
- [9] Das BC, Das SN. Cultivation of minor Fruits. Kalyani publishers, New Delhi. 2003;107111.
- [10] Yadav V. Arid horticulture for enhancing Productivity and economic empowerment. *Ind. Horti.* 2018;74-77.
- [11] Vidhya R, Narain A. Development of Preserved products using under exploited Fruit, wood apple (*Limonia acidissima*). *American J. Food Tech.* 2011;6(4):279-288.
- [12] Mani A, Satish V, Kuchi, Mitra S, Kumar F, Bauri, Das S. Identification and Conservation of elite wood (*Feronia Limonia* L.) genotypes from West Bengal. Society of Education, India. *Advan. Biores. Adv. Biores.* 2020;11(2):75-80.
- [13] Parvez G, Sarker RK. Pharmacological Potential of wood apple (*Limonia Acidissima*): A Review *IJMFM and AP.* 2021;7(2):40-47.
- [14] Ko SNN, Naing HH. Nutritional Compositions, antioxidant and Antimicrobial activities of exotic fruit *Limonia acidissima* , 3Rd Myanmar Korea Conference Research Journal. 2020;3(3).
- [15] Sujitha S, Venkata Lakshmi P. Insights into The In vitro antioxidant, anti-inflammatory And anticancer activities of *Limonia Acidissima* fruits. 2021;11(3):1-11
- [16] Pitre S, Shrivastav SK. Pharmacological, Microbiological and phytochemical studies on roots of *Aegle marmelos*. *Fitoterapi* 1987;58:194-197.
- [17] Yadav K, Singh N. In vitro propagation and Biochemical analysis of field established Wood apple (*Aegle marmelos* L.). *Analele Universității din Oradea – Fascicula Biologie Tom.* 2011;15(1):23-28.
- [18] Bhatt DK, Jha A. A study of incorporation of therapeutic values of wood apple (*Feronia limonia* Swingle) in fruit bar. *Int. J. Pharm. Sci. Res.* 2015;6(10):4398-4405.
- [19] Dowarah B, Deb P, Rana G, Sushmitha K. Fruit Physico-chemical studies of some Local wood apple (*Limonia acidissima* L.) Genotypes . *Int. J. Minor Fruits, Med. Aroma. Plants.* 2021;7(1):53-56.
- [20] Kirtikar KR, Basu BD, An ICS. *Indian Medicinal Plants Vol I* (Singh B and Singh M P Eds.), India. 1993;496-498.

- [21] Orwa C, Mutua A, Kindt R, Jamnadass R, Anthony S. *Feronia limonia* (L.) Swingle Rutaceae. Agroforestry Database: A tree Reference and selection guide version 4.0; 2009.
- [22] Veeraraghavathatham DM, Jawaharlal SJ, Rabindra K. Scientific Fruit Culture. In: Underutilized and underexploited Horticultural crops; K V Peter (ed.), NewDelhi Publishing Agency, New Delhi: 314; 1996.
- [23] Pandey AK, Pal AK, Shukla PK, Yadav MP. Germplasm evaluation of wood apple (*Feronia limonia* L.). *Prog. Hort.* 2013;45(1):76-79.
- [24] Kumar P, Prakash O, Srivastava AK, Thakur N, Chugh V, Singh RS. Correlation Studies in Wood Apple (*Feronia limonia* L.) in Bundelkhand region of Uttar Pradesh, Ind. *J. Pure App.Biosci.* 2021;9(1):316-321.
- [25] International Journal of Food and Ferment Technology Citation: *Int. J. Food Ferment. Technol.*: 11(1&2): 43-49, June & December 2021.
- [26] Krishnaiah D., Sarbatly R., Bano A. Phytochemical Antioxidants for health and medicine – A move Towards nature. *Biotechnology and Molecular Biology Reviews*, 2007, 1(4): 97104.
- [27] Saxena M., Saxena J., Nema R., Singh D., Gupta A. Phytochemistry of medicinal plants.
- [28] *Journal of Pharmacognosy and Phytochemistry*, 2013, 1(6):168-182
- [29] Pandey, S. Satpathy, G. Gupta, R.K. Evaluation of nutritional, phytochemical, antioxidant and antibacterial activity of exotic fruit "*Limonia acidissima*". *Journal of Pharmacognosy and Phytochemistry*, 2014, 3(2):81-88
- [30] Lamani S., Anu-Appaiah K.A., Murthy H.N., Dewir Y.H. Analysis of free sugar, organic acids, and fatty Acids of wood apple (*Limonia acidissima* L.) fruit Pulp. *Horticulture*, 2022, 8(67): 1-13.
- [31] Pradhan, D., Tripathy, G. and Patanaik, S. 2012 Anticancer Activity of *Limonia acidissima* Linn (Rutaceae) Fruit extracts on human Breast cancer cell lines. *Trop. J. Pharm. Res.*,11(3). Doi: 10.4314/tjpr. v11i3.
- [32] Eluru, J.R., Taranalli, A.D. and Kawatra, S. 2015. Anti-tumour activity of *Limonia acidissima* L. methanolic extract in Mice Model of Dalton's Ascitic Lymphoma. *Int. J. Pharmacogn. Phytochem.*,7(6):1094-1100.
- [33] Ilango, K. and Chitra, V. 2009. Hepatoprotective and antioxidant activities of fruit pulp of *Limonia acidissima* Linn. *Int J Health Res.*, 2:361-367.
- [34] Gupta, R., Johri, S. and Saxena, A.M. 2009. Effect of ethanolic extract of *Feronia elephantum* Correa fruits on blood glucose levels in normal and streptozotocin-induced diabeticrats. *Nat Prod Rad.*, 8:32-36.
- [35] Mishra, A. and Garg, G.P. 2011. Antidiabetic activity of fruit pulp of *Feronia elephantum* *Corr. Phcog J.*, 3(20):27-32
- [36] Mohana-Priya, E., Gothandam, K.M. and Karthikeyan, S. 2012. Antidiabetic activity Of *Feronia limonia* and *Artocarpus Heterophyllus* in streptozotocin induced Diabetic rats. *Am. J. Food Technol.*,7:43-49.
- [37] Ilango, K. and Chitra, V. 2010. Wound healing and anti-oxidant activities of the fruit pulp of *Limonia acidissima* Linn (Rutaceae) in rats. *Trop. J. Pharm. Res.*, 9:223-230.
- [38] Rupal, A., Amaravadi, V.R.L. and Narasimhacharya. 2013. *Limonia* fruit as a food supplement to regulate fluoride-induced hyperglycaemia and hyperlipidaemia. *J. Sci. Food Agric.*, 93(2):422-426. doi: 10.1002/jsfa.5762
- [39] Priya-Darsini, D.T., Maheshu, V., Vishnupriya, M., Nishaa. S. and Sasikumar, J.M. 2013. Antioxidant potential and amino acid analysis of underutilized tropical fruit *Limonia Acidissima* L. *Free RadicAntioxid.*, 3:62-69.
- [40] Attarde. 2016. Attarde, D.L, Chaudhari, B.J. and Bhambar, R.S. 2011. Phytochemical investigation and in vitro antioxidant activity of extracts from leaves of *Limonia acidissima* linn. *J Pharm Res.*, 4(3): 766-768.
- [41] Kerkar, S.P., Patil, S., Arya, S., Dabade, A. and Sonawane, S. 2020. *Limonia acidissima*: versatile and nutritional fruit of India. *Int. J. Fruit Sci.*, S405-S413.
- [42] Singhanian, N., Kajla, P., Bishnoi, S., Barmanray, A. and R. 2020. Development and storage studies of wood apple (*Limonia acidissima*) chutney. *Int. J. Chem. Stud.*, 8(1):24732476. doi: 10.22271/chemi.2020.v8.i1a1.8639.
- [43] Patel, D., Kumar, R., Laloo, D. and Hemalatha, S. 2012. Diabetes mellitus: an overview on its Pharmacological aspects and reported Medicinal plants having antidiabetic activity. *Asian Pac. J. Trop. Biomed.*, 25: 411-420.
- [44] Balamuruganvelu, S., Abilash, S.C., Shree Lakshmedevi, S., Geethavani, B., Premlal KR., Jaikumar, S. and Sengottuvelu, S. 2015. AntioxidantActivity of *Limonia acidissima* in high fat diet induced hyperlipidemic rats. *Int. J. Phytopharm.* 6(4):181-183.
- [45] Anacletus, F.C., Nwauche, K.T. and Ugwu, G.M. 2019. Hepatoprotective and nephroprotective potentials of aqueous leaves extract of *Limonia acidissima* in phenylhydrazine- induced anaemic wistar Rats. *Asian J. Pharma. Res. Develop.*, 7(1):27-33.
- [46] Parial, S., Jain, D.C. and Joshi, S.B. 2009. Diuretic activity of the extracts of *Limonia acidissima* in rats. *Rasayan J. Chem.*, 2:53-56.
- [47] Mishra, A., Arora, S., Gupta, R., Manvi, P.R.K. and Sharma, A.K. 2009. Effect of *Feronia Elephantum* (Corr) fruit pulp extract on Indomethacin-induced gastric ulcer in albino Rats. *Trop. J. Pharm. Res.*, 8:509-514.
- [48] Aneesha, A., Rao, R.N., Tejaswini, S.N., Durga, L.S.A., Haseena, S.K. and Maneesha, B. 2018. Phytochemical studies and anti-ulcer activity of *Limonia acidissima* linn. leaf in treating ethanol induced ulcer Albino rats. *Indian J. Res. Pharm. Biotec.*, 6(3):104-110.
- [49] Islam, F., Azad, A.K., Faysal, M., Azad, M.A.K., Islam, S., Amin, M.A. Nahida Sultana, Farhana Yeasmen Dola, Md. Mominur Rahman and Md. Zamshed Alam Begh. 2020. A Comparative Study of Analgesic, Antidiarrhoeal and Antimicrobial Activities of Methanol and Acetone Extracts of Fruits Peels of *Limonia acidissima* L. (Rutaceae). *J. drug deliv. ther.*, 10(1-s):62-65. doi.org/10.22270/jddt.v10i1-s.3882
- [50] Rakhunde, P.B., Saher, S. and Ali, S.A. 2014. Neuroprotective effect of *Feronia limonia* on ischemia reperfusion induced brain injury in rats. *Indian J. Pharm.*, 46:617-621.
- [51] Dhanapal, R., Ratna, J.V., Sarathchandran, I. and Gupta, M. 2012. Reversible antispermatic and antisteroidogenic activities of *Feronia limonia* fruit pulp in adult male rats. *Asian Pac. J. Trop. Biomed.*, 2:684-690.
- [52] Senthilkumar, K.L., Kumawat, B.K. and Rajkumar, M. 2010. Antidiarrhoeal activity of bark extracts of *Limonia acidissima* Linn. *Res. J. Pharm. Biol. Chem. Sci.*, 1
- [53] Joshi, R.K., Badakar, V.M., Kholkute, S.D. and Khatib, N. 2011. Chemical composition and antimicrobial activity of the essential oil of the leaves of *Feronia elephantum* (Rutaceae) from Northwest Karnataka. *Nat. Prod. Commun.*, 6:141-143.
- [54] Pandey, S., Satpathy, G. and Gupta, R.K. 2014. Evaluation of nutritional, phytochemical, antioxidant and antibacterial activity of exotic fruit *Limonia acidissima*. *J. Pharmacogn. Phytochem.*, 3(2):81-88.



- [56] Bagul, V., Dhabekar, S.S., Sansarode, D. and Dandekar, S. 2019. Wood Apple (*Limonia Acidissima* L.): A multipurpose herb in cosmetics. *Int. J. Sci. Dev. & Res.*, 4(7):172181.
- [57] Patil, B. and Taranath, T. 2018. *Limonia acidissima* L. leaf mediated synthesis of silver and zinc oxide nanoparticles and their antibacterial activities. *Microb. Pathog.*, 115: 227-232. doi: 10.1016/j.micpath.2017.12.035.
- [58] Sonawane, S., Bhagwat, A., and Arya, S. 2018. *Limonia acidissima* and *Citrullus lanatus* fruit seeds: Antimicrobial, thermal, structural, functional and protein identification study. *Food Biosci.*, 26: 8-14. doi: 10.1016/j.fbio.2018.09.001.
- [59] Bheemanagouda, N., Patil, T.C. and Taranath. 2018. *Limonia acidissima* L. leaf mediated synthesis of silver and zinc oxide nanoparticles and their antibacterial activities. *Microb. Pathog.*, 115:227-232. doi: 10.1016/j.micpath.2017.12.035
- [60] Bheemanagouda, N., Patil, T.C. and Taranath. 2016. *Limonia acidissima* L. leaf mediated synthesis of zinc oxide nanoparticles: A potent tool against *Mycobacterium tuberculosis*. *Int. J. Mycobacteriol.*, 5(2):197- 204. doi: 10.1016/j.ijmyco.2016.03.004.
- [61] Lairungruang, K., Itharat, A. and Panthong, S. 2014. Antimicrobial activity of extracts from a Thai traditional remedy called Kabpi for oral and throat infection and its plant components. *J. Med. Assoc. Thai.*, 97(8):108-115.
- [62] Shivakumar, P.S. and Vidyasagar, G.M. 2015. Antidermatophytic activity of ethanolic leaves extract of *Limonia acidissima* Groff. *Int. Lett. Nat. Sci.*, 39: 56-62.
- [63] Banerjee, S., Singha, S., Laskar, S. and Chandra, G. 2011. Efficacy of *Limonia acidissima* L (Rutaceae) leaf extract on larval immature of *Culex quinquefasciatus* Say 1823. *Asian Pac. J. Trop. Med.*, 4:711-716.
- [64] Reagan, A.D., Gandhi, M.R., Paulraj, M.G., Balakrishna, K. and Ignacimuthu, S. 2014. Effect of niloticin, a protolimonoind isolated From *Limonia acidissima* L. (Rutaceae) on the immature stages of dengue vector *Aedes aegypti* L. (Diptera: Culicidae). *Acta. Trop.* 139:6, 7-76.
- [65] Reagan, A.D., Gandhi, M.R., Paulraj, M.G. and Ignacimuthu, S. 2015. Ovicidal and Oviposition deterrent activities of medicinal Plant extracts against *Aedes aegypti* L. and *Culex quinquefasciatus* Say Mosquitoes (Diptera: Culicidae). *Osong Public Health Res. Perspect.*, 6(1):64-69. Doi: 10.1016/J.phrp.2014.08.009.
- [66] Sonal R. Zanwar, Amarjeet N. Satwase and Monika G. Phalphale. Wood Apple: An exploited Nutritional Fruit. *Just Agriculture*. Vol.3 Issue- 9, May 2023.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)