



# IJRASET

International Journal For Research in  
Applied Science and Engineering Technology



---

# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume:** 14    **Issue:** IV    **Month of publication:** April 2026

**DOI:** <https://doi.org/10.22214/ijraset.2026.80787>

[www.ijraset.com](http://www.ijraset.com)

Call:  08813907089

E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)

# A Comprehensive Review on the Oxidation of Inorganic Compounds by Cerium (IV): Kinetic and Mechanistic Perspectives

Dhanraj

Assistant Professor Department of Chemistry, Government College Kota, (Raj) University of Kota, Kota

**Abstract :** The oxidation studies with cerium(IV) is important because cerium(IV) is a significant single electron oxidant in aqueous acidic medium. This review paper investigated the kinetic and mechanistic perspective of oxidation of some inorganic compounds by cerium(IV) in aqueous medium. The present article highlights the different type of reactive species of cerium(IV) in various oxidation studies. The nature of these reactive species depends on the concentration and nature of medium used in the reaction. The order of reaction were identified with the respect of substrate, oxidants and metal ions i.e. catalyst.

**Keywords:** Oxidation. Inorganic Compounds, Cerium(IV), Kinetic, Mechanistic.

## I. INTRODUCTION

The kinetics and mechanism of oxidation reactions from the kinetic data have been always the most intriguing subjects in chemistry. The kinetics of the oxidation of inorganic and organic compounds is highly useful in determining the different factors that influence the rate of reaction and in understanding the mechanism of chemical reactions. Oxidation of organic and inorganic compounds to give specific products by certain selective catalyst under certain conditions has received much attention from a chemical as well as an industrial point of view. The oxidation of organic compounds by Ce(IV) in general seems to proceed via the formation of an intermediate complex[1-5]. Among the inorganic substrates which serve as ligands for cerium (IV) are Cl<sup>-</sup>, Br<sup>-</sup> and hypophosphite [6]. Kinetics of oxidation of organic and inorganic compounds by a variety of oxidants such as chloramine-T[7], H<sub>2</sub>O<sub>2</sub>[8], potassium permanganate[9] and O<sub>3</sub>[10], quinolinium dichromate[11], pyridiniumchlorochromate[12], sodium N-bromobenzenesulphonamide[13], imidazolium dichromate[14], Cetyltrimethylammonium Dichromate[15], Benzimidazolium Fluorochromate[16], benzyltrimethylammoniumchlorobromate[17], iodine[18], chlorite[19], hypochlorous acid[20], ferrate (VI) and ferrate (V)[21], hypochlorite and hypochlorous acid[22], HCl[23], peroxymonosulfate[24], N-bromosuccinimide[25], Cerium (IV)[26], peroxymonosulphate[27] in acidic and basic media has been reported in literature. Out of these cerium(IV) and peroxymonosulphate oxidants are of particular importance due to their eco-friendly nature and their capability to oxidize a variety of organic and inorganic compounds catalyzed by different transition metal catalysts[28-31].

### Cerium (IV) as an oxidant

The accessibility of the 4f<sup>0</sup> tetravalent oxidation state for molecular cerium complexes in solution is unique among the lanthanide elements. The thermodynamic basis for the isolation of molecular cerium(IV) compounds originates from the enhanced stability afforded by the [Xe] noble gas electronic configuration for the ion [32]. The existence of molecular cerium(IV) compounds and the ability to cycle between the +3 and +4 oxidation states plays a key role in the separation of cerium from lanthanide-containing ore, and more generally in studies of valence control for nuclear fuel reprocessing [33-35]. Cerium is the only rare earth element with a stable +4 oxidation state in addition to +3 states. The coloured Ce (IV) and the colourless Ce(III) in solution form a favourable one electron redox couple in aqueous acidic medium [36]. The redox potential of the Ce<sup>4+</sup>/Ce<sup>3+</sup> couple depends on the concentration and type of acid media used.

Cerium(IV) is a well-known oxidant in acid media [37] (the reduction potential [38] of the cerium(IV)/cerium(III) couple is 1.70 V) and is stable only in high acid concentrations.

In sulfuric acid and sulfate media cerium(IV) forms several sulfate complexes [39]. Similarly in perchloric acid solutions [40-41] cerium(IV) exists as Ce<sup>4+</sup>, [Ce(OH)]<sup>3+</sup>, [Ce(OH<sub>2</sub>)]<sup>2+</sup>, [Ce-O-Ce]<sup>6+</sup> and [HO-Ce-O-CeOH]<sup>4+</sup>. A potentiometric study [42] of cerium(IV) hydrolysis indicates [Ce(OH)]<sup>3+</sup> to be the predominant active species of cerium(IV), in the concentration range 0.30–2.0 mol dm<sup>-3</sup> HClO<sub>4</sub>, but its role has not received much attention so far.


Because of high stability, both kinetically and thermodynamically [43, 44] and easy availability, ceric sulfate has been widely used in synthetic [45] and mechanistic studies [46-50] in aqueous H<sub>2</sub>SO<sub>4</sub> media.

Actually, Ce(IV) is not so stable in aqueous HClO<sub>4</sub> and HNO<sub>3</sub> media [43, 44]. During the studies on kinetic and mechanistic aspects of Ce(IV) oxidation of different types of both organic and inorganic substrates in aqueous H<sub>2</sub>SO<sub>4</sub> media, the major problem arises from the standpoint of identification of kinetically active Ce(IV) species [46]. Ce(IV) can exist as a mixture of different types of sulfate species such as Ce(SO<sub>4</sub>)<sup>2+</sup>, Ce(SO<sub>4</sub>)<sub>2</sub>, Ce(SO<sub>4</sub>)<sub>3</sub><sup>2-</sup> and HCe(SO<sub>4</sub>)<sub>3</sub><sup>-</sup> in aqueous H<sub>2</sub>SO<sub>4</sub> media. Generally, from the bisulfate dependence of rate at fixed [H<sup>+</sup>], attempts [51-52] have been made to identify the kinetically active Ce(IV) species. Because of so many interdependent equilibria involving Ce(IV), SO<sub>4</sub><sup>2-</sup>, HSO<sub>4</sub><sup>-</sup>, and [H<sup>+</sup>], e.g. Ce(IV)-HSO<sub>4</sub><sup>-</sup>-H<sup>+</sup>; SO<sub>4</sub><sup>2-</sup>-HSO<sub>4</sub><sup>-</sup>-H<sup>+</sup>, it is very difficult [53] to compute the actual concentration of [HSO<sub>4</sub><sup>-</sup>] and [H<sup>+</sup>] under the experimental conditions. In fact, so many approximations are required to compute [HSO<sub>4</sub><sup>-</sup>] and [H<sup>+</sup>] and it leads to the uncertainty to identify the kinetically active Ce(IV) species from the dependence of rate on [HSO<sub>4</sub><sup>-</sup>] and [H<sup>+</sup>]. In fact, several conflicting results are presented in the literature. Hintz and Johnson [54] have reported the reactivity order, Ce(SO<sub>4</sub>)<sup>2+</sup> > Ce(SO<sub>4</sub>)<sub>2</sub> > Ce(SO<sub>4</sub>)<sub>3</sub><sup>2-</sup> in the oxidation of cyclic alcohols. In the oxidation of reducing sugars by Ce(IV) in aqueous sulfuric acid media, there is also a conflict among the workers [55,56] regarding the kinetically active Ce(IV) species.


In sulphuric acid and sulphate media Ce(IV) forms several sulphate complexes, but their role has not received much attention so far. The slow cerium(IV) oxidation reaction of many inorganic compounds are facilitated by a minute amount of metal ions in aqueous sulphuric acid medium. [57]. In this process the redox potential of different couples involved in the oxidation reaction in acidic medium make the metal ions catalysis for oxidation of inorganic compounds by cerium (IV) appear feasible.

## II. REACTIVE SPECIES OF CERIUM (IV)


According to Hardwick and Robertson [58], Cerium(IV) exists in various forms in sulphuric acid solution (2 M) and constant ionic strength, the following equilibrium exists.





The value of equilibrium constants are  $Q_3 = 3.5 \times 10^3$  (25°C),  $Q_2 = 2 \times 10^2$  (25°C) and  $Q_3 = 0.6$  (at=20°C). Without keeping the ionic strength constant, Buganko and Kaun Lin [59] investigated the nature of cerium(IV) spectrophotometrically in 0.1 to 1.76 M sulphuric acid. The equilibrium at 20±2°C is shown below.



They have suggested that Ce(SO<sub>4</sub>)<sub>2</sub> and HCe(SO<sub>4</sub>)<sub>3</sub><sup>-</sup> are the predominant species up to 2M and H<sub>3</sub>Ce(SO<sub>4</sub>)<sub>4</sub> exists above 2M H<sub>2</sub>SO<sub>4</sub>. Hargreaves and Sutcliffe have [60] suggested in addition the following equilibrium to explain mechanism, of certain reactions.



But the equilibrium constants were not calculated, though they may be useful in some mechanisms. In addition to the above equilibrium, the following were also proposed [61]



ZhaiYoung quin et al.[62]and Bai W. S. and Coworkers[63] have reported  $\text{Ce}(\text{SO}_4)_2$  as the kinetically active species of cerium(IV) in sulphuric acid medium in the oxidation of ethanol amine and ethylene glycol respectively. On the other hand Tripathy, S. N., Prasad, R. K. [64] and Kushwaha, U., & Singh, A. K.[65]have suggested reactive species as  $\text{Ce}(\text{SO}_4)_3^{2-}$ . While S. K. Mishra and Y. K. Gupta[66] have shown that  $\text{CeSO}_4^{2+}$  is the active species of cerium(IV), Hinz, H. L., and Johnsonr[67]also suggested  $\text{CeSO}_4^{2+}$  as the reactive species in the oxidation study of cyclic alcohol by cerium(IV) in mixed perchloric and sulphuric acid medium.

### III. OXIDATION STUDIES OF INORGANIC COMPOUNDS WITH CERIUM (IV)

To understand the mechanistic aspects of reduction of Ce(IV) to Ce(III), several kinetic studies of Ce(IV) sulphateoxidation of different types of organic substrates have been carried out by different workers [68-70].A number of inorganic ions are oxidised by Ce(IV), but none of them serves as ligand to Ce +4 and it was to investigate this possibility that hypophosphite was employed for the reduction of Ce(IV). Bernhart[71] has reported that hypophosphite can be quantitatively oxidised by Ce(IV).

The Kinetic study of oxidation of antimony (III) by cerium(IV) in presence of micro amounts of iodide and chloride in an aqueous sulphuric acid medium have been studied spectrophotometrically at 25°C and  $I=3.0\text{mol dm}^{-3}$ [72]. The stoichiometry of the reaction is 1:2, which shows that one mole of antimony (III) requires two moles of cerium(IV). In the presence of chloride the order of reaction was unity with respect to cerium(IV) and halide, whereas in the presence of iodide the order with respect to cerium(IV) was unity and with respect to iodide was more than one. The order with respect to antimony (III) is less than unity in presence of both halide ions. Increased sulphuric acid concentration increased the rate of reaction. The order with respect to  $[\text{H}^+]$  was less than unity. The active species of cerium (IV) was understood to be  $\text{H}_3\text{Ce}(\text{SO}_4)_4^-$ , whereas in the presence of chloride the active species of reductant was  $\text{SbCl}_3$  and in the case of iodide the reactive species was  $\text{SbI}_2$ .

The oxidation of antimony(III) by cerium(IV) in the presence of bromide in sulphuric acid medium has been studied[73]. The active species of cerium(IV) is identified as  $\text{H}_3\text{Ce}(\text{SO}_4)_4^-$ , whereas that of reluctant as  $\text{SbBr}_2^+$ .

The oxidation of antimony(III) by cerium(IV) has been studied spectrophotometrically in the presence of a minute amount of Mn(II) in aqueous sulphuric acid medium[74]. The stoichiometry of the reaction showed that one mole of antimony(III) required two mole of cerium(IV). The reaction was first order with respect to Ce(IV) and Mn(II). The order with respect to antimony (III) was less than unity. The rate of reaction decreased with increased concentration of sulphuric acid. The added product sulphate and bisulphate decreases the rate. The active species of Ce(IV), Sb(III) and Mn(II) was  $\text{Ce}(\text{SO}_4)_2$ ,  $[\text{Sb}(\text{OH})(\text{HSO}_4)]^+$  and  $[\text{Mn}(\text{H}_2\text{O})_4]^{2+}$  respectively.

The kinetics of oxidation of antimony(III) by cerium(IV) in perchloric acid medium has been reported[75]. The varying of  $\text{H}^+$  concentration the rate first increases, reaches maximum at  $[\text{H}^+] = 0.15 \text{ M}$  then decreases as the concentration of hydrogen ion increases. the active species of oxidant and reductant in perchloric acid medium are assumed to be the hydrolytic species of Ce(IV) and  $\text{SbO}^+$  based on the effect of  $\text{H}^+$  concentration and spectrophotometric measurements.

The oxidation of arsenic by cerium (IV) catalysed by chromium(III) has been studied in aqueous sulphuric acid medium[76]. The rate of reaction was increase with increasing the concentration of sulphuric acid in both catalysed and uncatalysed condition. The active species of cerium(IV) is understood to be the  $\text{H}_3\text{Ce}(\text{SO}_4)_4^-$ .

S. M. Tuwar and coworkers[77] has been reported the oxidation of mercury(I)catalysed by Pd(II) in aqueousperchloric acid medium. The stoichiometry resultsshowed that one mole of Ce(IV) required one mole of Hg(I). The active species of oxidant and catalyst are found to be  $\text{HCe}(\text{SO}_4)_3^-$  and  $\text{PdCl}^+$ respectively.

The Kinetic study of oxidation of selenium(IV) by cerium(IV) in perchloric acid medium has been reported by Dikshituluand coworkers[78]. They observed that the added product cerium(III) retards the reaction. In the presence of cerium(III), the order with respect to cerium(IV) found to be between one and two, while in the absence of cerium(III) the order with respect to cerium(IV) was first order and with respect to selenium(IV) was fractional order.

Ru(III) catalysed oxidation of selenium(IV) by Ce(IV) in perchloric acid medium also reported by Dikshitulu and coworkers[79]. In the another study of oxidation of tellurium (IV) by cerium(IV) in perchloric[80], nitric and sulphuric acid medium[81], The added product cerium(III) retard the reaction while other product tellurium(VI) accelerated the reaction. The order of reaction with respect to cerium(IV) was first order and with respect totellerium (IV) was fractional.

K. A. Thabajat. al[82] reported the oxidation of palladium(II) by cerium(IV) in perchloric acid medium at 40°C. The reaction was reported to be first order each in Ce(IV) and Pd(II). The order with respect to  $[H^+]$  was fractional. The active species of oxidant and reluctant are respectively  $H_3Ce(SO_4)_4$  and  $[PdCl_4]$ .

S. A. Chimatdar and coworkers[83] studied the Ru(III) catalyzed oxidation of thallium(I) by cerium(IV) in aqueous sulphuric acid medium. The reaction was reported to be first order in cerium(IV) as well as in Ru(III) and the order in thallium(I) varies from zero to unity. Increase in acid concentration accelerates the reaction rate. The added products cerium(III) and thallium (III) has a negligible effect on the reaction rate.

B. P. Sinha and H. P. Mathur reported[84] the Os(VIII) catalysed oxidation thallosion by ceric ion in sulfuric acid medium. The order with respect to both ceric ion and thallosion ion noticed unity. The rate of reaction directly proportional to the concentration of catalyst. The  $HSO_4^-$  inhibits the reaction and the rate of reaction varies inversely as the square of bisulphate ion concentration. Thallic and cerious ion does not have any effect on the rate. Another reaction of oxidation of Ti(I) by cerium(IV) in presence of Os(VIII) catalyst in perchloric acid medium reported by B. P. Sinha and H. P. Mathur[85]. In this case the order of the reaction is found to be a function of both Ti(I) and Ce(IV). Hydrogen ion inhibits the reaction rate.

Oxidation of hypophosphorous acid by cerium(IV) in aqueous nitric[86] medium has been reported. The kinetic studies of oxidation of Fe(II)[87], chloride[88], bromide[89], hypophosphite[90], bromous acid[91], Cr(III)[92], As(III)[93] (in presence of iodide) by cerium(IV) have been reported.

#### IV. CONCLUSION

Cerium (IV) in redox reactions reacts as an outer sphere or inner sphere mechanism in agreement with the type of the reluctant. This review study has correlated the different factors to control the mechanistic paths of catalysis in the title oxidation reactions. Several kinetically active cerium(IV) species like  $Ce(SO_4)_2$ ,  $CeSO_4^{2+}$ ,  $Ce(SO_4)_3^{2-}$ ,  $H_3Ce(SO_4)_4$  were involved in the oxidation reaction of different type of inorganic and organic compounds. The effect of substrates, oxidants, metal ions (as catalyst) and added products on the rate of reaction have been reported and observed that reaction rate varies with varying the concentration of these reagents.

#### REFERENCES

- [1] Young, L. B., & Trahanovsky, W. S. (1969). Oxidation of organic compounds with cerium (IV). VII. Formation constants for 1: 1 cerium (IV)-alcohol complexes. *Journal of the American Chemical Society*, 91(18), 5060-5068.
- [2] Trahanovsky, W. S., & Macaulay, D. B. (1973). Oxidation of organic compounds with cerium (IV). XVII. Relative rates of formation of allyl, benzyl, and tert-butyl radicals by oxidative cleavage of alcohols. *The Journal of Organic Chemistry*, 38(8), 1497-1499.
- [3] Voskresenskaya, O. O., & Skorik, N. A. (2015). Thermodynamics of the formation of intermediate complexes in the oxidation of oxalic acid by cerium (IV) and kinetics of their intramolecular redox decomposition. *Russian Journal of Physical Chemistry A*, 89, 1821-1827.
- [4] Duke, F. R., & Forist, A. A. (1949). The Theory and Kinetics of Specific Oxidation. III. The Cerate--2, 3-Butanediol Reaction in Nitric Acid Solution. *Journal of the American Chemical Society*, 71(8), 2790-2792.
- [5] Duke, F. R., & Bremer, R. F. (1951). The Theory and Kinetics of Specific Oxidation. IV. The Cerate 2, 3-Butanediol Reactions in Perchlorate Solutions. *Journal of the American Chemical Society*, 73(11), 5179-5181.
- [6] Carroll, R. L., & Thomas, L. B. (1966). The kinetics and mechanism of the oxidation of hypophosphorous acid by cerium (IV) in perchloric acid solution. *Journal of the American Chemical Society*, 88(7), 1376-1381.
- [7] Pareek, D., Rolaniya, A., Bhasin, M., & Sailani, R. (2023). Ruthenium (III) catalyzed oxidation of paracetamol by chloramine-T in aqueous alkaline medium—A kinetic and mechanistic pathway. *International Journal of Chemical Kinetics*, 55(1), 39-48.
- [8] Bowers, A. R., Cho, S. H., & Singh, A. (2021). Chemical oxidation of aromatic compounds: comparison of  $H_2O_2$ ,  $KMnO_4$  and  $O_3$  for toxicity reduction and improvements in biodegradability. In *Chemical Oxidation* (pp. 11-25). CRC Press.
- [9] KS, R., Shastry, V. A., & Kumar, M. (2023). Comparative kinetic study of palladium (II) catalyzed and uncatalyzed oxidation of ambroxol hydrochloride with potassium permanganate in an aqueous alkaline medium: A mechanistic approach. *Physical Chemistry Research*, 11(4), 865-875.
- [10] Wang, S., Du, L., Zhu, J., Tsona, N. T., Liu, S., Wang, Y., & Wang, W. (2018). Gas-phase oxidation of allyl acetate by  $O_3$ , OH, Cl, and  $NO_3$ : reaction kinetics and mechanism. *The Journal of Physical Chemistry A*, 122(6), 1600-1611.
- [11] Khan, A. A. P., Mohd, A., Bano, S., & Siddiqi, K. S. (2011). Kinetics and mechanism of deamination and decarboxylation of 2-aminopentanedioic acid by quinolinium dichromate (QDC) in aqueous perchloric acid medium. *Industrial & engineering chemistry research*, 50(17), 9883-9889.
- [12] M Heravi, M., Fazeli, A., & Faghihi, Z. (2016). Recent advances in application of pyridiniumchlorochromate (PCC) in organic synthesis. *Current Organic Synthesis*, 13(2), 220-254.
- [13] Vaz, P. N. (2023). Kinetics of oxidation of acidic amino acids by sodium N-bromobenzenesulphonamide in acid medium: A mechanistic approach. *Journal of Chemical Sciences*, 135.
- [14] Rao, A., Singh, N., Mahawar, D. K., & Mahawar, P. (2022). Kinetics and mechanism of the oxidation of cysteine by imidazolium dichromate. *Journal of the Indian Chemical Society*, 99(4), 100386.

- [15] Garnayak, S., & Patel, S. (2016). Oxidation of Antitubercular Drug Isoniazid by a Lipopathic Oxidant, Cetyltrimethylammonium Dichromate: A Mechanistic Study. *International Journal of Chemical Kinetics*, 48(1), 32-44.
- [16] Arora, B., Ojha, J., & Mishra, P. (2021). Kinetics and Mechanism of Oxidation of Aliphatic Secondary Alcohols by BenzimidazoliumFluorochromate in Dimethyl Sulphoxide solvent. *Oriental Journal of Chemistry*, 37(3), 626.
- [17] A Tiwari, V., Kumbhani, S., Shastri, I., & Sharma, V. (2008). Kinetics and mechanism of oxidation of some thioacids by benzyltrimethylammoniumchlorobromate.
- [18] Xu, L., Valkai, L., Kuznetsova, A. A., Makarov, S. V., & Horváth, A. K. (2017). Kinetics and mechanism of the oxidation of thiourea dioxide by iodine in a slightly acidic medium. *Inorganic Chemistry*, 56(8), 4679-4687.
- [19] Chigwada, T., Mbiya, W., Chipiso, K., & Simoyi, R. H. (2014). S-oxygenation of thiocarbamides V: oxidation of tetramethylthiourea by chlorite in slightly acidic media. *The Journal of Physical Chemistry A*, 118(31), 5903-5914.
- [20] Ueki, H., Chapman, G., & Ashby, M. T. (2010). Reactive sulfur species: Kinetics and mechanism of the oxidation of aryl sulfonates with hypochlorous acid. *The Journal of Physical Chemistry A*, 114(4), 1670-1676.
- [21] Sharma, V. K. (2010). Oxidation of inorganic compounds by ferrate (VI) and ferrate (V): one-electron and two-electron transfer steps. *Environmental science & technology*, 44(13), 5148-5152.
- [22] Kumar, K., & Margerum, D. W. (1987). Kinetics and mechanism of general-acid-assisted oxidation of bromide by hypochlorite and hypochlorous acid. *Inorganic Chemistry*, 26(16), 2706-2711.
- [23] Zhang, S., Zhang, Q., Díaz-Somoano, M., Dang, J., Xu, Y., Zhao, Y., & Zhang, J. (2022). A kinetic study on mercury oxidation by HCl over typical Mn-based SCR catalysts. *Fuel*, 329, 125421.
- [24] Li, J., Jiang, J., Zhou, Y., Pang, S. Y., Gao, Y., Jiang, C., & Guan, C. (2017). Kinetics of oxidation of iodide (I<sup>-</sup>) and hypiodous acid (HOI) by peroxymonosulfate (PMS) and formation of iodinated products in the PMS/I<sup>-</sup>/NOM system. *Environmental Science & Technology Letters*, 4(2), 76-82.
- [25] Eljack, N. D., & Sulfab, Y. (2011). Kinetics and mechanism of oxidation of aquaethylenediaminetetraacetatocobaltate (II) by N-bromosuccinimide in aqueous acidic solution. *Transition Metal Chemistry*, 36, 87-91.
- [26] Thriveni, M. K., Shivalingaswamy, T., Bellappa, S., & Mallikarjuna, I. (2022). Ruthenium (III) Catalyzed Oxidation of Methenamine by Cerium (IV) in Aqueous Sulfuric Acid Medium—A Kinetic and Mechanistic Approach. *Russian Journal of Physical Chemistry A*, 96(13), 2878-2886.
- [27] He, X., & O'Shea, K. E. (2020). Selective oxidation of H1-antihistamines by unactivated peroxymonosulfate (PMS): Influence of inorganic anions and organic compounds. *Water Research*, 186, 116401.
- [28] Tandon, P. K., Dwivedi, P. B., Singh, S. B., & Yadav, S. C. (2013). Kinetic study of iridium (III) catalyzed oxidation of 2-phenyl ethanol and 2-methyl cyclohexanol by cerium (IV) sulphate. *Journal of the Indian Chemical Society*, 90(12), 2237-2245.
- [29] Chimatadar, S. A., Nandibewoor, S. T., Sambrani, M. I., & Raju, J. R. (1987). Chromium (III)-catalysed cerium (IV) oxidation of arsenic (III) in aqueous sulphuric acid. *Journal of the Chemical Society, Dalton Transactions*, (3), 573-577.
- [30] Lavanya, C., Andal, P., & Ramachandran, M. S. (2015). Co (II) catalyzed oxidation of organic compounds by peroxomonosulphate.
- [31] Gupta, B., Gupta, K. S., & Gupta, Y. K. (1984). Kinetics and mechanism of the silver (I)-catalyzed oxidations of arsenic (III) and antimony (III) with peroxodiphosphate in acetate buffers. *Journal of the Chemical Society, Dalton Transactions*, (9), 1873-1876.
- [32] Cotton, F. A., Wilkinson, G., Murillo, C. A., & Bochmann, M. (1999). *Advanced inorganic chemistry*. John Wiley & Sons.
- [33] Ellis, R. J., & Antonio, M. R. (2012). Redox Chemistry of Third Phases Formed in the Cerium/Nitric Acid/Malonamide-n-Dodecane Solvent Extraction System. *ChemPlusChem*, 77(1), 41-47.
- [34] Wei, Y., Kumagai, M., Takashima, Y., Asou, M., Namba, T., Suzuki, K., ... & Ohe, S. (1998). The Application of an Advanced Ion Exchange Process to Reprocessing Spent Nuclear Fuels, (I) Separation Behavior of Fission Products from Uranium. *Journal of nuclear science and technology*, 35(5), 357-364.
- [35] Chiang, M. H., Williams, C. W., Soderholm, L., & Antonio, M. R. (2003). Coordination of actinide ions in Wells- Dawson heteropolyoxoanion complexes. *European Journal of Inorganic Chemistry*, 2003(14), 2663-2669.
- [36] Binnemans, K. (2006). Applications of tetravalent cerium compounds. *Handbook on the physics and chemistry of rare earths*, 36, 281-392.
- [37] Thabaj, K. A., Munavalli, D. S., Chimatadar, S. A., & Nandibewoor, S. T. (2007). Chloride and iodide mediated oxidation of antimony (III) by cerium (IV) in aqueous sulphuric acid medium. *Transition Metal Chemistry*, 32(7), 902-912.
- [38] Day M.C. and Selbin J., *Theoretical Inorganic chemistry*, Reinhold Pub.Corp., New York, 226 (1964)
- [39] Kharzeoua, S. E., & Serebrennikou, V. V. (1967). Study of sulphato complexes of cerium (IV) by infrared absorption spectra. *Russian Journal of Inorganic Chemistry*, 12, 1601-1605.
- [40] Hardwick, T. J., & Robertson, E. (1951). Ionic species in ceric perchlorate solutions. *Canadian Journal of Chemistry*, 29(10), 818-827.
- [41] Datt, N., Nagori, R. R., & Mehrotra, R. N. (1986). Kinetics and mechanisms of oxidations by metal ions. Part VI. Oxidation of  $\alpha$ -hydroxy acids by cerium (IV) in aqueous nitric acid. *Canadian journal of chemistry*, 64(1), 19-23.
- [42] Baker, F. B., Newton, T. W., & Kahn, M. (1960). The kinetics of the reaction between uranium (IV) and cerium (IV). *The Journal of Physical Chemistry*, 64(1), 109-112.
- [43] Grant, D. (1964). Thermal instability of cerium (IV) sulphuric acid solutions. *Journal of Inorganic and Nuclear Chemistry*, 26(2), 337-346.
- [44] Grant, D., & Payne, D. S. (1961). The effect of heat on solutions of cerium (iv) sulphate in sulphuric acid and some analytical implications. *Analytica Chimica Acta*, 25(1), 422-428.
- [45] Jwo, J. J., & Noyes, R. M. (1975). Oscillations in chemical systems. IX. Reactions of cerium (IV) with malonic acid and its derivatives. *Journal of the American Chemical Society*, 97(19), 5422-5431.
- [46] Das, A. K. (2001). Kinetic and mechanistic aspects of metal ion catalysis in cerium (IV) oxidation. *Coordination Chemistry Reviews*, 213(1), 307-325.
- [47] Das, A. K., & Das, M. (1995). Kinetics and mechanism of ruthenium (III) catalyzed oxidation of ethanol by cerium (IV) in aqueous sulfuric acid media. *International Journal of Chemical Kinetics*, 27(1), 7-16.
- [48] Sharma, J., & Singh, R. (1990). Kinetics of Oxidation Of Maltose And Lactose By Cerium (IV) In Sulfuric-Acid-Solution. *Journal Of The Indian Chemical Society*, 67(4), 316-317.
- [49] Lakshmi, S., & Renganathan, R. (1996). Kinetics of oxidation of certain pyrimidines by Ce (IV). *International journal of chemical kinetics*, 28(10), 713-720.

- [50] Mishra, S. K., & Gupta, Y. K. (1967). Kinetics and mechanism of oxidation of hypophosphite by Ce (IV) in sulphuric acid solution. *Journal of Inorganic and Nuclear Chemistry*, 29(7), 1643-1656.
- [51] Mishra, S. K., & Gupta, Y. K. (1970). Stoichiometry and kinetics of the reaction between cerium (IV) and hydrazine in acid solution. *Journal of the Chemical Society A: Inorganic, Physical, Theoretical*, 2918-2921.
- [52] Sankhla, P. S., & Mehrotra, R. N. (1972). Kinetics of cerium (IV) oxidation of formaldehyde in aqueous sulphuric acid solution. *Journal of Inorganic and Nuclear Chemistry*, 34(12), 3781-3788.
- [53] Das, A. K., Mondal, S. K., Kar, D., & Das, M. (1999). Kinetics and Mechanism of Chromium (III) Catalysed Oxidation of Propan-1-ol and Butan-1-ol by Cerium (IV) in Aqueous Sulphuric acid Media. *BioInorganic Reaction Mechanisms*, 1(2), 169-176.
- [54] Hinz, H. L., & Johnson, D. C. (1967). The mechanism of oxidation of cyclic alcohols. *J Org Chem*, 32, 556-564.
- [55] Mehrotra, R. N., & Amis, E. S. (1974). Kinetics of oxidation of aldo sugars by cerium (IV) in aqueous sulphuric acid. *The Journal of Organic Chemistry*, 39(12), 1788-1791.
- [56] Ali, M. S., & Khan, Z. (2006). Effect of surfactant micelles on the kinetics of oxidation of D-fructose by cerium (IV) in sulphuric acid medium. *International Journal of Chemical Kinetics*, 38(1), 18-25.
- [57] Chimatadar, S. A., Basavaraj, T., & Nandibewoor, S. T. (2002). Ruthenium (III) Mediated Cerium (IV) Oxidation of Thallium (I) in Aqueous Sulphuric Acid--A Kinetic and Mechanistic Study. *Inorganic Reaction Mechanisms*, 4(3-4), 209-219.
- [58] Hardwick, T. J., & Robertson, E. (1951). Association of ceric ions with sulphate (a spectral study). *Canadian Journal of Chemistry*, 29(10), 828-837.
- [59] Bugaenko, L. T.; Huaf, kuan Lin, Russ. *J. Inorg. Chem.* 8, 299 (1963).
- [60] Hargreaves, G., & Sutcliffe, L. H. (1955). The oxidation of aqueous formaldehyde by ceric ions. *Transactions of the Faraday Society*, 51, 1105-1111.
- [61] Suresh C. Yadav, Mahesh K. Gupta, Shalini Srivastava, Praveen K. Tandon (2023). Kinetic and mechanistic aspects of cerium(IV) as an oxidant in ruthenium catalysed oxidation reaction: Amini review, *Green Chemistry & Technology Letterse*, Vol 9, No 2, 2023, pp 01-06.
- [62] Yong-qing, Z., Hong-mei, L., Lin, Y., Guo-zhong, Y., Wen-yu, S., & Yu-kai, L. (2007). Kinetics and mechanism of iridium (III)-catalyzed oxidation of ethanol amine by cerium (IV) in sulphuric acid media. *Chemical Research in Chinese Universities*, 23(3), 333-338.
- [63] Bai, W. S., Zhai, Y. Q., Liu, H. M., Lü, H. Y., & Song, W. Y. (2008). Kinetics and mechanism of iridium (III) catalyzed oxidation of ethylene glycol by cerium (IV) in sulphuric acid media.
- [64] Tripathy, S. N., & Prasad, R. K. (1980). Kinetics of Oxidation of Citric Acid by Cerium (IV).
- [65] Kushwaha, U., & Singh, A. K. (2010). Kinetics and Mechanism of Oxidation of Citraconic Acid By Cerium (IV) in Acidic Medium Catalysed by Ruthenium (III). *Reviews in Inorganic Chemistry*, 30(3), 159-174.
- [66] Mishra, S. K., & Gupta, Y. K. (1968). Kinetics and mechanism of the oxidation of Sb (III) by Ce (IV) in sulphuric acid solution in presence of chloride ion. *Journal of Inorganic and Nuclear Chemistry*, 30(11), 2991-3008.
- [67] Hinz, H. L., & Johnson, D. C. (1967). The mechanism of oxidation of cyclic alcohols. *J Org Chem*, 32, 556-564.
- [68] K. C. Tiwari, and B. L. Krishna, *ChemSoc*, 1961, 3097.
- [69] S. A. Mishra, and P. S. RadhaKrishnamurti, *Ind. J. Chem. Sec. A*, 20 : (8), 297-301 (Eng) (1981).
- [70] Iftikar Ahmed and E.L..Fatin, *Arab Gulf, J. Sei. Res.* 2 (2), 435- 443 (Eng) (1984).
- [71] D. N. Bernhart, *Analyt. Chem.* 26, 1798 (1954).
- [72] Thabaj, K. A., Munavalli, D. S., Chimatadar, S. A., & Nandibewoor, S. T. (2007). Chloride and iodide mediated oxidation of antimony (III) by cerium (IV) in aqueous sulphuric acid medium. *Transition Metal Chemistry*, 32(7), 902-912.
- [73] Munavalli, D. S., Thabaj, K. A., Chimatadar, S. A., & Nandibewoor, S. T. (2007). Bromide mediated oxidation of antimony (III) by cerium (IV) in aqueous sulphuric acid medium
- [74] Patil, R. K., Chimatadar, S. A., & Nandibewoor, S. T. (2008). Mechanistic study of cerium (IV) oxidation of antimony (III) in aqueous sulphuric acid in the presence of micro amounts of manganese (II) by stopped flow technique. *Transition Metal Chemistry*, 33, 625-633.
- [75] Mishra, S. K., & Gupta, Y. K. (1970). Kinetics of oxidation of antimony (III) by cerium (IV) in media containing perchloric acid. *Journal of the Chemical Society A: Inorganic, Physical, Theoretical*, 260-264.
- [76] Chimatadar, S. A., Nandibewoor, S. T., Sambrani, M. I., & Raju, J. R. (1987). Chromium (III)-catalysed cerium (IV) oxidation of arsenic (III) in aqueous sulphuric acid. *Journal of the Chemical Society, Dalton Transactions*, (3), 573-577.
- [77] Tuwar, S. M., Nandibewoor, S. T., & Raju, J. R. (1990). Palladium (II) catalysed cerium (IV) oxidation of mercury (I) in aqueous perchloric acid
- [78] Dikshitulu, L. S. A., Vani, P., & Rao, V. H. (1981). Kinetics and mechanism of oxidation of Se (IV) by Ce (IV). *Journal of Inorganic and Nuclear Chemistry*, 43(6), 1261-1265.
- [79] Dikshitulu, D. S. A., Vani, P., & Rao, V. H. (1981). Kinetics and mechanism of Ru (III) catalyzed oxidation of Se (IV) by Ce (IV). *Inorganic and Nuclear Chemistry Letters*, 17(5-6), 187-191.
- [80] Dikshitulu, L. S. A., Rao, V. H., & Dindi, S. N. (1980). Kinetics of Oxidation of Tellurium (IV) by Cerium (IV) In Perchloric Acid Medium.
- [81] Dikshitulu, L. S. A., Rao, V. H., & Dindi, S. N. (1981). Kinetics & Mechanism of Oxidation of Tellurium (IV) by Cerium (IV) in Nitric & Sulphuric Acid Media.
- [82] Thabaj, K. A., Chimatadar, S. A., & Nandibewoor, S. T. (2006). Mechanistic study of oxidation of palladium (II) by cerium (IV) in aqueous acid. *Transition metal chemistry*, 31, 186-193.
- [83] Chimatadar, S. A., Basavaraj, T., & Nandibewoor, S. T. (2002). Ruthenium (III) Mediated Cerium (IV) Oxidation of Thallium (I) in Aqueous Sulphuric Acid--A Kinetic and Mechanistic Study. *Inorganic Reaction Mechanisms*, 4(3-4), 209-219.
- [84] Sinha, B. P., & Mathur, H. P. (1971). Kinetics and Mechanism of the Oxidation of Thallous Ion by Ceric Ion: Part VI. *Zeitschrift für Physikalische Chemie*, 246(1), 342-351
- [85] Sinha, B. P., & Mathur, H. P. (1971). Kinetics and mechanism of the oxidation of Tl (I) by Ce (IV) catalysed by Os (VIII) in perchloric acid medium. *Journal of Inorganic and Nuclear Chemistry*, 33(6), 1673-1679.
- [86] Nagori, R. R., Mehta, M., & Mehrotra, R. N. (1979). Kinetics and mechanisms of redox reactions in aqueous solution. Part 5. Oxidation of hypophosphorous acid by cerium (IV) in aqueous nitric acid. *Journal of the Chemical Society, Dalton Transactions*, (2), 216-220.



- [87] Adamson, M. G., Dainton, F. S., & Glentworth, P. (1965). Kinetics of the oxidation of iron (II) by cerium (IV) in aqueous perchloric acid. *Transactions of the Faraday Society*, 61, 689-701.
- [88] Duke, F. R., & Borchers, C. E. (1953). The Kinetics of the Cerium (IV)—Chloride Reaction. The Cerium (IV)—Thallos Reaction in the Presence of Chloride. *Journal of the American Chemical Society*, 75(21), 5186-5188.
- [89] King, E. L., & Pandow, M. L. (1953). The Rate Law and Mechanism for the Reaction of Cerium (IV) and Bromide Ion in Aqueous Sulfate Media 1, 2. *Journal of the American Chemical Society*, 75(13), 3063-3068.
- [90] Mishra, S. K., & Gupta, Y. K. (1967). Kinetics and mechanism of oxidation of hypophosphite by ce (iv) in sulphuric acid solution. *Journal of Inorganic and Nuclear Chemistry*, 29(7), 1643-1656.
- [91] Sullivan, J. C., & Thompson, R. C. (1979). Kinetic study of the cerium (IV)-bromous acid reaction in acid sulfate solution. Implications for the Belousov-Zhabotinskii oscillating reaction. *Inorganic Chemistry*, 18(9), 2375-2379.
- [92] Tong, J. Y. P., & King, E. L. (1960). The Kinetics and Mechanism of the Reaction of Cerium (IV) and Chromium (III) 1, 2. *Journal of the American Chemical Society*, 82(15), 3805-3809.
- [93] Yates, J. S., & Thomas, H. C. (1956). Kinetics of the Iodide-catalyzed Reaction between Cerium (IV) and Arsenic (III). *Journal of the American Chemical Society*, 78(16), 3950-3953.



10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)