



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 13 Issue: IV Month of publication: April 2025

DOI: <https://doi.org/10.22214/ijraset.2025.68766>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

A Review: Recent Advancement in Herbal Technology

Mr. Sudhanshu Dhananjay Kapase¹, Mr. Nilesh Zanzane², Mr. Nitin Gawai³, Mr. Akash Jayashing Ghadage⁴, Mr. Nikhil Kailas Tambe⁵

B. Pharmacy Department, Mahadev Kanchan College of Pharmaceutical Education and Research, Uruli Kanchan, Pune, Maharashtra, India

Abstract: The fashion ability of herbal products has also led to the emergence of certain dangerous ideas and particulars that have caught directors and consumers off guard and some of which have had mischievous effects. Herbal specifics have gained fashion ability lately because of their numerous advantages. Herbal remedies have proven to be extensively effective as treatments for a number of illnesses. Even if the maturity of these operation is unusual, over 80 percent of the world's population takes herbal treatments. One of the biggest challenges facing scientists is the creation of unique logical styles that can directly describe the phytochemical arrangement, including quantitative evaluations of marker/ bioactive mixes and other important elements. Both contemporary developments and pictorial convectional ways are described in the current review essay. Recent advancements includes DNA characteristic, metabolomics system, differential vibration polarography,- shaft diffraction etc. are practical. Capillary electrophoresis and chromatographic styles contributions towards standardization of herbal drugs is also proved.

Keyword: herbal, ways, chromatography, Authentication, Isolation

I. INTRODUCTION

The creative use of contemporary technologies and scientific principles to ameliorate the product, birth, expression, and distribution of herbal products is known as advanced herbal technology. These technologies combine state- of- the- art exploration in accoutrements wisdom, biotechnology, and pharmacology with traditional knowledge of medicinal shops. The ideal is to save the natural integrity of herbal- based products while enhancing their efficacy, safety, and thickness. This strategy incorporates slice-edge tactics including factory inheritable engineering, enhanced birth ways like supercritical fluid birth(SFE), and drug delivery systems grounded on nanotechnology. The need for herbal remedies has grown encyclopedically in recent times, leading to notable developments in herbal technology. High- performance liquid chromatography(HPLC) is one fashion that's being utilised to guarantee the energy and chastity of factory factors. The creative use of contemporary technologies and scientific principles to ameliorate the product, birth, expression, and distribution of herbal products is known as advanced herbal technology. These technologies combine state- of- the- art exploration in accoutrements wisdom, biotechnology, and pharmacology with traditional knowledge of medicinal shops. The ideal is to save the natural integrity of herbal- based products while enhancing their efficacy, safety, and thickness. This strategy incorporates slice- edge tactics including factory inheritable engineering, enhanced birth ways like supercritical fluid birth(SFE), and drug delivery systems grounded on nanotechnology. The need for herbal remedies has grown encyclopedically in recent times, leading to notable developments in herbal technology. currently, styles like high performance liquid chromatography(HPLC) are employed to guarantee the energy and chastity of factory ingredients, while screening bioinformatics and phytochemicals^[1]

II. IDEAL⁽³⁾

- 1) Formalized birth styles Development of standardized styles for rooting active composites from sauces.
- 2) Identification of bioactive composites concentrate on relating and understanding the bioactive composites present in sauces.
- 3) expression of effective delivery systems Creating innovative delivery systems to insure the effectiveness of herbal drugs.
- 4) Full eventuality Application The overarching thing is to harness the complete eventuality of herbal drugs.
- 5) Chemical structure diversity Herbal drugs parade diversity in their chemical structures.
- 6) Biodiversity application using the biodiversity of sauces for medicinal purposes.
- 7) Revolutionized Natural product webbing preface of new technologies has revolutionized the webbing of natural products.
- 8) Discovery of new medicines Easing the discovery of new medicines through advanced webbing styles.
- 9) Mortal health benefit The end is to contribute to mortal health by exercising the advancements in herbal technology.

III. PLANT SPECIES IDENTIFICATION WAYS

Identification of seasoning material is combination of art and wisdom. Correct botanical Identification is original and one amongst the foremost vital way for making certain smart quality end Product. If the morning chief is n't original or smart quality; also finished product quality Can not be secured. Identification of condiment will be done by fully different approach though there are Several ways offered, nothing methodology is applicable for all sauces. utmost of the time combination Of these ways will be used for proper identification⁽²⁾

- 1) Approaches grounded on leaves utmost of studies grounded on splint shape or color to prize features, there are some important splint features similar that " aspect rate ", " narrow factor ", " conciseness ", " centroid ", " curiosity ", " dissipation ", " area ", " original periphery ", " moments steady ", " etc. In J.Chaki and R.Parekh presented a system grounded on splint image to identify shops, two ways were used moments- steady and centroid- diameters. N.Valliammal and S.N.Geethalakshmi proposed a mongrel approach grounded on " discrepancy stretching " and " adaptive thresholding " that at the same time adjusts the intensity position of factory splint image. The bracket approach proposed by K. Gurpreet and K. Gurpinder was grounded on splint features " isoperimetric quotient ", " curiosity ", " aspect rate ", " splint area ", " splint border ", " length of the major and minor axes ", " reliability " and " upper and lower triangle area of the splint ". Tree identification system that presented by Itheriet al.detected boundary also described the detected boundary by using the directional scrap histogram, geometric features similar that " rectangularity ", " protuberance and reliability ", " circularity ", " Sphericity " and " cirque friction " are uprooted. It was noted that utmost of the exploration grounded on geometric features of leaves as in(6) in which geometric features were uprooted with hypersphere classifier to classify further than 20 species of shops.^[4]
- 2) Approaches grounded on flowers or fruits Some approaches fete shops grounded on flowers, by fastening on the flower region as a whole or on corridor of the flower. proposed flower shape descriptors " area ", " border ", " roundness ", and " aspect rate ". Some studies prize features like color, texture, and shape presented a system for classifying flower, RGB histogram was used as a point spices of input images were used which had good quality, for bracket arbitrary timber algorithm was used. presented a system for feting the orchid species grounded on image of flower, shape and color features were uprooted, for segmentation minimal similarity grounded on region incorporating was used which was easy to use and more accurate than others, for classificationsupport vector machine system was used.^[4]
- 3) Macroscopy: Macroscopy involves checking external look or sensitive characters Like color odor, taste, size, shape, fracture etc. botanic identification of condiment is generally done By trained person like biologist. For proper botanic identification, entire factory at the side of root And flower is needed. Botanic identification is rested on morphology that involves checking colorful rudiments of sauces like leaves, flower, root et al. Leaves and flowers nearly vital corridor that grease in identification of factory. Herb will be caught on for color size, shape and Arrangement of leaves and flower. Arrangement of leaves on stem AN branching is nominated Phyllotaxy. Differing types of arrangement of leaves like alternate distichous, contrary, Decussate, whorled kinds of leaves arrangement will be useful to spot condiment duly. Different types of shapes of leaves like round, oblong, obovate, globular direct, lanceolate, Elliptical, spatulate, cordate, unsubdividedar one amongst the stylish tool to spot shops. Indeed perimeters Of leaves will be caught on to spot sauces. perimeters like entire, serrated rough, sinuate, ciliate, Spinose grease in identification of condiment. In some cases, fully different species of shops will be known Only once flowering.
- 4) Microscopy: exploration plays terribly pivotal part in identification of medicine those ar Morphologically analogous. Magnifier will be used for checking sections of leaves, root And stem make sure identity of condiment. Research will be also used to check stomata, trichome, Calcium swab chargers, which can be distinctive thereto condiment. Sure splint constants like stomatal Index Palisade quantitative relation, tone islet variety ar vital for proper identification of condiment. Indian Senna and backcountry| Alexandriasenna| Alexandriansenna| truesenna| tinnevellysenna| Indiansenna| Sennaalexandrina| Cassiaacutifolia| Cassiaaugustifolia| senna} will be discerned by exploitation bitsy parameters. Indian Senna has vain islet variety nineteen.5 to 22.5 whereas tinnevellysenna has twenty five to twenty nine.5, Alexandrian senna Have stomatal indicator seventeen to twenty whereas tinnevellysenna have eleven.4 to 13.3. Likewise several indispensable shops will be known by exploration. Form of essence swab is salutary to spot factory for illustration- ensign formed demitasse is gift in Jamestown weed, needle formed demitasse ar Presents in German iris, rap helpers ar gift in squill, monoclinic prism form Is gift in black henbane, beach formed gift in deadly nightshade. Research is Especially helpful just in case of fine drug. bounce grains, size and length of filaments, staining responses like hard vascular towel and bast will be studied by exploitation magnifier^[2]

- 5) DNA characteristic: Use of biotechnological tools for correct identification of sauces is the forthcoming rearmost technology. Molecular labels like RAPD, ISSR, RFLP uses DNA characteristic for identification of condiment at molecular position. Molecular labels are nothing but sequence of DNA which is unique to each factory. First, factory DNA is insulated and also amplified with the help of PCR and also screened for analogous and different patterns. shops also have DNA patterns analogous to mortal. Pattern of this DNA can be linked in the for barcoding or DNA characteristic. DNA barcoding identify factory as specie species position or molecular position, indeed when small part of factory without flower More recent technology like genomics, proteomics, transcriptomics and metabolomics fashion can identify factory at inheritable position.^[2]

IV. BIRTH WAYS

In birth, the alloy of substances is separated, by dissolving each element with one or other cleaners, which yields two phases – Rabbinat Phase (rich in Feed cleaner) and prize Phase (rich in Solute). When the Relative Volatility is 1 the separation of the factors in the admixture isn't possible by Distillation and when relative Volatility is lower than 1 birth system is used for separation of the factors.^[5]

Type based on Types of Phases^[5]

Liquid-Liquid birth – Sample Phase (Liquid)

Excerpt Phase (Liquid)

Base for Separation (Partitioning)

Solid Phase birth or – Sample Phase (Gas, Liquid) Micro-extraction Extract Phase (Liquid, Solid, Stationary Phase) Base for Separation (Partitioning or adsorption)

Filtration or Solid Liquid – Sample Phase (Solid) birth Excerpt Phase (Liquid)

Base for Separation (Partitioning)

Supercritical Fluid- Sample Phase (Solid, Liquid) birth Extract Phase (Supercritical Fluid)

Base for Separation (Partitioning with applied heat)

A. Types of birth

- 1) Liquid birth Liquid – liquid birth, also improperly known as partitioning, is a system to separate composites based on their relative solubilities in two different immiscible liquids, generally water and an organic cleaner. The term cleaner birth can also be used for partitioning of substances between liquid and gas, liquid and solids and gas and solids. Liquid – liquid birth is an introductory fashion in chemical and radio chemical exploration laboratories. The simplest system is a homemade one using a separator channel. This type of process is generally performed after a chemical response, constantly but not always including chemical completion, as part of the work-up of the response assessment.^[6]
- 2) Solid Phase birth -Solid Phase birth is sample Preparation Method used for insulation, enrichment and sanctification of factors from thirsty results depending upon their physical and chemical parcels. This involves reaching of thirsty samples with a solid phase or sorbent, where the emulsion is adsorbed on the face of the solid phase former to elation. It also overcomes issues faced in the Liquid-Liquid birth Operation, similar as phase separation isn't satisfactory, less recovery, waste of large quantities of organic cleaners. Also, the tableware utilised is in liquid birth.^[5]

B. Birth Ways

Ultrasonic supported birth, Microwave oven supported birth, Mechano- chemical supported birth, Supercritical Fluid birth, Heat Influx birth, Pressurized Liquid/ fluid birth, Enzyme supported birth.^[5]

- 1) Microwave oven supported birth Method: Principle Electromagnetic swells correspond of two perpendicularly oscillatory fields videlicet Electric Field and glamorous Field, which can also be called as Microwave oven. Electromagnetic swells are absorbed by the material and converted to heat energy. This is a Microwave Energy. These swells are non-ionisable radiation. There are two mechanisms for conversion of electromagnetic energy to spicy energy or heat Ionic Conduction and Dipole Rotation. As there's change in sign of the electric field, the direction of ions changes which results in collisions of motes with each other and repel the inflow of ions. As the electric field diminishments, restoring of thermal motes takes place that in turn leads to release of thermal energy.^[5]
- 2) Soxhlet Birth: The Soxhlet birth system uses a fragile quantum of detergent and is veritably cost- operative. The detergent is appended to the solvent force beaker and mounted onto a heating mantle. After heating, the condensed vapors of the detergent come in connection with the sample greasepaint, and the answerable portion of the greasepaint gets mixed with the detergent

for birth. When the solvent face exceeds the ultimate height of the siphon, the detergent containing the excerpt is drained ago. The beaker is reiterated, rooting a portion of the substance each time so that the logical substance is constantly exercised as a pure detergent and the uprooted substance is concentrated in the beaker^[7]

C. Pressurized Liquid birth(PLE)

Pressurized fluid birth employs the use of detergents at high temperatures, above its boiling point and below its overcritical point, under enough pressure to conserve them in a fluid country. In this section, the essential criteria for the election of applicable functional parameters will be handed from a theoretical point of prospect. extension- supporter, the abecedarian principles of PLE for logical slices are described. Due to the terminations of marketable outfit, the only potentiality to deal with liquid slices is by transubstantiating them into solids, for illustration, by adding an ab/ adsorbent. The process of rooting analytes from circumfluous and logical slices can be described by the following five way:

- 1) Dampening the sample(analytes to be uprooted and matrix) with birth detergent
- 2) Desorption of composites from the matrix(involving or not the breakdown of chemical bonds)
- 3) Solvation of the composites in the detergent
- 4) Dissipation out of the matrix
- 5) Prolixity through the nearest solvent subcaste around the matrix to eventually reach the bulk detergent thus, birth forcefulness is told by three interrelated aspects matrix sequel, mass transfer, and solubility. These nonidentical features in PLE are limited by nonidentical considerations, similar as the election of inflow rate, pressure, temperature, and time exercised in the birth.

D. Ultrasonic-assisted Extraction

(UAE) is a swift and efficient extraction method that utilizes ultrasound to create quick solvent movement, leading to an increased mass transfer rate and faster extraction. In comparison to other sophisticated extraction methods, UAE is more cost-effective, environmentally friendly, and user-friendly. The assertions were backed by Boateng & Lee, 2013, which state that UAE is an efficient and quick method that uses less energy, time, and materials, resulting in purer products with higher yields. UAE can take place at low operating temperatures, preventing thermal degradation of the extracts and maintaining the properties of the bioactive compounds regarding their structure and molecules, making UAE an excellent choice for the edible oil sector. UAE not only prevents thermal damage to bioactive compounds but also protects plant materials. It can be utilized in two distinct methods: either through an ultrasonic bath or with an ultrasonic horn transducer. Ultrasonic bath that featured a temperature-regulated apparatus which assisted in the recovery of heat-sensitive compounds such as essential oils^[9]

Direct contact with the sample or indirect contact are the two methods of ultrasonic irradiation. Indirect contact, like in an ultrasonic bath system, occurs when contact occurs through the sample's walls. Because it can power up to 100 times better than indirect contact or an ultrasonic bath, the direct contact technology is more effective in the extraction process^[9]

V. METHOD OF PURIFICATION AND ISOLATION^[2]

A. Methods of General Isolation

- 1) Techniques for extraction
- 2) The extraction of plant material might be an essential technique for the extraction and purification of natural plant components.
- 3) Plant matrices are inherently well-developed, with a wide range of chemicals with different physical and
- 4) Chemical characteristics. Therefore, meticulously separating it from the rest of the plant, matrices, and
- 5) In order to assess plants, make pure chemicals of interest.
- 6) Categorized They must be described in this chapter according to the temperatures at which they operate.
- 7) Strategies involving moderate or temperature.
- 8) Cold extraction technique
- 9) Literature has provided representations of the process.

VI. CHROMATOGRAPHIC METHODOLOGY

In short, samples of dried plant parts that have been chopped, crushed, or otherwise processed Chromatographic Methodology: Overview and a Few Terms Chromatography, which translates to "color-writing," is a physical separation procedure that allows a mixture of substances to be separated, isolated, and purified into distinct molecules that rely on varying distribution rates based on

- 1) Solubility
- 2) Affinity (whether molecules are polar or non-polar)

Interaction with fixed material (the stationary phase, which we shall define later). The mixture's components are distributed among two phases: the mobile phase, which moves in a designated direction at different speeds, and the stationary phase and the mobile phase. It is known that Michael Tswett, the Russian botanist in 1901 discovered that chlorophyll pigments are split into different colored components when he uses a column containing CaCO_3 and moves its combination on it. As a result, he is referred to as the father and founder of chromatography. The three primary components of every chromatographic separation method are as follows:

- a) Sample
- b) Phase of mobility
- c) The stationary stage.

The stationary Phase

Which can only be a solid or a liquid, is the solid material at which the component mixture will be separated and isolated. A mobile phase is a solid or liquid material that transports a mixture of samples that need to be separated, isolated, and purified at the stationary phase's surface. Normal phase liquid chromatography (NPLC), which uses a polar stationary phase, is the first of two categories of chromatographic separation procedures. The second is reversed-phase liquid chromatography (RPLC). In contrast, the mobile phase is non-polar. In order to successfully separate using four chromatographic techniques.

We should select the appropriate parameter between the stationary and mobile phases when using chromatographic techniques for separation. Chromatography serves as a bridge between analytical methods that ascertain a sample's chemical composition and concentration and primitive methods that rely on separating and isolating only the mixture sample rather than determining the concentration of the purified sample.^[10]

A. Thin Layer Chromatography

The solid phase in the process of thin layer chromatography is a thin glass plate covered with silica gel or aluminum oxide. The solvent used as the mobile phase is selected based on the characteristics of the mixture's constituent parts. Just above the bottom of the TLC plate, a starting point is treated with a tiny quantity of a compound or mixture. The plate is then developed in the developing chamber, which features a shallow solvent pool that is somewhat lower than the sample application level. As the solvent passes over the mixture, each component will either dissolve in the solvent and go up the plate or stay with the solid phase. The capillary action draws the solvent up through the plate's particles. Especially, starting at a place just above the TLC plate's bottom, a tiny quantity of a compound or mixture is applied.

After the plate has been applied, it is developed in a developing chamber with a shallow pool of solvent somewhat below the sample application level.

Capillary action draws the solvent up through the plate's particles, and when the solvent passes over the mixture, each component either dissolves in the solvent or progresses up the plate or stays within the solid phase. Particularly depending on its physical traits, which are determined by its molecular structure and group function, chemicals could travel up a plate or remain behind. The physical characteristics of each particular molecule, and consequently its molecular structure and functional groups, determine whether the compound advances along the plate or remains behind.^[11]

B. Thin-layer high-performance chromatography (HPTLC)

The upgraded and automated version of thin-layer chromatography (TLC) known as High Performance Thin Layer Chromatography (HPTLC) has improved separation efficiency and detection limits. Other names for it include flat-bed chromatography, planar chromatography, and high pressure thin layer chromatography. For both qualitative and quantitative analytical tasks, this potent analytical technique is equally appropriate. Adsorption, partitioning, or both can cause separation, depending on the type of adsorbents utilized on the plates and the solvent system used for development.

Applications: Various facets of HPTLC foundations: principle, theory, and comprehension; instrumentation: automation, validation, optimization, and implementation; and qualitative and quantitative analysis. There have been reports of hyphenation potential analytical analysis, phytochemical analysis, herbal medication quantification^[12].

C. High-performance Chromatography using Liquids

One type of column chromatography that is frequently used in biochemistry and analysis to separate, identify, and quantify the active substances is high-performance liquid chromatography, also known as high pressure liquid chromatography, or HPLC. A column that contains packing material (the stationary phase), a pump that circulates the mobile phase or phases through the column, and a detector that displays the primary components of HPLC. The stationary phase, the molecules being examined, and the solvent or solvents utilized all affect retention time. Certain chemical or physical interactions with the stationary phase slow down the introduction of the sample to be studied in a tiny amount into the stream of mobile phase. Both the stationary and mobile phase compositions and the type of analyte determine how much retardation occurs. The retention time of a particular analyte is the moment at which it elutes, or exits the end of the column. Miscible mixtures of water or organic liquids (most frequently methanol and acetonitrile) are employed as solvents. Depending on the analyte's affinity for the present mobile phase, the gradient separates the analyte mixtures. The characteristics of the analyte and stationary phase influence the choice of solvents, additives, and gradient.

D. Types of HPLC

HPLC types are often determined by the process's phase system. The HPLC types listed below are frequently employed in analysis [13].

- 1) Normal phase chromatography: This technique, sometimes referred to as Normal phase HPLC (NP-HPLC), separates analytes according to their polarity. The polar analyte and the polar stationary phase interact to lengthen the elution time, and adsorption intensities rise with analyte polarity.
- 2) Chromatography using reversed phase: A non-polar stationary phase and an aqueous, moderately polar mobile phase are features of HPLC (also known as RP-HPLC or RPC). When the analyte molecule associates with the ligand in the aqueous eluent, the contact surface area surrounding the non-polar section of the molecule determines how much the analyte will bind to the stationary phase.
- 3) Size exclusion chromatography (SEC): Also referred to as gel permeation chromatography or gel filtration chromatography, SEC primarily uses size to separate particles. It is also helpful in figuring out the quaternary and tertiary structures of amino acids and proteins. The molecular weight of polysaccharides is frequently determined using this method.
- 4) Excluded are ions with the same charge. Ligand-exchange chromatography, ion-exchange chromatography of proteins, high-pH anion-exchange chromatography of carbohydrates and oligosaccharides, and water purification are all common applications for this type of chromatography.
- 5) Gas chromatography: This method is unique and widely applicable. Its early development involved the analysis of gases and vapors from highly volatile components. Different components of the sample are analyzed using a device known as a gas chromatograph (GC). One common form of chromatography in analytical research is gas chromatography (GC), which is used to separate and investigate substances that can evaporate without dissolving. Finding the individual components of a mix or assessing a substance's purity are common applications for GC. Martin and Synges work enabled gas-liquid chromatography (GLC), and James and Martin's contributions to the field transformed chemical separations and analysis. [14]

Benefits The following are some benefits of gas chromatography [14].

It is a dependable method that offers quick analysis.

- 1) It produces great resolution and is very efficient.
- 2) It uses detectors that are sensitive.
- 3) A tiny sample was needed.
- 4) It is non-destructive because it makes it possible to connect a mass spectrometer, which determines the masses of individual molecules that have been electrically charged and transformed into ions.
- 5) It offers excellent quantitative precision.
- 6) It is a tried-and-true method with a wealth of applications and literature.

The drawback One drawback of gas chromatography is the following [14].

- 1) Only applicable to volatile samples.
- 2) Thermolabile samples (those that break down at high temperatures) are not appropriate for it.
- 3) It works well with preparative chromatography. Because the majority of non-MS detectors are destructive,
- 4) An MS detector is necessary for the structural elucidation of the analyte.

VII. WHO SETS THE STANDARDS FOR HIGH-QUALITY, STANDARDIZED HERBAL FORMULATIONS ^[15]

The bioactive extract should be standardized using the chromatographic fingerprints (TLC, HPTLC, HPLC, and GC) and active principles or main components. Nine The following actions are part of the process of standardizing raw drug materials:

- 1) Authentication (plant parts obtained, regional status, botanical identification such as phytomorphology, microscopical and histological analysis, taxonomical identity, etc.)
- 2) Foreign matter (collected herbs should not contain dirt, bug pieces, animal excrement, etc.) The organoleptic evaluation includes the drug's color, taste, look, odor, and feel, among other sensory characteristics.
- 3) Important tissues for diagnosis found in the medication powder. e. Ash and extractive values.
- 4) amount of moisture and volatile matter
- 5) Identification of heavy metals, such as arsenic, lead, and cadmium.
- 6) Chromatographic and spectroscopic evaluation: TLC, HPTLC, and HPLC techniques will yield semi-quantitative and qualitative data regarding the primary active ingredients in the crude medication. The spectroscopic fingerprint can also be used to evaluate the drug's quality.
- 7) Pesticide residue: Typically found in herbs, pesticides are subject to restrictions specified by the Food and Agricultural Organization (FAO) and WHO. When the herbs are being grown, these insecticides are combined with them. Pesticides such as DDT, BHC,

VIII. MEDICATION FOR DEVELOPING TECHNOLOGY

A. *Jasminum, Or Jasmine*

The limbic system, which is in charge of affecting the nervous system, sends messages to your body when you inhale the chemicals from jasmine. You can use jasmine as an essential oil to add to a diffuser to catch the aroma, or you can keep it in your room as a plant to help your anxiety and depression systems. In addition to reducing anxiety and depression, jasmine can help you focus better, sleep better, balance your hormones, and reduce your risk of infection. This demonstrates how the jasmine plant has several uses and can raise your standard of living. ^[2]

REFERENCES

- [1] Gavhane P, Avhad P, Sanap T, Badwar O. A review on advanced herbal technology. *Int J Pharm Sci*. 2015;2(11):1130-1137.
- [2] Salunke NS, Kazi SM, Bais SK. Review on advance herbal technology. *Int J Adv Res Sci Commun Technol (IJARSCT)*. 2023;3(2).
- [3] Dubey TS, Shaikh SS, Sawale V, Kakade R, Vishe H. Advanced herbal technology for the development of herbal drug molecule - A review. *Int J Compr Adv Pharmacol*. 2024;9(1):7-16.
- [4] Hssoon IM, Kassir SA, Altaie SM. A review of plant species identification techniques. *Int J Sci Res (IJSR)*. 2018;7(8):325-328.
- [5] Patel K, Panchal N, Ingle P. Review of extraction techniques: microwave, ultrasonic, pressurized fluid, Soxhlet extraction, etc. *Int J Adv Res Chem Sci (IJARCS)*. 2019;6(3):6-21. doi:10.20431/2349-0403.0603002.
- [6] Bjørnstad T. Introduction to liquid-liquid extraction. *Lecture Compendium - Nuclear Chemistry*, University of Oslo.
- [7] Hussain G. Soxhlet extraction principal, working & usage [Internet]. ResearchGate; 2023 Jul. Available from:
- [8] Alvarez-Rivera G, Bueno M, Mendiola JA, Ibáñez E. Pressurized liquid extraction. 2020:375-398. doi:10.1016/B978-0-12-816911-7.00013-X.
- [9] Syahir A, Sulaiman S, Mel M, Othman M, Sulaiman SZ. An overview: analysis of ultrasonic-assisted extraction's parameters and its process. *IOP Conf Ser Mater Sci Eng*. 2020; 778:012165. doi:10.1088/1757-899X/778/1/012165.
- [10] Sayed MA. A review of chromatography: principles, classification, applications. doi:10.13140/RG.2.2.22113.43361.
- [11] Kumar S, Jyotirmayee K, Sarangi M. Thin layer chromatography: A tool of biotechnology for isolation of bioactive compounds from medicinal plants. *Int J Pharm Sci Rev Res*. 2013;18(1):126-132.
- [12] Ramu B. High performance thin layer chromatography and its role in the pharmaceutical industry: review. 2018;5(3):29-34.
- [13] Malviya R, Bansal V, Pal OP, Sharma PK. High performance liquid chromatography: A short review. *J Glob Pharma Technol*. 2010;2(5):22-26.
- [14] Patil P, Yele T, Jawale J, Pawar S. Gas chromatography: a review. *Int Res J Mod Eng Technol Sci*. 2022;4(12).
- [15] Umamaheswari D, Muthuraja R, Kumar M, Venkateswarlu BS. Standardization of herbal drugs – An overview. *Int J Pharm Sci Rev Res*. 2021;68(1):213-219.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)