



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 11 Issue: VIII Month of publication: Aug 2023 DOI: https://doi.org/10.22214/ijraset.2023.55245

www.ijraset.com

Call: 🕥 08813907089 🔰 E-mail ID: ijraset@gmail.com



# A Study on the Nutrient Content, Phytochemicals and Antioxidant Potential of Raw, Germinated and Fermented *Moringa Oleifera* Seeds

Shyama Reji<sup>1</sup>, Dr. S. Janaki Alias Priya<sup>2</sup>

Research Scholar<sup>1</sup>, Assistant Professor<sup>2</sup>, Department of Home Science, University of Madras

Abstract: Diet and Nutrition are important factors in the promotion and maintenance of good health throughout the life. A plant-based diet can prevent and treat chronic diseases. Health is "Absence of Disease". Moringa oleifera is rich in macro and micronutrients like protein, carbohydrate, calcium, phosphorus, phosphorus, iron, vitamins, beta carotene and other bioactive compounds which are important for normal functioning of the body, prevention and curing of certain diseases. M. oleifera has excellent therapeutic properties including anticancer, antiulcer, antimicrobial, antioxidant. Various researches have concluded that Moringa should be used as a functional ingredient in food products. This study was taken up to investigate the effects of germination and fermentation on the nutritional, bioactive, and antimicrobial characteristics of moringa seed and it is found to have essential phytochemicals, micro and macro nutrients which can be used as natural remedies to prevent various diseases and deficiencies.

Keywords: Moringa Oleifera Seeds, Germinated, Fermented, Nutrient, Phytochemicals, Antioxidant Potential

# I. INTRODUCTION

Diet and nutrition are important factors in the promotion and maintenance of good health throughout the life. Nowadays, with continuously changing socio-economic status, people have become more concerned about their health [1]. Mankind has always been interested in food. Several studies over the last 50 years have clearly shown that diets dominated plant-based foods such as fruits, vegetables and dietary fibres prevent and reduce the risk of chronic diseases such as cardiovascular diseases, obesity, diabetes, and so on and promote human health [2]. The generation of scientific research linking foods of plant origin and health worldwide has resulted that plant bioactive compounds have antioxidant and other healthy properties [3]. Ayurveda and traditional medicine have utilized natural resources such as plants and trees as part of their regime to treat different illness and diseases with positive outcomes. As an effect of growth in population and agricultural expansion, only a few forests are remaining, and they are highly degraded, causing losses in traditionally important nutritious foods, medicines and other useful products leading to food insecurity [4].

*Moringa oleifera* (Moringaceae) is one tree having enormous nutritional and medicinal benefits [1]. Nearly all parts have shown to be effective against several ailments including cancer which was attributed to the bioactive constituents [5]. *M. oleifera* is commonly known as the drumstick tree or horse radish tree [6]. It is used as a vegetable and also in Indian folk medicine for the treatment of various illnesses. The length of plant can reach 12m with soft and brittle stems. The diameter is about 20-40 cm and lifespan are about 30 years. Moringa leaves are compound, pinnate double, and of small round or oval shape. The fruit, called "drumstick," is long and angular, its sides form a triangle; the drumsticks are about 15-45 cm long [7]. Taller plants bear creamwhite, small size flowers in clusters throughout the season, which subsequently develop into slender dark-green, three-sided, edible seed pods with tapering ends. Each pod contains 20 fleshy pulp and round pea-sized seed encased inside a wing shaped coat (hull).

*M. oleifera*, native to sub-Himalayan Mountain of Northern India grows in the tropical and subtropical areas of the world. *M. oleifera*, the native of sub-Himalayan Mountain of Northern India; is now cultivated for a variety of purposes in the whole tropical and sub-tropical regions of the world [8]. India is one of the large producers of moringa, with an annual production of 1.1 to 1.3 million tons of fruits from an area of 380 km<sup>2</sup> [9]. The cultivation of moringa in India occurs mainly in the southern states of Tamil Nadu, Karnataka, Kerala, and Andhra Pradesh. In India, Andhra Pradesh stands at the first position in both area and production followed by Karnataka and Tamil Nadu. Moringa is commonly found in countries of Southeast Asia, the Philippines, the Middle East, Africa, and the Caribbean.



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 11 Issue VIII Aug 2023- Available at www.ijraset.com

*M. oleifera* is an extremely valued plant, which has an effective range of medicinal uses in traditional folk medicine with high nutritional value. Many pharmacological researches have shown the ability of this plant to exhibit analgesic, anti-inflammatory, antipyretic, anticancer, antioxidant, nootropic, hepatoprotective, gastro protective, anti-ulcer, cardiovascular, anti-obesity, antiepileptic, anti-asthmatic, anti-diabetic, anti-urolithiasis, diuretic, local anaesthetic, anti-allergic, anthelmintic, wound healing, antimicrobial, immune-modulatory, and anti-diarrheal properties [9].

Deficiencies in micronutrients across the globe can be prevented or eliminated if adequate nutrients included in the diet [10]. *M. oleifera* is rich in macro and micronutrients like protein, carbohydrate, calcium, phosphorus, phosphorus, iron, vitamins, beta carotene and other bioactive compounds which are important for normal functioning of the body and prevention and curing of certain diseases [1]. Evidence defines clear and concrete information on the benefits of moringa seeds regarding human consumption, which is beneficial to human health. *M. oleifera* seed contain a significant amount of oil with a high-quality fatty acid thus Moringa oil could be a good substitute for olive oil in the diet. Our aim was to germinate and ferment moringa seeds to compare the benefits with the raw seeds.

According to our knowledge, there is not much comparative study done on raw, germinated and fermented *M. oleifera* seeds. Therefore, the present study on the chemical and nutritional composition of raw, germinated, fermented *M. oleifera* seeds, phytochemical content and their potential uses for human health which would be helpful to the mankind in many ways. Moringa is globally available plant and the seeds are a promising resource for food and non-food applications,

#### **II. MATERIALS AND METHODS**

The Family, genus and Species of *M. oleifera* was identified in the Plant Biotechnology Department, Women's Christian College, Chennai. Seeds of *M. oleifera* were collected from the market at Madhavaram, Chennai, Tamil Nadu. Raw *M. oleifera* seeds were sorted, rinsed, sundried, dehulled and milled using a mechanical blender. The sample is sieved to obtain raw *M. oleifera* (RMO) seed flour.

#### A. Germinated M. oleifera Seed (GMO) Flour

*M. oleifera* seeds were sorted and soaked in potable water for 2 hours. Excess water was removed and the seeds were tied in a wet cloth for 72 hours (3 days) at room temperature. Germinated Seeds were selected and washed thoroughly, sundried, dehulled and milled using a mechanical blender and sieved to obtain germinated *M. oleifera* (GMO) seed flour.

#### B. Fermented M. oleifera Seed (FMO) Flour

*M. oleifera* seeds were sorted, dehulled [11], and soaked in water for 72 hours (3 days) to ferment. The fermented seeds were sundried and milled using a mechanical blender and sieved to obtain fermented *M. oleifera* (FMO) seed flour.

All the three samples were packed in a plastic container sealed and stored at room temperature prior to analyses.

#### C. Preparation of Extract

About 20 g of each sample was soaked in 100ml of ethanol for 72 h. The pale-yellow supernatant liquid was filtered by Whatman filter paper. The extract was condensed in a hot plate at 50°C, which yields gummy extract.

Qualitative and quantitative analysis of phytochemicals, identification of volatile bioactive compounds using gas chromatography mass spectrometry (GC-MS), estimation of macro nutrients and selected micro nutrients and the antioxidant potential of the samples was estimated with the condensed sample. Table I gives the details regarding the analysis phytochemicals.

		-		
Sn.No	Compound	Name of test	Observation	Interpretation
1	carbohydrates	Fehling's test	A brick red precipitate	Presence
2		Molish test	Reddish brown ring at the junction of two layers	Presence
3	Alkaloids	Mayer's test	creamy white precipitate	Presence [12]
4		Hager's test	A prominent yellow precipitate	Presence [13]
5	glycosides	Borntrager's test	Pink color	Presence [12]
6	Saponins	Foam test	appearance of turbid foam	Presence [13]
7	phenolic compounds	Ferric chloride test	A dark green or blue colour	Presence [14]
8		Flavonoids	white precipitate	Presence
9	Terpenoids	Salkowski test	A reddish-brown coloration	Presence





ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538

Volume 11 Issue VIII Aug 2023- Available at www.ijraset.com

# **III.RESULTS**

A. Qualitative Assessment of Phytochemicals in RMO, GMO and FMO Seeds

Phytochemical screening was carried out for *in RMO*, *GMO and FMO* seeds to identify the presence and absence of major natural chemical groups such Carbohydrate, Protein, Alkaloids, Flavonoids, Saponins, Terpenoids, Tannins, Phenols, Steroids.

•				
S. No	Phytochemical test	RMO	GMO	FMO
1	Carbohydrate test	Positive	Positive	Positive
2	Protein test	Positive	Positive	Positive
3	Alkaloid test	Positive	Positive	Positive
4	Flavonoid test	Positive	Positive	Positive
5	Saponin test	Positive	Positive	Positive
6	Terpenoid test	Positive	Positive	Positive
7	Tannin test	Positive	Positive	Positive
8	Phenol test	Positive	Positive	Positive
9	Steroid test	Positive	Positive	Positive

#### TABLE II QUALITATIVE RESULTS FOR VARIOUS PHYTOCHEMICALS

- Presence of phytochemicals (Positive)
- Absence of phytochemicals (Negative)

### B. Quantification of Phytochemicals in in RMO, GMO and FMO Seeds

Table III shows the total the total flavonoid content in 100 grams of RMO, GMO and FMO is 129.37, 326.36 and 268.4  $\mu$ g QE/g respectively. 221.00, 464.19 and 339.14 mg/g gallic acid equivalent phenols is present in 100 grams of each sample.

S No	Phytochomical	Quantity Present			
5. 110	Fnytochennical	RMO	GMO	FMO	
1	Phenols (mg/g gallic acid equivalent)	221.00	464.19	339.14	
2	Flavonoids (µg QE/g) QE-Quercetin Equivalent)	129.37	326.36	268.4	

 TABLE III

 Estimation Of Total Phenol And Flavonoids Content in *in RMO*, *GMO AND FMO* SEEDS

# C. Identification of volatile bioactive compounds using Gas Chromatography Mass Spectrometry (GC-MS) in in RMO, GMO and FMO Seeds

The interpretation of GC - MS was elucidated using National Institute Standard and Technique (NIST). The particular compounds present in the sample were identified by matching their mass spectral fragmentation patterns of the respective peaks in the chromatogram with those stored In the National Institute Standard and Technology Mass spectral database library. The spectrum of the unknown components was compared with the spectrum of the known components stored in NIST library.

### 1) Volatile Bioactive Constituents Identified in RMO Seeds Using GC – MS

The following table IV presents the compounds that were identified by comparing the spectrum of the unknown components with that of the spectrum of the known compounds stored in the NIST library. Eight volatile bioactive components were identified in RMO. The compound name molecular formula, Structure, Molecular mass and the property are presented in table IV.

According to table IV the predominant volatile constituents in the RMO seeds are: Ethyl 2-quinoxalinecarboxylate, Methyl tetradecanoate, Pentadecanoic acid methyl ester, 1H-indene, 2-decyloctahydro, Hexadecanoic acid methyl ester, 8 – octadecenoic acid methyl ester, Z, Z-4,16-Octadecadien-1-ol acetate, Methyl eicosa-5,8,11,14,17 pentaenoate.



Volume 11 Issue VIII Aug 2023- Available at www.ijraset.com

# TABLE IV

Compounds Identified In Rmo Seeds Through Gas Chromatography – Mass Spectrometry Components That Eluted In The Chromatogram In The Increasing Order Of Polarity

				Molecular	
S.No	Peak	Formulae	Compound name	Mass	Property
				(g/mol)	
1	13.7	C11H10N2O2	Ethyl 2- quinoxalinecarboxylate	201.3	Antipsoriatic, Antineoplastic
2	15.0	C15H30O2	Methyl tetradecanoate	242.4	Not available
2	157		Pentadecanoic acid,		Antiacne agent, Antipsoriatic,
5	13.7	C16H32O2	methyl ester	256.4	Antiviral, Antioxidant
4	16.6		1H-indene, 2-	264.0	Not evoilable
4	10.0	C19H36	decyloctahydro	204.0	Not available
5	17.0		Hexadecanoic acid methyl	270.0	Antiasthmatic, antidepressant,
3	17.2	C17H34O2	ester	270.0	antibacterial, antiviral Agent
6	10.0		8 – octadecenoic acid,	206.0	Antionidant
0	18.9	C19H36O2	methyl ester	290.0	Antioxidant
7	21.1		Z, Z-4,16-Octadecadien-1-	200.0	NI-4
/	21.1	C20H36O2	ol acetate	508.0	INOL AVAIIABLE
0 000			Methyl eicosa-	216.0	Antipsoriatic, Antineoplastic,
ð	22.3	C21H32O2	5,8,11,14,17- pentaenoate	510.0	Anti-allergic

# 2) Volatile Bioactive Constituents Identified in GMO Seeds Using GC – MS

According to table V, the predominant volatile constituents in the GMO seeds are Undecanoic acid, 3-Methyl-5-(1, 4, 4-trimethylcyclohex-2-enyl) pentan-1-ol), Methaqualone, 9-Hexadecenoic acid, Hexadecanoic acid, methyl ester, Hexadecanoic acid, ethyl ester, 8-octadecenoic acid, methyl ester, Benzimidazole. Three compounds eluted are 9-Hexadecenoic acid of molecular mass 268.00 g/mol, Hexadecanoic acid, methyl ester of molecular mass 270.00 g/mol and 8- octadecenoic acid, methyl ester of molecular mass 269.00 g/mol.

TABLE V

Compounds Identified In Gmo Seeds Through Gas Chromatography – Mass Spectrometry Components That Eluted In The Chromatogram In The Increasing Order Of Polarity

S. No	Peak	Compound name	Formula	Molecular mass (g/mol)	Property
1	13.1	Undecanoic acid	C11H22O2	200.0	Antioxidant
2	14.1	3-Methyl-5-(1,4,4- trimethylcyclohex-2- enyl) pentan-1- ol)	C13H20O	224.0	Antioxidant
3	14.6	Methaqualone	C16H14N2O	250.0	Antibacterial, Anti Allergic
4	15	9-Hexadecenoic acid	C16H30O2	268.0	Antioxidant, Anticonvulsant, Antibacterial Agents
5	15.2	Hexadecanoic acid, methyl ester	C17H34O2	270.0	Antiasthmatic, Antidepressant, Antibacterial, Antiviral Agent
6	15.9	Hexadecanoic acid ethyl ester	C18H36O2	284.0	Antidepressant, Antibacterial
7	16.9	8-octadecenoic acid, methyl ester	C19H36O2	269.0	Antioxidant
8	17.9	Benzimidazole	C7H6N2	308.0	Anti – Infective



# International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538

Volume 11 Issue VIII Aug 2023- Available at www.ijraset.com

# 3) Volatile Bioactive Constituents Identified in FMO Seeds Using GC – MS

from table VI, the predominant volatile constituents in the *FMO Seeds* are: Vinyl lauryl ether, 5-Octadecene, Benzoic acid, 3,5-bis (1,1-dimethylethyl)-4- hydroxy, 9-Hexadecenoic acid, Hexadecanoic acid, 3-Eicosene, 9-Octadecenoic acid, 5- [4(Diethylamino) benzylideneamino] benzimidazol-2 (3H) one, 1-Tetradecene, 2-decyl.Three compounds peaked eluted are 9-Hexadecenoic acid of molecular mass 268.00 g/mol, Hexadecanoic acid, methyl ester of molecular mass 270.00 g/mol and 9- Octadecenoic acid, of molecular mass 296.00 g/mol.

### TABLE VI Compounds Identified In Fmo Seeds Through Gas Chromatography – Mass Spectrometry Components That Eluted In The Chromatogram In The Increasing Order Of Polarity

			U		U	5
	S. No	Peak	Compound name	Formula	Molecular mass (g/mol)	Property
Ī	1	13.0	Vinyl lauryl ether	C14H28O	212.0	Anti-infective, antiseptic
	2	13.7	5-Octadecene	C18H36	212.0	Not available
	3	14.6	Benzoic acid 3,5-bis (1,1- dimethylethyl)- 4-hydroxy	C15H22O 3	250.0	Anti-asthmatics, anticonvulsants, antidepressants
	4	15	9-Hexadecenoic acid	C16H30O 2	268.0	Antioxidant, antiseptic
	5	15.3	Hexadecanoic acid	C16H32O 2	270.0	Antidepressants, Antioxidant, Anti-infective
Ī	6	15.9	3-Eicosene	C20H40	280.0	Not available
	7	16.9	9-Octadecenoic acid	C18H34O 2	296.0	Anti-Parkinson drugs
	8	17.9	5 [4(Diethylamino) benzylideneam ino] benzimidazol-2(3H) one	C7H6N2	308.0	Anti-asthmatics, Antioxidant
	9	19.6	1-Tetradecene, 2- decyl	C24H48	336.0	Antioxidant

### D. Estimation of Macro Nutrients and Selected Micro Nutrients in RMO, GMO and FMO Seeds

From table VII, it is clear that 100g of each RMO, GMO and FMO seeds contain 379.49 kcal, 349.45 kcal and 354.44 Kcal respectively. RMO provides more energy than GMO and FMO seeds. Also, it was observed that 100 grams of each RMO, GMO and FMO seeds provide 33.22, 49.13 and 29.28g of carbohydrate. GMO seeds provide more carbohydrates than RMO and FMO seeds. On top of this, GMO seeds have a higher amount of protein than RMO and FMO seeds. Finally, it is evident that 100 grams of each RMO, GMO and FMO, GMO and FMO seeds contain 36.46 g, 25.98g and 31.19g respectively. All three samples provide approximately the same amount of fat. The main energy source of *M. oleifera* seeds is from fat. Thus *M. oleifera* seeds are used for extracting oil for various uses.

From table VII, 100 grams of each RMO, GMO and FMO seeds contain 5.6 g, 5.4 g and 5.9 g crude fibre respectively. All three samples provide approximately the same amount of crude fibre. But out of the three, FMO seeds provide more crude fibre. Thus, it is evident that GMO and FMO seeds have more nutritional value than the RMO seeds. On top of that. GMO seeds provide more phosphorus, iron, Ascorbic acid and calcium. Every part of *M. oleifera* is a storehouse of important nutrients and anti-nutrients. The seeds of *M. oleifera* are rich in minerals like calcium, phosphorus, zinc, magnesium, iron and copper are essential for growth and development [15].



# International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538

Volume 11 Issue VIII Aug 2023- Available at www.ijraset.com

Quantification Of Macro Nutrients in RMO, GMO and FMO Seeas						
		Quantity present in 100g of				
S. No	Nutrients		sample			
		RMO	GMO	FMO		
1	Energy (Kcal)	379.5	349.4	354.4		
2	Carbohydrates (g)	33.2	49.1	59.3		
3	Protein (g)	36.5	56.6	31.1		
4	Fat (g)	37.0	25.9	31.2		
5	Crude fiber (g)	5.6	5.4	5.9		
6	Phosphorus (mg)	33.3	34.2	32.8		
7	Calcium (mg)	132.2	137.9	124.8		
8	Iron (mg)	7.4	7.5	6.5		
9	Ascorbic acid (mg)	24.8	26.1	17.7		

	TA	BLE V	II					
Quantif	ication Of Macro Nutrie	ents In	RMO,	GMO	and	FMO	Seed	ls
		0				100	c	

# E. The Antioxidant Potential of RMO, GMO and FMO Seeds

Antioxidant has increased interest in order to prevent disastrous effects of the free radicals and prevent the deterioration of fats and other food items. In both cases, antioxidants have a natural preference compared to synthetic sources [16]. There is therefore a drastic increase in the use of methods for estimating the efficiency of such substances as antioxidants **Error! Reference source not found.** 

# 1) Free Radical Activity of the RMO, GMO and FMO Seeds using DPPH Assay

In the present study, the ethanol extract of *RMO*, *GMO* and *FMO* Seeds was used to investigate the antioxidant activity by DPPH (2,2-diphenyl-1- picryl hydrazyl) method and FRAP assay based on the rapid reduction in ferric-tripyridyltriazine (Fe<sup>III</sup>-TPTZ) by antioxidants present in the samples forming ferrous-tripyridyltriazine (Fe<sup>II</sup>-TPTZ), a blue-coloured product [18].

#### TABLE VIII

# COMPARISON OF RADICAL SCAVENGING ACTIVITY OF RMO, GMO AND FMO WITH STANDARD ASCORBIC ACID USING DPPH ASSAY AND FRAP (FE<sup>3+</sup>) ASSAY

	DPPH Assay			FRAP (Fe <sup>3+</sup> ) Assay		
Concentration	RMO	GMO	FMO	RMO	GMO	FMO
(µg/mL)	Mean (SD)	Mean	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)
	Mean (SD)	(SD)				
50	46.4 (1.2)	62.5 (0.9)	31.8 (1.2)	22.5 (1.2)	56.5 (1.2)	48.5 (1.2)
100	56.3 (1.0)	74.3 (0.6)	42.6 (1.5)	32.6 (0.3)	71.9 (0.6)	56.5 (0.6)
150	59.2 (0.7)	86.1 (1.2)	52.9 (0.7)	41.8 (0.9)	75.8 (0.9)	73.6 (1.2)
200	63.8 (1.3)	86.7 (0.8)	65.6 (0.2)	53.8 (1.0)	85.3 (1.0)	79.0 (0.7)
250	74.4 (1.5)	87.9 (0.6)	78.5 (1.0)	70.6 (0.7)	87.7 (1.7)	82.6 (1.0)
300	83.6 (1.0)	89.7 (1.0)	88.4 (0.8)	75.6 (1.0)	90.4 (0.2)	85.7 (0.7)
F Value	396.1	436.9	1174.1	1522.66	419.15	790.45
P Value	0.0001	0.0001	0.0001	0.0001	0.0001	0.0001

From the table VIII, the radical scavenging activity of RMO, GMO and FMO seeds increase with the concentration. It is found that the P value for RMO, GMO and FMO is 0.0001 (p<0.01) which is significant at 1 per cent level. Thus, there is a significant difference between the RMO, GMO and FMO was comparable to the percentage of inhibition of the reference standard (ascorbic



acid) at all concentrations.

2) IC50 (inhibition concentration 50) and RC50 (Reduction concentration at 50 %) in RMO, GMO and FMO Seeds

IC50 (inhibition concentration 50) is the antioxidant concentration ( $\mu$ g/mL) which is able to inhibit 50 per cent of the activity of free radicals. From the table 15, it is evident that GMO seed have the lowest IC50 value. 40.00  $\mu$ g/mL of the Moringa seed extract can have 50 per cent of inhibiting property. The IC50 value of RMO seed is 53.82  $\mu$ g/ml whereas the IC50 value of the FMO seed is 141.72  $\mu$ g/mL which is the highest. The samples when compared to the IC50 value of the standard ascorbic acid (2.9  $\mu$ g/mL) RMO, GMO and FMO samples requires higher concentration to reduce 50 per cent of free radical

COMPARISON OF 1C30 VALUE OF KIND, UND AND FIND SEEDS					
SAMPLE	IC50 VALUE	RC50 VALUE			
RMO	53.8 µg/Ml	179.5 μg/mL			
GMO	40.0 µg/mL	44.2 µg/mL			
FMO	141.7 μg/mL	551.6 µg/mL			
STANDARD ASCORBIC ACID	2.9 µg/mL				

TABLE IX COMPARISON OF IC50 VALUE OF RMO, GMO AND FMO SEEDS

RC50 (Reduction concentration at 50 %) is the antioxidant concentration ( $\mu$ g/ml) which is able to reduce 50 per cent of the activity of free radicals. From the table, it is evident that compared to the RC50 value of the standard ascorbic acid (2.88  $\mu$ g/mL) the test samples require a higher concentration to reduce 50 per cent of free radical. The RC50 value of Germinated Moringa seeds 44.21  $\mu$ g/mL which is very low compared to raw and fermented which is 179.46  $\mu$ g/mL and 551.56  $\mu$ g/mL respectively.

# **IV.DISCUSSION**

Natural plant products have been on the increase leading to the identification and improvement of plant products beneficial to mankind. According to Bagchi [19] 'Foods or dietary components that may provide a health benefit beyond basic nutrition,' he also stated that functional foods are 'Foods that by virtue of physiologically vital food elements provide health benefits beyond basic nutrition'. There is great scope for foods that can impart health benefits beyond traditional nutrients. Within more functional perspectives, health has been defined in terms of a certain quality of life which enables individuals to live happily, successfully, fruitfully, and creatively [20]. The current World Health Organization (WHO) definition of health, describes health as "a state of complete physical, mental and social well-being and not merely the absence of disease or infirmity.

Germination of seeds helps to add on to the nutrient content of the seeds, especially the amount of protein. Thus, Sprouts help to get protein from the least amount of seed consumption. Even though there are many other benefits to germinated seeds in food, more research must be done to improve texture and sensory properties to gain community acceptance [21].

Fermented foods are digested easily and reactive foods into health – giving foods. Fermentation destroys most of the harmful microorganisms and chemicals in foods and adds beneficial bacteria in the gut. The enzymes developed and bacteria grown during fermentation aids in the digestion of foods. But not all forms of fermentation or fermented foods are safe and nutritious. For example, certain microorganisms such as fungi can contaminate the foods and may enhance the production of N-nitroso compounds with potential carcinogenic properties [22]. Still fermenting seeds will help to increase the nutrient content and the probiotic concentration which have a lot of health benefits.

*M. oleifera* is a miracle plant in which all the parts of the plant are used for medicinal purposes. Leaves, pods, flower and seeds of *M. oleifera* are considered as a functional food. Root and bark acts as a cardiac stimulant, anti-ulcer and anti-inflammatory agent [23]. The root and bark contain minerals like calcium, magnesium and sodium and alkaloids like morphine, moriginine which helps the bark to be antiulcer and helps to relax the muscles [21]. The leaves of *M. oleifera* are rich in minerals like phosphorus , calcium, zinc, iron, magnesium and copper [15] and Vitamins like beta-carotene, vitamin B such as folic acid, pyridoxine and nicotinic acid, vitamin C, D and E [25]. Moringa leaves treat asthma, hyperglycaemia, Dyslipidaemia, flu, heart burn, syphilis, malaria, pneumonia, diarrhoea, headaches, scurvy, skin diseases, bronchitis, eye and ear infections [26] Also reduces, blood pressure and cholesterol and acts as an anticancer, antimicrobial, Antioxidant, anti-diabetic and anti- atherosclerotic agents, neuro-protectant [27]. The presence of minerals and vitamins help in boosting the immune system and cure a myriad of diseases [28]. Various amino acids and Phytochemicals provide anti-diabetic and antioxidant properties to the leaves.

Moringa seeds are rich in antioxidant thus it reduces the oxidative stress which is a risk factor to heart disease, heart failure, and

Volume 11 Issue VIII Aug 2023- Available at www.ijraset.com

high blood pressure [29].

High cholesterol in the blood is a risk factor of heart disease. In traditional Thai medicine, Moringa seeds are used as a cardiotonic which is a drug or herb used to improve the heart health [30]. A single serving of Moringa seeds has almost three times the amount of iron as spinach. This is especially important for vegetarians and vegans and those who suffer from iron deficiency anaemia [31]. Moringa seeds are beneficial for diabetes and blood sugar regulation. The scientists believe that the compounds which help in lowering the blood sugar level are isothiocyanates [32].

Fibre is an important factor for the maintenance of healthy cardiovascular system [33]. Moringa seeds are rich in soluble fibre, so that, it can help in maintaining a healthy digestive system [34]. Studies show that Moringa seed powder has an anti – proliferative effect on the breast cancer cell. [35]. Iron deficiency will lead to anaemia. Moringa seeds are high in iron. Moringa seeds can be taken as a substitution for iron supplements [36]. A Study done on experimental rats showed that the Moringa seed extract given twice daily had healed liver injury induced with liver fibrosis over the course of eight weeks [37]. The process by which the dormant embryo wakes up, grow out of the seed coat and establishes itself as a seedling is called germination [30].

Research revealed that the water extracts of *M. oleifera* seed kernel exhibited significant free radical scavenging activity. Thus, the study shows that the water extract possessed potent antioxidant activity which supports the use of *M. oleifera* seed kernel as a natural antioxidant [38]. Different parts of the plant such as bark, leaves, seeds, flowers, roots, and immature pods and mature pods contains large number of important bioactive constituents such as terpenoids, alkaloids, tannins, steroidal aglycones and reducing sugars [39]. The active components found in the seeds in the Aqueous and Hydro-alcoholic extract of *M. oleifera* seeds are Methionine, cysteine, 4-(alpha-L-rhamnopyranosyloxy) benzylglucosinolate, Moringine, benzylglucosinolate, niazimicinniazirin [4].

Hence, germinating and fermenting the Moringa seeds could be a viable approach to increase the nutritional benefits which would reduce hidden hunger and other health complications

### V. LIMITATIONS OF THE STUDY

- 1) All the phytochemicals could not quantify due to unavailability of reagents, cumbersomeness of the process and time constraints.
- 2) Quantification of all the micro nutrients was not done due to time and financial constraints.
- 3) There are various methods to assess antioxidant potential of the plant samples. However, in the present study, only preferred methods were used to analyse each of the activities. These assessments can also be carried out by various other methods which can provide more accuracy to the result.

#### VI.PROSPECTS FOR FUTURE RESEARCH

- 1) A study on the other plant parts of *M. oleifera* can be done.
- 2) The cancer prevention potential of raw, germinated and fermented *M. oleifera* seeds against various cancer types can be analysed.
- 3) Antifungal and anti-inflammatory activities of raw, germinated and fermented *M. oleifera* seeds can also be studied.
- 4) A study on *M. oleifera* seed oil can be done.

#### REFERENCES

- Sahay, S., Yadav, U., & Srinivasamurthy, S. Potential of Moringa oleifera as a functional food ingredient: A review. Int. J. Food Sci. Nutr.Vol. 2, Issue 5. 2017.
   Huang, W. Y., Cai, Y. Z., & Zhang, Y. Natural phenolic compounds from medicinal herbs and dietary plants: potential use for cancer prevention. Nutr Cancer.
- 62(1):1-20. 2010.
- [3] Hossan, S. J., El-Sayed, M., & Aoshima, H. Antioxidative and anti α-amylase activities of four wild plants consumed by nomads in Egypt. Orient Pharm Exp Med; 9(3): 217-224. 2009.
- [4] Paikra, B. K., Dhongade, H. K., & Gidwani. B. Phytochemistry and Pharmacology of Moringa oleifera Lam. J Pharmacopuncture. 20(3): 194–200. 2017.
- [5] Tiloke, C., Anand, K., Gengan, R M., & Chuturgoon, A. A. Moringa oleifera and their phytonanoparticles: Potential antiproliferative agents against cancer. Biomed Pharmacother. 108:457-466. 2018.
- [6] Padayachee. B., & Baijnath, H. An overview of the medicinal importance of Moringaceae. J Med Plants Res 6:5831–5839. 2012.
- [7] Sengupta A., & Gupta M. P. Studies on seed fat composition of Moringaceae family. FetteSeifenAnstrichm. 72:6–10. 1970.
- [8] Leone, A., Spada, A., & BatteZZati, A. Cultivation, genetic, ethnopharmecology, phytochemistry and pharmachology of Moringa oleifera leaves: an overview. Int J Mol sci 16:12791 – 12835. 2015.
- Bhattacharya, A. I., Tiwari, P., Sahu, P. K., & Kumar, S. A Review of the Phytochemical and Pharmacological Characteristics of Moringa oleifera. JPharmBioallied Sci. 10(4):181-191. 18. 2018.



# International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538

Volume 11 Issue VIII Aug 2023- Available at www.ijraset.com

- [10] Ramasubramania, R., Sreenivasulu, M., Vaishnavi, S., Muni, D., Navyasri, G., Raven, P. H., Evert, R. F., & Eichhon, S. E. Germination In: Biology of Plants, 7th Edition. W.H. Freeman and Company, New York, USA, 504- 508. 2005.
- [11] Bakebain, D. A., & Giami, S. Y. Proximate composition of full fat fluted pumpkin (Telfairia occidentalis) seed flour. J. Sci. Food Agric. 59:321–325. 1992.
- [12] Evans. Trease and evans pharmacognosy. 14th edition.Harcourt Brace and Company. Asia Pvt.Ltd. Singapore. 343. 1997.
- [13] Kokate, A. Phytochemical Methods. Phytotherapy, 2<sup>nd</sup> edition. 78:126-129. 1999.
- [14] Mace, M. E. Histochemical Localization of phenols in healthy and diseased tomato roots. Phytopathology 16:915-925. 1963.
- [15] Kasolo, J. N., Bimenya, G. S., Ojok, L., Ochieng, J., & Ogwal-okeng, J. W. Phytochemicals and uses of Moringa oleifera leaves in Ugandan rural communities. J. Med. Plants Res., 4, pp. 753-757. 2010.
- [16] Abdalla, A. E., & Roozen, J. P. Effect of plant extracts on the oxidative stability of sunflower oil and emulsion. Food Chemistry. 64: 323-329. 1999.
- [17] Moreno, S. C. Review: methods used to evaluate the free radical scavenging activity in foods and biological systems. Food Sci. Tech. Int., 8(3): 121-137. 2002.
- [18] Benzie, I. F., & Strain, J. The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay. J Anal Biochem. 239(1):70-6. 1996.
- [19] Bagchi, D. Neutraceutical and Functional Food Regulations. Elsevier: New York. 2008.
- [20] Lalas, S., & Tsaknis, J. Characterization of Moringa oleifera seed oil variety Periyakulam-1. J. Food Compos. Anal. 15, pp. 65-77. 2002.
- [21] Finar, I. L. Stereo chemistry and the chemistry of natural products. Vol.2. Longman, Singapore. pp 518. 1986.
- [22] Islami, F., Ren, J. S., Taylor, P. R., & Kamangar, F. Pickled vegetables and the risk of oesophageal cancer: a meta-analysis. Br J Cancer. 3; 101(9):1641-7. 2009.
- [23] Monera, T. G., & Maponga, C. C. Prevalence and patterns of Moringa oleifera use among HIV positive patients in Zimbabwe: a cross-sectional survey.J. Public Health Africa, 3, pp. 6-8. 2012.
- [24] Adeyemi. O. S., & Elebiyo, T. C. Moringa oleifera supplemented diets prevented nickel-induced nephrotoxicity in Wistar rats. J. Nutr. Metab. 4(3) pp. 1-8. 2014.
- [25] Mbikay, M. Therapeutic potential of Moringa oleifera leaves in chronic hyperglycemia and dyslipidemia: a review Front. Pharmacol. 3. pp. 1-12. 2012.
- [26] Ijarotimi, O. S., Adeoti, O., & Ariyo, O. Comparative study on nutrient composition, phytochemical, and functional characteristics of raw, germinated, and fermented Moringa oleifera seed flour. Food Sci. Nutr., pp. 452-463. 2013.
- [27] Rockwood, J. L., Anderson, B. G., & Casamatta, D. A. Potential uses of Moringa oleifera and an examination of antibiotic efficacy conferred by M. oleifera seed and leaf extracts using crude extraction techniques available to underserved indigenous populations. Int. J. Phytothearpy Res., 3, pp. 61-71. 2013.
- [28] Choudhary, M. K., Bodakhe, S. H., & Gupta, S. K. Assessment of the antiulcer potential of Moringa oleifera root-bark extract in rats. J. Acupunct. Meridian Stud, 6 (2013), pp. 214-220. 2013.
- [29] Nadeem, M., & Imran, M. Promising features of Moringa oleifera oil: recent updates and perspectives. Lipids Health Dis. 15:212. 10. 2016.
- [30] Dutta, A. C. The Seed In: Botany for Degree Students, 6thEdition. Published in India by Oxford University Press. PP:90-100. 2004.
- [31] Fuglie, J. L. The Moringa Tree: A local solution to malnutrition Church World Service in Senegal. Int J Mol sci. 4(1): pp :145 234. 2005.
- [32] De Vogel, E. F. Morphological types in dicot seedlings, with reference to their origin. Bull. Soc. Bot. France, 126 Actual. Bot., no. 3:17 3-182. 1980.
- [33] Moussa, L., Bézirard, V., Salvador, C., Bacquié. V., Lencina, C., Lévêque, M., Braniste, V., Ménard, S., Théodorou, V., & Houdeau, E. A low dose of fermented soy germ alleviates gut barrier injury, hyperalgesia and faecal protease activity in a rat model of inflammatory bowel disease. PLoS One. 7(11):47 – 49. 2012.
- [34] Oduro, I., Ellis, W. O., & Owusu, O. Nutritional potential of two leafy vegetables: Moringa oleifera and Ipomoea batatas leaves. Sci. Res. 3, pp. 57-60. 2008
- [35] Adebayo, I. A., Arsad, H., & Samian, M. R. Antiproliferative Effect on Breast Cancer of Moringa Oleifera Seed Extracts. Altern Med. 13;14(2):282-287. 2017.
- [36] Saini, R. K., Sivanesan, I., & Keum, Y. Phytochemicals of Moringa oleifera: a review of their nutritional, therapeutic and industrial significanceBiotech. 6(2). Pp-203. 2016.
- [37] Hamza, A. A. Ameliorative effects of Moringa oleifera Lam seed extract on liver fibrosis in rats. Food Chem Toxicol. 48(1):345-55. 2010.
- [38] Jahan, I. A., Hossain, H., Ahmed, K. S., Sultana, Z., Biswas, P., & Nada K. Antioxidant activity of Moringa oleifera seed extracts. Oriental Pharmacy and Experimental Medicine. 18(1). 2018.
- [39] Sharma, V. R., Paliwal, R., & Sharma, S. Phytochemical analysis and evaluation of antioxidant activities of hydro-ethanolic extract of Moringa oleifera Lam. J Pharm Res. 4(2):554–7. 2001.











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24\*7 Support on Whatsapp)