



IJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 10 **Issue:** XII **Month of publication:** December 2022

DOI: <https://doi.org/10.22214/ijraset.2022.48434>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Antibacterial effect of *Caesalpinia bonducella* Leaf Extract on *Salmonella typhi*

Rajesh Kumar

Research Scholar, University Department Of Zoology, B.N.M.University, Madhepura, BIHAR

Abstract: Plants are important source of treatment for various diseases. We are using them from prehistoric times in our folk medicines. Many plants have their different medicinal properties, among them one is *Caesalpinia bonducella*. It has many local names in different languages, it is known as *kantkaranja* in Hindi and *fever nut* in English. It is prickly shrub widely distributed all over the world specially in tropical areas. All parts of plant have medicinal properties, so it is utilized in traditional system of medicines. It is used for the treatment of several diseases. It is very popular in Ayurveda, Siddha, Unani and Homeopathic system of medicines. This review attempts to encompass the antityphoidal property of *Caesalpinia bonducella*.

Keywords: *Caesalpinia bonducella*, *Salmonella typhi*, Antityphoidal properties, *Kantkaranja*

I. INTRODUCTION

Caesalpinia bonducella is a medicinal plant belonging to the family caesalpiniaceae. It is prickly shrub widely distributed all over the world specially in India, Srilanka and Andaman and Nicobar Island's. In India this plant is specially found in Tropical regions. It is also widely found in Koshi and Purnia commissionery of Bihar. In India traditional plant medicine has been considered as an important remedy for the treatment of several diseases. It is popular in various indigenous system of medicines like Homeo-pathic, Ayurveda, Siddha and Unani. All parts of the plants have medicinal properties. So it is a very valuable traditional plant, which is utilized in traditional system of medicine. The plant has been reported to possess anxiolytic, antidiarrheal, antidiabetic, apoptogenic, anthelmintic, antiestrogenic, anti-inflammatory, antimalarial, antimicrobial, antifungal, antispasmodic, antioxidant, antiproliferative, antipsoriatic, antitumor, larvicidal, muscle contractile, hepatoprotective, anticonvulsant and antifilarial activities. Phytochemical analysis of *Caesalpinia bonducella* has revealed the presence of alkaloids, flavonoids glycosides, tanins and triterpenoids. This review attempts to encompass the available literature on and *Caesalpinia bonducella* with respect to this pharmacognostic characters, chemical constituents, summary of its various pharmacological activities and traditional uses.

Salmonella typhi is a genus of rod-shaped (bacillus) Gram-negative bacteria of the family *Enterobacteriaceae*. The two species of *Salmonella* are *Salmonella enterica* and *Salmonella bongori*. *S. enterica* is the type species and is further divided into six subspecies that include over 2,600 serotypes. *Salmonella* was named after Daniel Elmer Salmon (1850-1914), an American veterinary surgeon. *Salmonella* species are non-spore-forming predominantly motile enterobacteria with cell diameters between about 0.7 and 1.5 μm, lengths from 2 to 5 μm, and peritrichous flagella (all around the cell body, allowing them to move). They are chemotrophs, obtaining their energy from oxidation and reduction reactions, using organic sources. They are also facultative anaerobes, capable of generating ATP with oxygen ("aerobically") when it is available, or using other electron acceptors or fermentation ("anaerobically") when oxygen is not available. The genus *Salmonella* is part of the family of *Enterobacteriaceae*. Its taxonomy has been revised and has the potential to confuse. The genus comprises two species, *S. bongori* and *S. enterica*, the latter of which is divided into six subspecies: *S. enterica*, *S. e. salamae*, *S. e. arizonae*, *S. e. diarizonae*, *S. houtenae*, and *S. indica*. The taxonomic group contains more than 2500 serotypes (also serovars) defined on the basis of the somatic O (lipopolysaccharide) and flagellar antigens (the Kauffman-White classification). The full name of a serotype is given as, for example, *Salmonella typhimurium*. Further differentiation of strains to assist clinical and epidemiological investigation may be achieved by antibiotic sensitivity testing and by other molecular biology techniques such as pulsed-field gel electrophoresis, multilocus sequence typing, and increasingly, whole genome sequencing. Historically, *Salmonellae* have been clinically categorized as invasive (typhoidal) or non-invasive (nontyphoidal *Salmonellae*) based on host preference and disease manifestations in humans.

II. METHOD

A. Collection of Plant Materials

The leaves of *Caesalpinia bonducella* were collected in the month of April and May from adjacent areas of Rampur village of Hasanganj Block of Katihar District, Bihar, India.

B. Preparation of Powder

The collected leaves were washed shade dried and pulverized into a coarse powder using a suitable grinder for size reduction. It was passed through mesh (size 40) and the fine powder was stored in air tight containers and used for the experiment and preparation of extract.

C. Preparation of Extracts

The leaves were dried in shade and crushed in an electric grinder and by using soxhelt extraction using methanol as solvent. aqueous extract is prepared from standard methods. The antibacterial potentialities of methanol extract and its fraction of *C. bonducella* leaves were investigated by the disc diffusion method against *Salmonella typhi*, 500µg and 800 µg/disc. Kanamycin (30µg/disc) was used as the standard drug. Antimicrobial activities of leaf extracts were evaluated their cytotoxicity..

III. RESULT AND DISCUSSION

Culture test of *Salmonella typhi* shows different zones of inhibition with kanamycin and methanol extracts of 500 and 800µg/disc in disc diffusion method in petri dishes. Kanamycin shows 30 mm and *C.boduc* 800µg/disc shows 6 mm zone of inhibition but *C.boduc* 500 µg/disc shows no zone of inhibition.

Table 1 Antibacterial effect of *Caesalpinia bonducella* leaf extract fractions on *Salmonella typhi*

Name of bacteria	Zone of inhibition(mm)	Zone of inhibition(mm)	Zone of inhibition(mm)
	Kenamycin 30 mg /disc	C.boduc 500 µg/disc	<i>C.boduc</i> 800µg/disc
<i>Salmonella typhi</i>	30	-	6

A variety of constituents present in *Caesalpinia bonducella* such as saponin, tannin, polyphenols, flavonoids, and alkaloids, may be present in the fractions, further extensive investigations are required to determine the antimicrobial, antidiarrheal, and cytotoxic properties present in the leaf extracts.

IV. CONCLUSION

From the above study indicates that *Caesalpinia bonducella* extracts have medicinal potential of preventing culture of *Salmonella typhi*. These results further support the traditional use of this plant in folk medicine and more and more work have to be done in this field to develop new phytomedicine which do not develop drug resistant and have very less or no side effects of antibiotics.

REFERENCES

- [1] Gupta N.R.; Viswas K.; Pathak M.; Singh P.S.; Gupta A. Antibacterial activities of ethanolic extracts of plants use in folk medicine, International Journal of Research in Ayurveda and Pharmacy, 2010, 1(2), 529-535.
- [2] Halliwell, B.; Gutteridge, J.M.C. Free Radicals in Biology and Medicine; Oxford University Press: Oxford, UK, 1999.]
- [3] Hong, C.H.; Hur, S.K.; Oh, O.J.; Kim, S.S.; Nam, K.A Lee, S.K. Evaluation of natural products on inhibition of inducible cyclooxygenase (COX-2) and nitric oxide synthase (iNOS) in cultured mouse macrophage cells. J. Ethnopharmacol. 2002, 83, 153–159.]
- [4] Iyenger, M.A. Pendse, G.S. 1965. Anti- Diarrheal activity of the nuts of *Caesalpinia bonducella* FLEM. Indian Journal of Pharmacology 27:307.
- [5] Jabbar, A.; Zaman, M.A.; Iqbal, Z.; Yassen, M.; Shamin A. Antihelmintic activity of *Chenopodium album* (L.) and *Caesalpinia crista* (L.) against trichostrongylid nematodes of sheep. J. Ethnopharmacol. 2007, 114, 86–91. Jeong, G.S.; Lee, D.S.; Kwon, T.; Lee, H.; An, R.; Kim, Y. Cytoprotective constituents of the heartwood of *Caesalpinia sappan* on glutamate-induced oxidative damage in HT22 cells. Biol. Pharm. Bull. 2009, 32, 945–949.
- [6] Jiang, R.W.; Ma, S.; But, P.P.; Mak, T.C.W New antiviral cassane furanoditerpenes from *Caesalpinia minax*. J. Nat. Prod. 2001, 64, 1266–1272. [Jiang, T.; But, P.P.H.; Ma, S.; Ye, W.; Chan, S.; Mak, T.C.W. Structure and antiviral properties of macrocaesalmin, a novel cassane furanoditerpenoid lactone from the seeds of *Caesalpinia minax* Hance. Tetrahedron Lett. 2002, 43, 2415–2418.
- [7] Jun, H.; Xiaoling, Y.; Wei, W.; Hao, W.; Lei, H.; Li, J.D Antioxidant activity in vitro of three constituents from *Caesalpinia sappan* L. Tsinghua Sci. Technol. 2008, 13, 474–479. Kannur, D.M.; Hukkeri, V.I.; Akki, K.S. Adaptogenic activity of *Caesalpinia bonduc* seed extracts in rats. J. Ethnopharmacol. 2006, 108, 327–331.
- [8] K. Harini, J. Jerlin Shownwya and N. Geetha; Photochemical constituents of different extracts from the leaves of *Chromolaena odorata* (L) King and Robinson; International Journal of Pharmaceutical Sciences and Business Management, Dec.2014; Vol.2. Issue.12; 13-20.
- [9] K. Karthiashwaran, S. Minrunalini, G. Dhamodharan, M. Krishnaveni and V. Arulmozi; Phytochemical Investigation of methanolic extract of the leaves of *Pergularia daemia* of Biological science, 2010, 1-5.
- [10] K. Sivasnakari, S. Janaky and T. Sekar, Evaluation of phytochemicals in select medicinal plants of the *Caesalpinia* species, Indian Journal of Science and Technology; 3(12); 1118-1121.
- [11] Kannur, D.M.; Hukkeri, V.I.; Akki, K.S. Antidiabetic activity of *Caesalpinia bonducella* seed extracts in rats. Fitoterapia 2006, 77, 546–549.] Kapoor, L.D. 2009. Hand of Ayurvedic Medicinal Plants, CRC Press 87.



- [12] Karthi J, Thamizhmozhi M, Srabanan C, Ahamed Ayas K, Chakravarthy N Invitro anthekminitic activity of leaves extracts of Casealpinia bonducella (L.)Scholars Research library . 2011; 3(4):317-319
- [13] Keusch,GT, Bennish ML: Shigellosis.p.593 In Evans A S ,Brachman PS (ed): Bacterial Infections of Humans, Epidemiology and Control .Plenum , New York , 1991.
- [14] Khan , R; Islam ,B ., Akram , M., Shakil, S.; Ahmad , A., Ali , S.M.,Siddiqui, M.; Khan, A.U., Antimicrobial activity activity of five herbal extracts against multi drug resistant strains of bacteria and fungus of clinical origin, Molecules , 2009, 14, 586-597.
- [15] Khan, H.U., Ali. Antibacterial , Antifungal ,Antispasmodic and Ca++Antagonist Effects of Casealpinia bonducella Journal of Phutochemical , 25(4): 444-449.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)