



# **iJRASET**

International Journal For Research in  
Applied Science and Engineering Technology



---

# **INTERNATIONAL JOURNAL FOR RESEARCH**

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume: 13    Issue: V    Month of publication: May 2025**

**DOI: <https://doi.org/10.22214/ijraset.2025.70413>**

**[www.ijraset.com](http://www.ijraset.com)**

**Call:  08813907089**

**E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)**

# Antiphytopathogenic Potential of *Allium sativum* L.: An In Vitro Study

Shaikh Farah T<sup>1</sup>, Sahera Nasreen<sup>2</sup>.

<sup>1</sup>Department of Botany, B.S. Patel ACS College, Pimpalgaon kale, Jalgaon jamod, Dist-Buldhana, Maharashtra, India

<sup>2</sup>Department of Botany, Govt. Institute of Science, Nagpur, Maharashtra, India

**Abstract:** The available information on the phytochemical composition and antifungal activity of garlic clove against phytopathogens is very limited. The main aim of this research paper is to determine the in vitro antifungal activity of *Allium* cloves extract against some isolated fungi viz., *F. oxysporum*, *F. solani*, *F. moniliforme*, *H. sativum*, *C. lindemuthianum*, *C. lunata*, *R. solani*, *A. solani* from different vegetables. Both the aqueous and alcoholic extracts of *Allium* were screened for the phytochemical constituents and were tested against fungal pathogens. Result of phytochemical screening, revealed the presence of different metabolites in different extracts. *A. sativum* found to be highly effective against all eight fungal pathogens and highest inhibition of mycelial growth was recorded in *F. oxysporum*, *F. solani*, *F. moniliforme*, *H. sativum*, *C. lindemuthianum*, *C. lunata* with 92.63% While methanol extract of *A. sativum* when tested highest inhibition on mycelial growth showed by *F. solani* with 89.58%

**Keywords:** *Allium sativum* L, Phytochemical Screening, Fungal Pathogens, Aqueous extract, Methanolic Extract.

## I. INTRODUCTION

Agriculture is the backbone of Indian Economy. But in India, chemicals like fertilizers and pesticides are extensively used to boost crop yields and protect against pests and diseases. However, the excessive and unregulated use of these chemicals has led to significant environmental and health problems. These chemicals in addition kill various beneficial organisms and their toxicity can persist in the soil (Shaikh and Sahera.,2021). Excessive use of synthetic pesticides has been implicated in their negative effects on the environment such as soil and water pollutions, long periods of degradation, residual accumulation in the food chain, and less control efficacy against pathogenic microbes with long-term usage (Nega, 2014; Bhavaniramy et al., 2019).

The increasing resistance by these microorganisms against these chemicals has been a great concern. Among the various alternatives, natural plant products are used having no side effect and are been used by scientists. Extracts obtained from these valuable plants have gained attention as scientific interest for having antifungal activity (Santas et al., 2010). *Allium sativum* commonly known as garlic, is a member of Liliaceae family (Shruti S, 2018). When crushed, *Allium sativum* yields alliin and phytonicide which serve as antimicrobial compounds (Najeeb et al. 2016).

It also contains other compounds including alliin, ajoene, diallylsulfide, B-vitamins, proteins and many phytochemicals (Momoh et al.2016). Phytochemicals are bio-active chemicals of plant origin. They are regarded as secondary metabolites because the plant that manufactures them may have little need for them. They are naturally synthesized in all parts of the plant body; bark, leaves stem, root, flower, fruits, seeds, etc. i.e. any part of the plant body may contain active components (Tiwari et al. 2010). Bioactive compounds in the plants which confer protection on them against bacteria, fungi and viruses have been linked to the antimicrobial activities of the extracts of such plants. (El-Mahmood and Amey, 2007). *Allium sativum* consists of various kinds of sulfur-containing compounds with different significances and these compounds give garlic its characteristic odor and favor as well as some favorable effects. Sulfur-containing compounds present in garlic include alliin, alliin, ajoene, 2- Vinyl-4H-1,3-dithiin, Diallyl sulfide, Diallyl disulfide, Diallyl trisulfide and Allyl methyl sulfide (Mardomi, et al. 2017).

## II. MATERIAL AND METHOD

### A. Preparation of Aqueous Plant Extract

Thoroughly washed fresh plant material (50 g) was macerated with 50 ml sterile distilled water in a blender for 10 min. The macerate was first filtered through double-layered muslin cloth, and then centrifuged at 4000 g for 30 min; the supernatant was filtered through Whatman No. 1 filter paper and sterilized at 120°C for 30 min. The extract was preserved aseptically in a brown bottle at 5°C until further use.

**B. Preparation of solvent plant extract:**

Extracts were prepared in methanol at room temperature by simple extraction method (Deshpande *et al.*, 2004). Collected plant parts were shade dried and ground to a fine powder using grinder mixer. Dried powder of plant parts (10g) was mixed with 100 ml solvent in 250 ml conical flask. The flasks were plugged tightly with cotton and wrapped with papers. All conical flasks were kept on shaker for 24 h then it was allowed to stand for five hours to settle the plant materials. Thereafter, it was filtered and centrifuged at 5000 rpm for 15 min. The supernatant was collected and the solvent was evaporated at 45 °C in vacuum evaporator to make the final volume 1/5th of the original volume. It was stored at 4 °C in airtight bottles for further studies.

These aqueous and methanolic extracts were further used for screening of antifungal activity against eight fungal pathogens causing vegetable fruit diseases.

**C. Antifungal Activity of Plant Extracts**

The plant extracts were evaluated *in vitro* through Poison food technique (Nene and Thapliyal, 2000). The supernatant was taken as standard plant extract solution (100%). Further, the extract was diluted by adding sterilized water to get different per cent concentrations. The plant extracts were subjected to boiling temperature of 50°C in water bath to avoid contamination and then incorporated into PDA media by transferring 2ml of each type of plant extract in to a Petridish containing 20 ml melted warm PDA medium and gently shaken for thorough mixing of the extract. The PDA plates containing the plant extracts were inoculated aseptically with different pathogens by transferring 6mm diameter agar disc of 07 days old culture of the pathogen to the centre of PDA medium in Petridish. Three replications were maintained for each treatment. The basal medium (PDA) without any phytoextract served as control. All the inoculated Petridishes were incubated at 25±1°C. The radial growth of the test fungus in the treated plates was measured in all treatments when the pathogen growth touched the periphery in the control Petridishes. The per cent inhibition of fungal growth was estimated by using the formula given by Vincent (1927).

$$I = \frac{C - T}{C} \times 100$$

Where, I = per cent inhibition.

C = Colony diameter in control.

T = Colony diameter in treatment.

**III. RESULT**

Present study reveals that tannins, alkaloids and flavonoids were absent in aqueous extract of *A. sativum* bulbs. *A. sativum* found to be highly effective against all eight fungal pathogens and highest inhibition of mycelial growth was recorded in *F. oxysporum*, *F. solani*, *F. moniliforme*, *H. sativum*, *C. lindemuthianum*, *C. lunata* with 92.63% but *A. solani* showed resistant with 87.57% inhibition. Whereas *R. solani* showed 92.60% inhibition of mycelial growth over control. While methanol extract of *A. sativum* when tested highest inhibition on mycelial growth showed by *F. solani* with 89.58% and lowest inhibition by *H. sativum* with 61.38%. Methanol extract of *A. sativum* were significantly effective and revealed 73.94% inhibition of mycelial growth of all eight tested pathogens.

Table 1: Phytochemical Analysis of *Allium* extracts.

Secondary Metabolites	Extracts	
	H2O	MeOH
Tannin	A	A
Alkaloid	A	P
Saponins	P	P
Cardiac Glycosides	P	P
Steroids	P	P
Flavonoids	A	A

A-Absent, P-Present

Table 2- Inhibitory effect of plant extracts of different concentrations on mycelial growth of fungal pathogens of vegetables.

Pathogens/Medicinal Plant	Mycelial growth and percent inhibition	<i>Allium sativum</i>	
		H2O Extract	MeOH Extract
<i>A. solani</i>	Mycelial Growth (mm)	6.6	21.3
	% Inhibition over control	87.57±0.54	76.22±1.43
<i>C. lindemuthianum</i>	Mycelial Growth (mm)	6.6	22.6
	% Inhibition over control	92.63±0.54	70.91±0.94
<i>C. lunata</i>	Mycelial Growth (mm)	6.6	31.3
	% Inhibition over control	92.63±0.54	64.94±0.71
<i>F. moniliforme</i>	Mycelial Growth (mm)	6.6	13.3
	% Inhibition over control	92.63±0.54	85.10±1.18
<i>F. oxysporum</i>	Mycelial Growth (mm)	6.6	23.3
	% Inhibition over control	92.63±0.54	73.99±0.71
<i>F. solani</i>	Mycelial Growth (mm)	6.6	9.3
	% Inhibition over control	92.63±0.54	89.58±0.54
<i>R. solani</i>	Mycelial Growth (mm)	6.6	27.3
	% Inhibition over control	92.60±0.94	69.42±0.97
<i>H. sativum</i>	Mycelial Growth (mm)	6.6	34.6
	% Inhibition over control	92.63±0.54	61.38±1.18

Value expressed in mean ± S. E. M of triplicate




Fig- Aqueous Extract of *Allium sativum* against different plant pathogens




Fig- Methanolic Extract of *Allium sativum* against different plant pathogens

#### IV. DISCUSSION

Present study revealed the presence of Saponin, Steroids and cardiac glycosides in different extracts [methanol extract and distilled water extract of garlic. A recent study on phytochemical analysis of *Allium sativum* (garlic) revealed the presence of saponins, terpenoids, flavonoids, amino acids and cardiac glycosides (Arify et al.,2018). Another study on phytochemical profile of aqueous extract of *Allium sativum* L., bulbs showed the presence of saponins, steroids, tannins, carbohydrates and cardiac glycosides (Mikail H.G.2010).

The extract possesses antimicrobial properties that can suppress the growth of fungi due to the presence of active sulfur-containing bioactive compounds such as allicin (Bayan *et al.*, 2014). These compounds act by interfering with fungal cell metabolism and inhibit spore germination (Redondo-Blanco *et al.*, 2020),

The finding of this study correlate with the finding of (Abaoab *et al.*2011) which found that clove extract possessed a broad spectrum of antimicrobial activity exhibited for both bacteria and fungi due to presence saponin, tannin, flavonoid and terpenoid. The result of this study on Phytochemistry of Garlic supported the study conducted by (Deresse.,2010).

This inhibition was concentration dependent. A similar survey by Alhussaeen *et al.*, (2011) indicated that garlic extract had a concentration dependent activity against *Pythium ultimum* isolated from tomato seedlings. Undiluted garlic extract showed a high control activity with no growth as compared to the biotic control without the extract whereas diluted garlic extracts 10% and 5% reduced the fungal growth to 15.5% and 41% respectively (Alhussaeen *et al.*, 2011).

The mechanism of action of phytochemicals against microorganism vary and depend on these phytochemicals (Aly and Bafiel.,2008). In conclusion, we have shown that garlic extracts can have a significant effect on preventing the growth of almost all targeted pathogens. However, growth of *H. sativum* was not significantly suppressed by the methanolic extract of garlic. The solvent used for dilution concentrations (water or Methanol) had an influence on the antifungal activity of the garlic extract. Aqueous dilutions of the extract had greater antifungal activity than Methanol diluted extracts, possibly because the longer the extracts were exposed to ethanol, the more the antifungal activity was reduced which was in support of the findings of Chanel *et al.*2014.

#### V. CONCLUSION

In conclusion, the in vitro results of this study confirmed the potentiality of *A. sativum* as one of the best sources for controlling fungal growth. Hence, further work is necessary to evaluate its potentiality in vivo on targeted pathogens. This can provide an alternative means for the control of vegetable diseases by farmers. Investigations are also needed to characterize, formulate and market the active principles of these extracts which may provide avenues for the discovery of novel antifungal compounds. These biofungicidal botanicals are environmentally safe; therefore, they could successfully replace the toxic and hazardous synthetic compounds and be exploited as ideal treatment for future plant disease management programs.

## REFERENCES

- [1] Shaikh Farah T and Sahera Nasreen., (2021). Phytochemical screening and evaluation of antimicrobial activity of Zingiber Officinale L. International Journal of Creative Research Thoughts. 9(5): e829-e834.
- [2] Nega, A., (2014). Review on Concepts in Biological Control of Plant Pathogens. J. Biol. Agric. Healthc. 4, 33–55.
- [3] Santas J, Almajano MP, Carbo R. (2010). Antimicrobial and antioxidant activity of crude onion (*Allium cepa* L.) extracts. International Journal of Food Science Technology, 45: 403–409.
- [4] Shruti S., (2018) Studies on antimicrobial and phytochemical properties of *Allium sativum* extracts, International Journal of Innovative Research in Science, Engineering and Technology, 7, 5371-5376.
- [5] Najeeb U., Abida P., Rahat B., Iqra Z., Mukharma M., Sadia J., Amna L. and Sohail A., (2016) In vitro and in vivo protocols of antimicrobial bioassay of medicinal herbal extracts: A review, Asian Pacific Journal of Tropical Disease, 6, 660-667
- [6] Momoh J., Oluremi N.O. and Odetunde S.K., Antimicrobial and antioxidant properties of aqueous garlic (*Allium sativum*) extract against *Staphylococcus aureus* and *Pseudomonas aeruginosa*, British Microbiology Research Journal, 14, 1-11 (2016)
- [7] El-Mahmood, A.M., Amey J.M. 2007. In vitro antibacterial activity of *Parkia biglobosa* root bark extract against some microorganisms associated with urinary infections. Afr. J. Biotechnol., 6(11): 1272-1275.
- [8] Mardomi R (2017) Determining the Chemical Compositions of Garlic Plant and its Existing Active Element. IOSR Journal of Applied Chemistry 10(1): 63-66.
- [9] Nene, Y. L. And P.N. Thapliyal. (1979), Fungicides in Plant Disease Control, Oxford and IBH Publishing House, New Delhi, pp 163.
- [10] Vincent, J. M., (1927). Distortion of fungal hyphae in presence of certain inhibitors. Nature, 59: 850.
- [11] Arify T., Ezhilvalavan S., Varun A., Sundaresan A. and Manimaran K., (2018) Qualitative phytochemical analysis of garlic (*Allium sativum*) and nilavembu (*Andrographis paniculata*), International Journal of Chemical Studies, 6, 1635-1638
- [12] Mikail H.G., (2010) Phytochemical screening, elemental analysis and acute toxicity of aqueous extract of *Allium sativum* L. bulbs in experimental rabbits, Journal of Medicinal Plants Research, 4: 322-326
- [13] L. Bayan, P.H. Koulivand and A. Gorji. Garlic: a review of potential therapeutic effects. Avicenna J Phytomed. 2014; 4(1):1-14. PMID: 25050296; PMCID: PMC4103721
- [14] S. Redondo-Blanco, J. Fernández, S. López-Ibáñez, E.M. Miguélez, C.J. Villar, and F.Lombó. Plant phytochemicals in food preservation: Antifungal bioactivity: A review. J. of Fd Prot. 2020. 1;83 (1):163-71. DOI: 10.4315/0362-028X.JFP-19-163
- [15] Aboaba OO, Ezeh AR, Anabuike CL (2011) Antimicrobial activities of some Nigerian spices on some pathogens. Agriculture and Biology Journal of North America 2(8): 1187-1193.
- [16] Deresse D (2010) Antibacterial Effect of Garlic (*Allium sativum*) on *Staphylococcus aureus*: An in vitro Study. Asian Journal of Medical Sciences 2(2): 62-65.
- [17] Alhussaen, K. Hussein, E.L., Khalid, A.M., Mahmoud, A., Wesan, A., Jacob, H.J., Shatnawi, A.M., Ashraf, K., Hegazy, M.I. 2011. Identification and controlling *Pythium* species infecting tomato seedlings cultivated in Jordan valley using garlic extract. Asian J. Plant Pathol., 5: 84-92.
- [18] Aly, M.M., Bafiel, S. 2008. Screening for antimicrobial activity of some medicinal plants in Saudi Arabia. World conference on Medicinal and Aromatic plants.
- [19] Chanel K. Daniel, Cheryl L. Lennox and Filicity A. Vries. 2014. In-vitro effects of garlic extracts on pathogenic fungi *Botrytis cinerea*, *Penicillium expansum* and *Neofabraea alba*. South African Journal of Science. 111(7/8):1-8



10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)