



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 11 Issue: I Month of publication: January 2023

DOI: https://doi.org/10.22214/ijraset.2023.48824

www.ijraset.com

Call: 🕥 08813907089 🔰 E-mail ID: ijraset@gmail.com



Assessment of *Jangal Desha* in Western Geo-Climatic Region of Rajasthan Based on *Ayurvedic* Parameters

Dr. Shubham Shukla¹, Dr. Rajesh Kumar Sharma², Dr. Dinesh Chandra Sharma³ ¹P.G. Scholar, Deptt. of Kriya Sharir, U.P.G.I.A.S.&R. Jodhpur ²Professor, Deptt. of Kriya Sharir, U.P.G.I.A.S.&R. Jodhpur ³Associate professor, Deptt. of kriya Sharir, U.P.G.I.A.S.&R. Jodhpur

Abstract: Desha and Disha are concepts used in Ayurveda to describe both health and sickness. Desha has been used in two ways. The first is for land/any geographical place, and the second is for body and mind. Atura Desha is the name given to humans. Both forms of Desha are discussed in Ayurvedic texts, and their interrelationship is also acknowledged. Desha is the name for a geographical area's topography-climate-vegetation profile in both thoughts (Ayurveda & modern). So, there was a need to explore this gap by a study that what part will be consider as a Jangal Desha. The paper intends to explain the clinical perspective & application of concept of Jangal desha with both thoughts. Keywords: Ayurveda, Desha, Jangal Desha, Western Rajasthan.

I. INTRODUCTION

Ayurveda is the ancient medical science of India. Apart from giving information on medical conditions and their treatment; it also gives valuable information on geographical, and social conditions of India. In Ayurveda the term Desha has been applied for both form as Disha (direction) and Desha (place) directed or divided part. In reference of site and location Desha has been applied and described in two-dimensional way in Ayurvedic classics known as Bhumi Desha (Specific part of land or geographical part) and Atura Desha (the site of disease body and mind). Both are important for the maintenance of health prevention and cure of disease too).

Commonly Desha term is in sense of site location, land etc. Desha is one of the factors which have to be assessed in ascertaining the *Hetu* (etiological factors) and also in deciding the treatment.

Medical geography which is gaining significance in contemporary medicine has its origin from Hippocrates. However, Charaka Samhita which was written centuries before Hippocrates elaborately speaks about medical geography. In reference of architecting, collection of drugs, quality of river water property of air etc. are described on the basis of Bhumi Desha. The air, water coming from specific direction has specific property. Bhumi desha - land region is of three kinds viz, ⁽¹⁾ Jangala desha (Dry / Arid land), Anupa desha (Marshy land) and Sadharan desha (Mixed land). Jangala desha (Dry / Arid land) is Characterized by rocky, sandy with full of stones, appearance of mirages in summer, small ponds, open spaces, land is like the sky, hot weather, less water, scanty rainfall, abundance of sunlight and air. Predominance of dry wind and arid desert.Plants are mostly thorny, harsh, strong and rough in nature.

A. Aims and Object

To understand correlation between western geo - climatic region of Rajasthan and Jangal Desha.

B. Material and Methods

Review of literature related to western geo – climatic region of Rajasthan and *Jangal Desha* were collected from *Brihatrayi*, Modern literature, available commentaries on it and research article are also searched from various authentic digital sources.

C. Concept of Jangal Desha

Desha is not only provided medical conditions and treatments, but it also provides essential information on India's geographical and social situations.



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 11 Issue I Jan 2023- Available at www.ijraset.com

II. ETYMOLOGY

The word Jaangla Pradesh is derived from two words, i.e, Jaangla + Pradesh

A. Jaangla

According to a Sanskrit english dictionary of m. monier Williams $\overline{vl}, \overline{s}, \overline{cl}$ means is Arid, sparingly grown with trees and plants (though not stead of in unfertile)¹. The word "Jaangal" in "Vaidak sabd sindhu" means "desert air countries"².

B. Desha

Desha means "habitat," which refers to the place where we live. *Desha* has a significant influence on a person's health. In *Rogi, Desa* is also known as one of the *Parikshya Bhavas*. It plays a crucial function in the aetiology of illness. The word "*Desha*" in "Vaidak sabd sindhu" means "*Sthane*", which is called "*Desha*" in the universe². According to Paarijaat Sanskrit hindi sbd kosh "*Desha*" means *Sthan, Pradesh, Vibhag*³. According to Rajsthani hindi sankshkrit kosh means *Desha* is territory considered under a rule or area near birth place⁴. After lots of observation, it was found that geographical variations and climate of the place of origin of the drug is the major factor in influencing the potency of drug, the reference of *Desh* (region) is present since the *Vedic Kaal* in the form of *Aarya- Aanarya Desh*. Later in *Samhita* as *Trividha Desh : Jangam Desh, Anup Desh* and *Sadharan Desh*.

C. Classification of Desha on the basis of Panchamahabhuta:

The character of that particular *Mahabhuta* dominates in that particular *Desha*. This factor is most important in classification of *Trivida Desha*. Acharya Sushruta has given *Panchabhautika* classification of *Bhumi* on the basis of overall appearance of that area⁵.

- 1) Parthiva Bhumi Desha: Full of heavy rocks, greyish or blackish coloured soil and huge trees are present⁶.
- 2) Apya Bhumi Desha: Smooth, full of water and grass, delicate trees and whitish soil are present⁷.
- 3) Agneya Bhumi Desha: Different coloured of soil, mixed with plenty of small stones and smaller trees are present⁸.
- 4) Vayaviya Bhumi Desha: Rough and ash-coloured stones, with plenty of dry small trees with holes are found⁹.
- 5) Akasiya Bhumi Desha: The land is soft, even with holes in it and is having tasteless huge trees and mountains. The colour of the soil is greyish black in colour¹⁰.

D. Classification of Desha / Bhumi on the basis of Rasa

- The soil is of six types on the basis of Rasa i.e.
- 1) Madhure Bhumi
- 2) Amla Bhumi
- 3) Lavana Bhumi
- 4) Katu Bhumi
- 5) Tikta Bhumi and
- 6) Kashaya Bhumi

III. FEATURES OF DESHA ACCORDING TO DIFFERENT ACHARYAS

Charak Samhita	Shushrut	Ashtang Sangrah	Ashtang Hridya	Sarangdhar	Harit Samhita
	Samhita			Samhita	
	By doing such	From a medical	There are two sorts	Three forms of	In a Desha where
Two types of	anoop, jangle and	standpoint, the	of Desha, according	Desha	there is a wide
countries have been	Sadharn, the	Desha is divided	to Ayurvedic theory:	distinctions have	range of
described by Acharya	desha becomes of	into two	1. Bhumi Desh 2.	been described by	mountains
Charak.	three types.	categories. Bhumi	Deha Desh.	Sharangadhar	covered with
One of these is known	It is called	and <i>Deha Desh</i> .	Deha desh refers to	acharya.	sharp, raucous
as Bhumi Desh, while	'Jangal Desh'	There are three	the various sections	Jangal Desh	and thorny, where
the other is known as	because it is a	sorts of land	of the body.	possesses the	the land which
Aatur Desh.	country that is	among these. For	The Bhumi Desh	attributes listed	produces deer and
Jangal, Anoop, and	equal to the sky,	instance, Jangal	been said to be	below. arid	leafless stunted



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 11 Issue I Jan 2023- Available at www.ijraset.com

Sadharan are the three	devoid of height	(a place where	of three types -	desert: for	trees stand in the
types of <i>Bhumi Desha</i> .	and low, where	there is only	1. Jaangle	example. India's	direction.
Jangal Desh is	there are few.	forest). Anoon (a	Desh	Bikaner and	scorching by the
Paryakash-Bhuvishtha	small and thorny	place where there	2. Anoon Desh	Jaisalmer regions.	intense rays of the
in these three types of	trees, water rains	is a lot of rain or	3. Sadharan	as well as	sun, the land and
countries, meaning it	less and water is	numerous rivers).	desh	Arabia's and	ponds have dried
has a clear sky or an	less in springs	and <i>ordinary</i> (a	Jaanole Desha	Africa's outer	up without water
open sky. It contains	wells, stepwells.	place where there	here the wind	regions, are	where only the
densely developed	etc., and a hot and	is neither much	blows more ¹³ .	examples. Often.	water of wells
Kadar (Bitstdir).	strong wind blow.	forest nor much		these are	Only when cows
Khadir. Asan	where small	water).		countries with	and buffaloes fill
(Vijavasar).	mountains are	Medicines, birds.		little or no water	their stomachs by
Ashwakarna (Sakhua	located at some	and humans		and few trees	chewing dry
with expanded leaves)	point where the	which are		There are more	grains there is no
Pay. Tinish Salai. Sal.	people are mostly	frequently		Pittaja, Raktaj,	tenderness in the
Somvask Ber Tendu	strong and have	generated in the		and Vaatai	inice and meat
Peepak Vat and	delicate bodies	Jangal pradesh		illnesses here ¹²	where cold air is
Amla trees There are	and where	are generally			not circulating
also Shami Ariun	Vaat and	Vata-Pradhan ¹⁴			where there is no
and Sheesham trees	<i>Pitta</i> diseases are	vana i raanan .			cultivation of
Due to the great	more prevalent ¹⁵				sugarcane then
velocity of the wind	more prevalent .				there is an
the stiff and dry young					outbreak of blood
trees of this country					and hile soon
are seen dancing. The					Such a state is
Mrigamrichika is read					called <i>Jangal</i>
here, and the land is					$Desh^{11}$.
abundant in <i>Tanu</i> .					2000
Khar. Paraush. Sikta					
(sand), and					
Sharkra (Kankari).					
This region is visited					
by Lav, Titar, and					
<i>Chakor</i> . This country					
is dominated by the					
Vata and Pitta					
Doshas, and the					
people who dwell here					
are stable and hardy ⁱ .					
Acharya <i>Charak</i> while					
describing the Agrava					
substances has said					
that <i>Maru</i> land is a					
healthy Desha.					
Having been born in a					
place where people					
are naturally powerful,					
such as Sindh, where					
residents are naturally					



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 11 Issue I Jan 2023- Available at www.ijraset.com

strong. This is due to			
the unique properties			
of certain locations. A			
man's strength is			
enhanced by his birth			
in such a location.			
Diseases are occurring			
at a lower rate ¹⁶ .			

A. Contemporary Thoughts on Jaangal Desh

According Morden classification of climate (Koppen classification), Due to the similarity in the characteristics of *Jangal Desh* and Arid or semi-arid region, the boundary of *Jangal Desh* can be determined from the Arid and Semi-Arid regions.

Ayurveda	Contemporary View	Rainfall (cm)	Temperature	
Jangala Desha	Tropical semiarid steppe	38.10 - 72.2	$20^{\circ}\mathrm{C} - 28.8^{\circ}\mathrm{C}$ In	
			December 32.8 [°] C in May	
	Tropical desert climate	<30.5	Above 34.5 [°] C	

IV. MODERN CONCEPT OF WESTERN GEO – CLIMATIC REGION OF RAJASTHAN¹⁷

In modern main geographical features of Rajasthan's arid and semi-arid regions are described below according to Resource Atlas of Rajasthan, State Remote Sensing Application Centre Department of Science & Technology Government of Rajasthan Jodhpur. The current Flora covers an area of around 25,000 square kilometres in the N.W. section of Rajasthan State, between 250 2' and 28° 10' N and 690 30' to 740 E. The arid and semiarid plains of the districts of Jodhpur (in the east), Barmer (in the south), and Jaisalmer (in the west) are included (in the W. & N.W.). The shape of the area, which is around 100 metres above sea level, is roughly rhomboidal. Pakistan shares its western boundary over the entirety of its length. Bikaner and Nagaur districts form the north-eastern boundary, while Pali and Jalore districts form the south-eastern boundary. In terms of geography, it is the eastern extension of the huge Saharao-Thar desert. The majority of the area is made up of a dry undulating plain of hardened sand, while the remainder is mostly made up of a rolling plain of loose sand that forms shifting sand-dunes of longitudinal and transverse types with lengths ranging from 2 to 10 kilometres and heights ranging from 30 to 80 metres. These dunes are oriented in the directions of North-East, East-West, and South-West, and are frequently generated at right angles to the direction of the south-westerly wind. Rivers are scarce, tiny, torrential, and only flow during certain seasons. A substantial portion of this region is said to have remained under the tremendous seas of the great Indian ocean in the distant past. Indeed, in many parts of the Jaisalmer district, the illusion of a receding tide that has 'forgotten' to return persists, leaving vast swaths of thirsty land interrupted by systematized arcs of sand-dunes. The arid conditions are thought to have developed as a result of climatic oscillations that occurred after the last phase of glaciation during the Pleistocene period. The area is a desolate wasteland of sand, unfriendly and sterile, and becoming more so as one travels westward. Agriculture, as one might imagine, is underdeveloped and limited due to the low water table and weather circumstances. Human settlements, in the form of small villages and villages, are few and far between, clearly reliant on the availability of potable water. The average population density in Barmer district is 30 people per square kilometre, 46 people per square kilometre in Jaisalmer, and 73 people per square kilometre in Jodhpur districts, with occasional lows of 1.30 and 1.36 people per square kilometre in Ramgarh and Sam sub-divisions of Jaisalmer district. A considerable portion of the population lives a nomadic or seminomadic lifestyle, with livestock as their primary source of income. With the exception of a few metalled roads linking the key oasis towns of Jaisalmer and Barmer, the region is largely traversed by foot or cart trails. Only a single railway line connects Jodhpur and Jaisalmer, the district headquarters, about half way west.

A. Climate

Extremes of temperature, severe drought accompanied by high wind velocity, low relative humidity, evaporation greatly surpassing precipitation, and insufficient rainfall to support any appreciable vegetation characterise the climate of this region.



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 11 Issue I Jan 2023- Available at www.ijraset.com

The hot season lasts from April to June, with May often being the hottest month.During these months, the average maximum temperature ranges from 39 to 42° C, while the average minimum temperature ranges from 25 to 28 C. The average maximum temperature for the period is between 28.50 and 27.50 degrees Celsius. The cold weather in the region occurs from December through January, with the mean lowest temperature ranging from 6.00 to 10.5° C in January. As a result, rainfall in this area is uncommon. In comparison to Barmer (14.5) and Jodhpur (12.6), there are fewer such days in Jaisalmer (12.6). The average rainfall in Jodhpur over the last 10 years has been 315.26 mm, 305.14 mm in Barmer, and 202.82 mm in Jaisalmer. As one travels westward, a declining pattern emerges. The monsoon season, which runs from June through September, sees the most rainfall. Winter showers are uncommon. Droughts occur on a regular basis. The relative humidity is lowest during the summer months, such as March, April, and May, and highest during the monsoon months, such as July, August, and September. It is, however, at its lowest point in April and at its maximum point in August. The relative humidity varies dramatically throughout the day, which is an essential but predictable aspect of the desert. Over the course of the year, the wind tends to blow from the south-west to the west. Dust storms, locally known as andhi, are common and can reach speeds of up to 136 km/h in hot weather. During the winter, though, the wind speed is much lower. Easterly breezes are frequently connected with rain.

B. Geology

The rocks in western Rajasthan are mostly covered by wind-blown sand, but they are exposed near the surface in a few spots. The Kailana-Jodhpur-Mandore plateau is a broad sandstone plateau that rises suddenly from the plains. It is of Vindhyan origin and often consists of a fine, gritty, reddish substance that is good for construction. The Jaisalmer plateau and outliers, which are rich in fossils, are composed of yellowish, brittle brown sandstone with a large quantity of well-preserved sea shells. Rajasthan's physiography is the result of many years of erosion and depositional processes. The geological formations and structures have had a significant impact on the current landforms and drainage systems.

Four major physiographic regions can be identified within the state.

- 1) The western desert (Thar),
- 2) The Aravalli hill region,
- 3) The eastern plains and
- 4) The south-eastern plateau region.

C. Vegetation (Trees and Shrubs)

The state of Rajasthan is home to a diverse range of flora that may be divided into two separate groups: dry vegetation found in the western portion of the state, and semi-arid to sub-humid vegetation found in the eastern and southern parts of the state.

D. Vegetation of Western Zone

Western Rajasthan is known for its scant vegetation, which is impacted by a variety of climatic, edaphic, and biotic conditions. The vegetation of the dry tract has adapted admirably to the harsh environment. The potential vegetation of this region has been altered by intense biotic forces. What is visible today is the deteriorated stage, and in extreme degradation, one can see relict vegetation. When considering a greater area of vegetation coverage, however, the vegetation type is determined by species dominance and co-dominance. The dominant and co-dominant species of the vegetation have largely been identified using two indices: density and frequency.

- *1)* Calligonum-Haloxylon-Leptadenia Type
- 2) Salvadora oleoides-Euphorbia caducifolia Type
- 3) Zizyphus nummularia-Capparis decidua Type
- 4) Suaeda Fruticosa-Salsola baryosma Type.
- 5) Prospois-Capparis-Zizyphus Type
- 6) Prosopis-Tecomella Type
- 7) Prosopis cineraria-Acacia nilotica Type
- 8) Salvadora oleoides-Prosopis cineraria-Capparis decidua Type
- 9) Acacia nilotica Type
- 10) Acacia Senegal-Euphorbia cadducifolia Type



Volume 11 Issue I Jan 2023- Available at www.ijraset.com



V. RAINFALL PATTERN

Rajasthan has a diverse climate that ranges from desert to sub-humid. Hyperthermic conditions characterise the entire state. The state's annual rainfall varies dramatically. Isohytes have a typical north-west to south-east tendency. West of the Aravalli range, rainfall decreases rapidly and dramatically, making western Rajasthan the driest area of the state. The average annual rainfall in this region ranges from less than 10 cm in the northwestern part of Jaisalmer (the lowest in the state), to 20 to 30 cm in the Ganganagar, Bikaner, and Barmer regions, 30 to 40 cm in the Nagaur, Jodhpur, Churu, and Jalor regions, and more than 40 cm in the Sikar, Jhunjhunun, and Pali regions, as well as along the western fringes of the Aravalli the rainfall on the eastern slope of the Aravalli ranges from 55 cm in Ajmer to 102 cm in Jhalawar. The districts of Banswara (92.0 cm) and Jhalawar (95 cm) receive the most yearly rainfall in the plains. Mount Abu (Sirohi district) in the state's south-west, on the other hand, receives the most rainfall (163.8 cm). The months of July and August have the highest average monthly rainfall in most regions. The number of rainy days during this time varies greatly depending on where you are. From ten in Jaisalmer to forty in Jhalawar and forty-eight in Mount Abu. The rest of the season sees rainfall ranging from 2.1 cm in Jaisalmer to 7.2 cm in Jaipur, spread out over 2.5 to 6 wet days.

A. Temperature Regimes

All over the state, considerable changes in diurnal and seasonal temperature ranges occur, demonstrating the most typical phenomena of the warm-dry continental climate. The month of March heralds the start of summer, with temperatures gradually climbing throughout April, May, and June. During this time, the temperature rise is nearly uniform across the state. The maximum daily temperatures in the western region of the state, primarily in Bikaner, Phalodi, Jaisalmer, and Barmer, range from 40°C to 45°C. During the summer, it can reach 49°C on rare occasions. Summers have a greater diurnal temperature variation. At night, the minimum daily temperature decreases significantly and remains between 20°C to 29°C. Day temperatures are similar on the eastern side of the Aravalli range, but night temperatures are around 26°C. Temperatures are lower at Udaipur and Mount Abu, where the mean daily maximum temperature in July is 38°C and 31.5°C, respectively. These two stations have daily minimum temperatures of roughly 25°C and 22°C, respectively.Summer isotherms (June) show temperatures ranging from 42°C (Sirohi district) to over 47°C (Ganganagar district). The dry zone in the west and the semi-arid zone of the mid-west, which make up a large portion of the state, had average maximum temperatures of 45°C this month. The isotherms in the south-eastern sections, however, show a downward tendency from 43°C to 37°C. The month of January is the coldest of the year.

The winter season lasts from December through February, with minimum temperatures dropping significantly in December and January. Some areas, including as Sikar, Churu, Pilani, and Bikaner, may see nighttime temperatures as low as minus 2°C. The rapid release of heat radiation from the sandy soil just after nightfall causes a dramatic drop in night temperatures in the dry and semi-arid zones of western Rajasthan. Secondary Western disturbances that primarily sweep western, northern, and eastern Rajasthan during the winter months, causing light rainfall, can also reduce night temperatures and increase wind speed, resulting in a wind chill effect. the majority of Rajasthan experiences mean temperatures of over 10°C.



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 11 Issue I Jan 2023- Available at www.ijraset.com

B. Agro-Climatic Zones

During the winter months, secondary depressions (western disturbances) accompanied by ice-cold winds from higher latitudes may bring the entire state of Rajasthan under the grip of a cold wave for 2 to 5 days.

The agricultural cropping pattern is influenced by the climatic conditions of a place, and different areas produce different crops as a result. Rainfall, temperature, humidity, wind velocity, and sunshine duration, among other meteorological conditions, have a considerable impact on cropping patterns. Annual rainfall and its distribution throughout the year, as well as diurnal and annual temperature regimes, are by far the most important elements affecting agriculture and people's lifestyles.

C. Ground Water Resources

Rajasthan is heavily reliant on ground water resources due to a paucity of surface water. The state's accessible ground water resources were first assessed in the mid-1970s, and the report has since been updated. The depth to water varies greatly across the state; shallow water levels have been observed in the canal command areas of Ganganagar, Banswara, Kota, and Bundi districts, while higher depth to water values have been observed in Rajasthan's western districts, particularly Jaisalmer, Bikaner, Barmer, and Jodhpur. The eastern part of Rajasthan has substantially more ground water development than the western part. Due to low and unpredictable rainfall, a lack of surface water sources, and significant evapotranspiration, yearly ground water recharge in western Rajasthan is quite low. However, because the storage in some thick aquifers is many times the yearly recharge, steady pumping can be performed even during a dry spell without inflicting harm.

VI. SIMILARITY BETWEEN WESTERN GEO – CLIMATIC REGION OF RAJASTHAN AND JANGAL DESHA

Acharya Charak has clearly mentioned the features of jangal desha in kalpa sthan. According to these features we can mark a Bhoomi or Desh as a jangal desh.

(Samhita)Atlas of Rajasthan)Less number of water bodies (lake, pond etc.) (ch.vi 3/47),Less in numberAbundance of sunlight (ch.vi.3/47)Mean duration of bright sun in this region is 8.0 to 8.8 hr /day. Maximun sunshine period of 10 to 10.5 hr/ day in mayAbundance of air (ch.vi.3/47)Average wind speeds of more than 8.8 miles per hourWater rains less (sh.su. 35/50)With yearly rainfall ranging from 10 to 40 cmSmall and thorny trees (sh.su. 35/50)Haloxylon-Leptadenia Type Euphorbia caducifolia Type Capparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C	Fetures of Jangal Desha	Features of arid or semi-arid region (Resource
Less number of water bodies (lake, pond etc.) (ch.vi 3/47),Less in numberAbundance of sunlight (ch.vi.3/47)Mean duration of bright sun in this region is 8.0 to 8.8 hr /day. Maximun sunshine period of 10 to 10.5 hr/ day in mayAbundance of air (ch.vi.3/47)Average wind speeds of more than 8.8 miles per hourWater rains less (sh.su. 35/50)With yearly rainfall ranging from 10 to 40 cmSmall and thorny trees (sh.su. 35/50)Haloxylon-Leptadenia Type Capparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C	(Samhita)	Atlas of Rajasthan)
Less number of water bodies (lake, pond etc.) (ch.vi 3/47),Less in numberAbundance of sunlight (ch.vi.3/47)Mean duration of bright sun in this region is 8.0 to 8.8 hr /day. MaximunSunshine period of 10 to 10.5 hr/ day in maysunshine period of 10 to 10.5 hr/ day in mayAbundance of air (ch.vi.3/47)Average wind speeds of more than 8.8 miles per hourWater rains less (sh.su. 35/50)With yearly rainfall ranging from 10 to 40 cmSmall and thorny trees (sh.su. 35/50)Haloxylon-Leptadenia Type Euphorbia caducifolia Type Capparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C		
Less number of water bodies (lake, pond etc.) (ch.vi 3/47),Less in numberAbundance of sunlight (ch.vi.3/47)Mean duration of bright sun in this region is 8.0 to 8.8 hr /day. MaximunAbundance of sunlight (ch.vi.3/47)Mean duration of bright sun in this region is 8.0 to 8.8 hr /day. MaximunAbundance of air (ch.vi.3/47)Average wind speeds of more than 8.8 miles per hourWater rains less (sh.su. 35/50)With yearly rainfall ranging from 10 to 40 cmSmall and thorny trees (sh.su. 35/50)Haloxylon-Leptadenia Type Euphorbia caducifolia Type Capparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C <td></td> <td></td>		
pond etc.) (ch. vi 3/47),Mean duration of bright sun in this region is 8.0 to 8.8 hr /day. MaximunAbundance of sunlight (ch.vi.3/47)Mean duration of bright sun in this region is 8.0 to 8.8 hr /day. MaximunAbundance of air (ch.vi.3/47)Average wind speeds of more than 8.8 miles per hourAbundance of air (ch.vi.3/47)Average wind speeds of more than 8.8 miles per hourWater rains less (sh.su. 35/50)With yearly rainfall ranging from 10 to 40 cmSmall and thorny trees (sh.su. 35/50)Haloxylon-Leptadenia Type Euphorbia caducifolia Type Capparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C	Less number of water bodies (lake,	Less in number
Abundance of sunlight (ch.vi.3/47)Mean duration of bright sun in this region is 8.0 to 8.8 hr /day. MaximunSunshine period of 10 to 10.5 hr/ day in maysunshine period of 10 to 10.5 hr/ day in mayAbundance of air (ch.vi.3/47)Average wind speeds of more than 8.8 miles per hourWater rains less (sh.su. 35/50)With yearly rainfall ranging from 10 to 40 cmSmall and thorny trees (sh.su. 35/50)Haloxylon-Leptadenia Type Euphorbia caducifolia Type Capparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C	pond etc.) (ch.vi 3/47),	
Abundance of sunlight (ch.vi.3/47)Mean duration of bright sun in this region is 8.0 to 8.8 hr /day. MaximunSunshine period of 10 to 10.5 hr/ day in maysunshine period of 10 to 10.5 hr/ day in mayAbundance of air (ch.vi.3/47)Average wind speeds of more than 8.8 miles per hourWater rains less (sh.su. 35/50)With yearly rainfall ranging from 10 to 40 cmSmall and thorny trees (sh.su. 35/50)Haloxylon-Leptadenia Type Euphorbia caducifolia Type Capparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C		
Abundance of sunlight (ch.vi.3/47)Mean duration of bright sun in this region is 8.0 to 8.8 hr /day. Maximun sunshine period of 10 to 10.5 hr/ day in mayAbundance of air (ch.vi.3/47)Average wind speeds of more than 8.8 miles per hourWater rains less (sh.su. 35/50)With yearly rainfall ranging from 10 to 40 cmSmall and thorny trees (sh.su. 35/50)Haloxylon-Leptadenia Type Euphorbia caducifolia Type Capparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C		
region is 8.0 to 8.8 hr /day. Maximunsunshine period of 10 to 10.5 hr/ day in mayAbundance of air (ch.vi.3/47)Average wind speeds of more than 8.8 miles per hourWater rains less (sh.su. 35/50)With yearly rainfall ranging from 10 to 40 cmSmall and thorny trees (sh.su. 35/50)Haloxylon-Leptadenia Type Euphorbia caducifolia Type Capparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C	Abundance of sunlight (ch.vi.3/47)	Mean duration of bright sun in this
Abundance of air (ch.vi.3/47)Sumshine period of 10 to 10.5 hr/day in mayAbundance of air (ch.vi.3/47)Average wind speeds of more than 8.8 miles per hourWater rains less (sh.su. 35/50)With yearly rainfall ranging from 10 to 40 cmSmall and thorny trees (sh.su. 35/50)Haloxylon-Leptadenia Type Euphorbia caducifolia Type Capparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C		region is 8.0 to 8.8 hr /day. Maximun
sunshine period of 10 to 10.5 hr/ day in mayAbundance of air (ch.vi.3/47)Average wind speeds of more than 8.8 miles per hourWater rains less (sh.su. 35/50)With yearly rainfall ranging from 10 to 40 cmSmall and thorny trees (sh.su. 35/50)Haloxylon-Leptadenia Type Euphorbia caducifolia Type Capparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C		
mayAbundance of air (ch.vi.3/47)Average wind speeds of more than 8.8 miles per hourWater rains less (sh.su. 35/50)With yearly rainfall ranging from 10 to 40 cmSmall and thorny trees (sh.su. 35/50)Haloxylon-Leptadenia Type Euphorbia caducifolia Type Capparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C		sunshine period of 10 to 10.5 hr/ day in
Abundance of air (ch.vi.3/47)Average wind speeds of more than 8.8 miles per hourWater rains less (sh.su. 35/50)With yearly rainfall ranging from 10 to 40 cmSmall and thorny trees (sh.su. 35/50)Haloxylon-Leptadenia Type Euphorbia caducifolia Type Capparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C		may
Abundance of air (ch.vi.3/47)Average wind speeds of more than 8.8 miles per hourWater rains less (sh.su. 35/50)With yearly rainfall ranging from 10 to 40 cmSmall and thorny trees (sh.su. 35/50)Haloxylon-Leptadenia Type Euphorbia caducifolia Type Capparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C		
Item and the set of the set	Abundance of air (ch.vi.3/47)	Average wind speeds of more
Water rains less (sh.su. 35/50)With yearly rainfall ranging from 10 to 40 cmSmall and thorny trees (sh.su. 35/50)Haloxylon-Leptadenia Type Euphorbia caducifolia Type Capparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C		than 8.8 miles per hour
10 to 40 cmSmall and thorny trees (sh.su. 35/50)Halox ylon-Leptadenia TypeEuphorbia caducifolia TypeEuphorbia caducifolia TypeCapparis decidua TypeCapparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C	Water rains less (sh.su. 35/50)	With yearly rainfall ranging from
Small and thorny trees (sh.su. 35/50)Haloxylon-Leptadenia TypeEuphorbia caducifolia Type Capparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C		10 to 40 cm
Euphorbia caducifolia Type Capparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C	Small and thorny trees (sh.su. 35/50)	Haloxylon-Leptadenia Type
Capparis decidua TypeHot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C		Euphorbia caducifolia Type
Hot and strong wind blows (sh.su. 35/50)Can reach speeds of up to 136 km/h in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C		Capparis decidua Type
in hot WeatherScorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C	Hot and strong wind blows (sh.su. 35/50)	Can reach speeds of up to 136 km/h
Scorching by the intense rays of the sun (hs 4/5-6)Average maximum temperature ranges from 39 to 42° C		in hot Weather
sun (hs 4/5-6) from 39 to 42° C	Scorching by the intense rays of the	Average maximum temperature ranges
	sun (hs 4/5-6)	from 39 to 42° C



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 11 Issue I Jan 2023- Available at www.ijraset.com

VII. DISCUSSION

Ayurveda is a health science that has evolved over time to care for human life. It is a holistic science that focuses on an individual's complete well-being. This upbeat attitude stems from the fact that a person is born and raised in a specific location. The same concept dictates that the nature of land be evaluated in order to determine an individual's unique characteristics. *Desha* is one of the *Pareekshyabhavas*¹⁸ (factors to be studied) concepts that must be addressed with great care. The determination of a diseased person's birthplace and growth provides a clear picture of the disease's prognosis and hence aids in treatment protocol; conversely, in a healthy individual, it aids in excavating the susceptible *Nidanas* and thus disease susceptibility. As a result, it helps *Ayurveda* achieve its primary goals of *"swasthasyaswasthyarakshanam*" and *"Aturasyavikaraprashamanam"*¹⁹. Modification of current lifestyles in response to external factors is urgently needed in order to maintain an individual's homeostasis and, as a result, improve health. *Disha* comes from the word *'Dish Nirdesh,'* which is mentioned in nine *'Karana Dravyas'* in Indian philosophy and *Ayurveda. Desha*, which is directed or split from one part to another, is also known as *Disha* (direction).

VIII. CONCLUSION

Desha is a term used to describe a site's location, land, and so on. *Desha* has been applied and discussed in two dimensions in Ayurvedic classics known as *Bhumi Desha* (Specific part of land or geographical part) and *Atura Desha* with regard to place and location (the site of disease body and mind). Geographical land is divided into three categories based on its unique characteristics and ecological status: *Jangala Desha* (dry land), which is dominated by *Vata, Anoopa* (marshy land), which is dominated by *Kapha*, and *Sadharana* (normal land), which is dominated by all three *Doshas*. The ayurvedic name for a certain geographical area's topography-climate-vegetation profile is '*Desha*.' Individual human constitution is defined by the term '*Prakriti*'. *Desha* and *Prakriti* are made up of comparable source materials, and as a result, they interact frequently and have unique characteristics arising from their common origins. A specific geographical area's terrain, climate, and vegetation are profiled by *Desha*. *Each Desha* is dominated by one of the *Mahabhootas*²⁰, and as a result, each *Desha* has a natural affinity for the matching *dosha*. *Anoopa desha* (wet and cold regions), for example, has a natural connection with *Kapha dosha* because both are products dominated by *Jala mahabhoota*. In addition, the qualities of the *desha* themselves reveal the source *mahabhootas* along their virtue. So, regarding this fact i considered some of district Rajasthan as a Jangal Pradesh. These districts are Barmer, Bikaner, Churu, Ganganagar, Hanumangarh, Jaisalmer, Jalor, Jodhpur and Nagaur which are parts of western rajasthan.

REFERENCES

- [1] Sanskrit english dictionary of m. monier Williams Pg. no. 417.
- [2] Vaidak sabd sindhu Pg. no. 490
- [3] पारिजात संस्कृत-हिन्दी शब्दार्थकोशः सम्पादक: प० ईश्वरचन्द्रः
- [4] राजस्थानी-हिन्दी संक्षिप्त शब्दकोश Pg. no. 675
- [5] Shastri AD, editor, (1st Ed.). Vol-1, Sushruta Samhita of Sushruta, Sutrasthana; Bhumipravibhgiy Adhyaya: Chapter 37, verse 5. Varanasi: Chaukhambha Sanskrit Sansthan, reprint 2017; page no 180.
- [6] Shastri AD, editor, Vol-1, Sushruta Samhita of Sushruta, Sutrasthana; Bhumipravibhgiy Adhyaya: Chapter 37, verse 5. Varanasi: Chaukhambha San-skrit Sansthan, reprint 2017; page no 180.
- [7] Shastri AD, editor, Vol-1, Sushruta Samhita of Sushruta, Sutrasthana; Bhumipravibhgiy Adhyaya: Chapter 37, verse 5. Varanasi: Chaukhambha San-skrit Sansthan, reprint 2017; page no 180.
- [8] Shastri AD, editor, Vol-1, Sushruta Samhita of Sushruta, Sutrasthana; Bhumipravibhgiy Adhyaya: Chapter 37, verse 5. Varanasi: Chaukhambha San-skrit Sansthan, reprint 2017; page no 180.
- [9] Shastri AD, editor, Vol-1, Sushruta Samhita of Sushruta, Sutrasthana; Bhumipravibhgiy Adhyaya: Chapter 37, verse 5. Varanasi: Chaukhambha San-skrit Sansthan, reprint 2017; page no 180.
- [10] Shastri AD, editor, Vol-1, Sushruta Samhita of Sushruta, Sutrasthana; Bhumipravibhgiy Adhyaya: Chapter 37, verse 5. Varanasi: Chaukhambha San-skrit Sansthan, reprint 2017; page no 180.
- [11] Tripathi Hari Prasad, editor, Harit Samhita; Deshkaalbalabalm Adhyaya: Chapter 4, verse 5-6. Varanasi: Chaukhambha Krishandas Acadami, reprint 2017; page no 17-18.
- [12] Dr Tripathi Brahmanand, editor, Sharangdhar Samhita, Purvkhand; Pratham Adhyaya: Chapter-1, verse 64. Varanasi: Chaukhambha Subharti Prakashan, Reprint 2020; page no.16.
- [13] Dr Tripathi Brahmanand, editor, Astanghridya Samhita, Sutrasthan; Ayushkamiya Adhyaya: Chapter-1, verse 23. Varanasi: Chaukhambha Subharti Prakashan, Reprint 2014; page no.18-19.
- [14] Dr Tripathi Ravidutt, editor, Astangsangrah Samhita, Sutrasthan; Ayushkamiya Adhyaya: Chapter-1, verse 46. Varanasi: Chaukhambha Sanskrit Pratishthan, Reprint 2011; page no.20.
- [15] Shastri AD, editor, (1st Ed.). Vol-1, Sushruta Samhita of Sushruta, Sutrasthana; Aturopkramniya Adhyaya: Chapter 35, verse 49-50. Varanasi: Chaukhambha Sanskrit Sansthan, reprint 2017; page no 175.
- [16] Chaturvedi G, Pandey K, editor, Charak Samhita of Agnivesha Vol-2, Kalpasthana; Madanphalkalp Adhyay: Chapter-1, verse 8. Varanasi: Chaukhambha bharti Acadami, Reprint 2012; page no 893.



- [17] Resource Atlas of Rajasthan, Government of Rajasthan. Pg. no. 18-149
- [18] Chaturvedi G, Pandey K, editor, Charak Samhita of Agnivesha Vol-1, Vimansthana; Rogvishagjitiya Adhyay: Chapter-8, verse 64. Varanasi: Chaukhambha bharti Acadami, Reprint 2012; page no 764.
- [19] Chaturvedi G, Pandey K, editor, Charak Samhita of Agnivesha Vol-1, Sutrasthana; Arthedashmahamuliya Adhyay: Chapter-30, verse 26. Varanasi: Chaukhambha bharti Acadami, Reprint 2012; page no 587.
- [20] Shastri AD, editor, Vol-1, Sushruta Samhita of Sushruta, Sutrasthana; Bhumipravibhgiy Adhyaya: Chapter 37, verse 5. Varanasi: Chaukhambha San-skrit Sansthan, reprint 2017; page no 180.











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24*7 Support on Whatsapp)