



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 14 **Issue:** X **Month of publication:** October 2026

DOI: <https://doi.org/10.22214/ijraset.2026.85214>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

Connecting Primordial Gravitational Waves with the Cosmic Microwave Background: Analytical and Observational Insights

Rupesh Kumar¹, Md. Mushtaque Khan²

¹Research Scholar, ²Professor and Head University Department of Mathematics, J.P. University, Chapra, Bihar

Abstract: Primordial gravitational waves provide a direct test of the dynamics and energy scale of the early Universe, but their observable imprint is filtered through the thermal history, radiative transfer, Galactic emission, and weak gravitational lensing. This study develops the connection from a quantized transverse–traceless metric perturbation to cosmic microwave background temperature and polarization spectra. The inflationary tensor spectrum is normalized by the tensor-to-scalar ratio r ; its propagation is governed by a damped wave equation with a calculable neutrino anisotropic-stress correction; and Thomson scattering projects its local quadrupole into parity-odd B-mode polarization. The resulting signal is concentrated in a reionization feature at multipoles below about 10 and a recombination feature near multipole 80. A multifrequency likelihood must distinguish this signal from polarized dust, synchrotron radiation, instrumental leakage, and lensing-induced B modes. Published analyses give $r_{0.05} < 0.036$ from BK18 alone and $r_{0.05} < 0.032$ when Planck PR4, BK18, baryon-acoustic-oscillation, and CMB-lensing information are combined. Numerical evaluations of the slow-roll tensor spectrum, the inflationary energy scale, and the exact matter-era transfer function clarify how these bounds propagate backward to inflation. The analysis shows that further progress depends jointly on low-frequency and high-frequency foreground leverage, stable polarization calibration, large-scale sky coverage, and delensing rather than on map depth alone.

Keywords: primordial gravitational waves; cosmic microwave background; B-mode polarization; inflation; tensor-to-scalar ratio; foreground separation; delensing.

I. INTRODUCTION

A stochastic background of tensor perturbations is a generic consequence of amplifying quantum vacuum fluctuations in a time-dependent geometry. Grishchuk first identified the parametric amplification of gravitational waves in an expanding isotropic universe [1], and Starobinsky calculated an early nearly scale-invariant relic spectrum [2]. The gauge-invariant perturbation framework later placed scalar and tensor production within a common mathematical description [3]. These results make primordial gravitational waves unusual among inflationary observables: their amplitude is set directly by the expansion rate during horizon exit, without an inflaton sound-speed or matter-transfer normalization.

The cosmic microwave background (CMB) is sensitive to tensor wavelengths whose present frequencies are roughly 10^{-18} – 10^{-16} Hz. Tensor perturbations alter the temperature quadrupole and generate a polarization pattern with nonzero curl. The parity decomposition of linear polarization showed that this curl, conventionally called the B mode, is produced by tensor perturbations but not by linear scalar perturbations [4], [5], [6]. This selection rule is the central observational advantage. It does not, however, make the measurement background-free: gravitational lensing converts scalar E modes into B modes, and the Milky Way produces polarized dust and synchrotron emission across every usable sky region. The inference problem is consequently a chain rather than a single proportionality. A primordial power spectrum must first be evolved across radiation and matter domination, projected through the visibility function at recombination and reionization, convolved with the instrument beam and sky mask, and fitted simultaneously with foreground and calibration parameters. An upper limit on r is meaningful only after these operations have been included in the covariance and likelihood. The same chain also explains why a deeper map at one frequency cannot by itself establish a primordial origin.

II. PRIMORDIAL TENSOR PERTURBATIONS

A. Geometry and canonical dynamics

In conformal time η , introduce a transverse–traceless perturbation h_{ij} of a spatially flat Friedmann–Lemaître metric. Spatial indices are raised with δ_{ij} , a prime denotes $d/d\eta$, and the reduced Planck mass is $M_{\text{Pl}} = (8\pi G)^{-1/2}$.

$$ds^2 = a^2(\eta) \{-d\eta^2 + [\delta_{ij} + h_{ij}(\eta, \mathbf{x})]dx^i dx^j\}, \quad \partial_i h_{ij} = 0, \quad h_{ii} = 0. \quad (1)$$

Decomposition into the two helicities $\lambda=+$ and $\times$ gives the quadratic Einstein action. With $v_\lambda = a M_{\text{Pl}} h_\lambda / 2$, each Fourier mode is a canonical parametric oscillator [3].

$$S_t^{(2)} = \frac{M_{\text{Pl}}^2}{8} \sum_{\lambda} \int d\eta \frac{d^3 k}{(2\pi)^3} a^2 (|h'_\lambda|^2 - k^2 |h_\lambda|^2), \quad v_\lambda = \frac{a M_{\text{Pl}}}{2} h_\lambda. \quad (2)$$

$$v_k'' + \left(k^2 - \frac{a''}{a}\right) v_k = 0, \quad v_k \longrightarrow \frac{e^{-ik\eta}}{\sqrt{2k}} \quad (k \gg aH). \quad (3)$$

The term a''/a amplifies a mode when its wavelength exceeds the comoving Hubble radius. In quasi-de Sitter expansion the growing solution makes h_k nearly constant outside the horizon. The factor of two in the dimensionless tensor spectrum counts the two helicities.

B. Spectrum, consistency relation, and energy scale

At a pivot $k^* = 0.05 \text{ Mpc}^{-1}$, let A_s denote the scalar curvature amplitude and define r as the tensor-to-scalar ratio. To first order in the slow-roll parameter $\epsilon = -\dot{H}/H^2$, canonical single-field inflation gives

$$\mathcal{P}_t(k) = \frac{2H_*^2}{\pi^2 M_{\text{Pl}}^2} \left(\frac{k}{k^*}\right)^{n_t}, \quad r = \frac{\mathcal{P}_t(k^*)}{A_s} = 16\epsilon^*, \quad n_t = -2\epsilon^* = -\frac{r}{8}. \quad (4)$$

The last equality is a model-specific consistency relation, not part of the definition of r . If the tensor tilt is fitted independently, a CMB limit applies over the measured wavenumber interval and cannot be extrapolated across many decades without specifying the spectrum. Under the slow-roll relation and potential domination, the Hubble and potential scales follow algebraically:

$$H_* = \pi M_{\text{Pl}} \sqrt{\frac{A_s r}{2}}, \quad V_*^{1/4} = M_{\text{Pl}} \left(\frac{3\pi^2 A_s r}{2}\right)^{1/4}. \quad (5)$$

Because the potential scale varies only as $r^{1/4}$, a large improvement in polarization sensitivity produces a more modest change in the inferred energy scale. The scientific gain is nevertheless substantial: r separates broad classes of inflationary potentials and, with n_s , constrains the trajectory in slow-roll parameter space.

III. FROM TENSOR MODES TO CMB POLARIZATION

Propagation through the thermal history

After inflation, each helicity obeys a damped wave equation. The conformal Hubble rate is $\mathcal{H} = a'/a$ and $\Pi_\lambda T$ is the transverse-traceless anisotropic stress.

$$h_\lambda'' + 2\mathcal{H}h_\lambda' + k^2 h_\lambda = 16\pi G a^2 \Pi_\lambda^T. \quad (6)$$

A mode is nearly frozen for $k \ll \mathcal{H}$ and begins to oscillate and redshift after horizon entry. Free-streaming neutrinos supply a nonlocal anisotropic stress; for modes entering during radiation domination, Weinberg found a 35.6% reduction of the squared tensor amplitude [7]. Equality, recombination, reionization, and late acceleration then determine the transfer function sampled by the CMB. In a purely matter-dominated era with negligible anisotropic stress, the normalized exact solution is

$$T_h(k, \eta) := \frac{h(k, \eta)}{h(k, 0)} = \frac{3j_1(k\eta)}{k\eta}, \quad T_h \rightarrow 1 \quad (k\eta \rightarrow 0). \quad (7)$$

The oscillatory subhorizon decay in (7) is important because polarization responds mainly to the time variation h' at a scattering surface. It also limits the simple identification $\ell \approx k\chi^*$, where χ^* is the comoving distance to last scattering: the observed multipole receives a finite line-of-sight kernel rather than a single Fourier mode.

Spin-2 projection and the two tensor features

Thomson scattering produces linear polarization only when the incident radiation has a local quadrupole. The Stokes fields $Q \pm iU$ transform with spin ± 2 and may be expanded, in one common phase convention, as

$$(Q \pm iU)(\hat{\mathbf{n}}) = \sum_{\ell m} (a_{\ell m}^E \pm ia_{\ell m}^B)_{\pm 2} Y_{\ell m}(\hat{\mathbf{n}}). \quad (8)$$

E has parity $(-1)^l$ and B has parity $(-1)^{l+1}$. Linear scalar perturbations possess no handed pattern around their wavevector and therefore generate E but not B; tensor helicities generate both [4]–[6]. Statistical isotropy makes the angular covariance diagonal in ℓ and m and defines the spectra $C_{\ell XY}$. The tensor B-mode spectrum is a positive projection of the primordial spectrum:

$$C_{\ell}^{BB,t} = 4\pi \int d\ln k P_t(k) [\Delta_{\ell}^B(k)]^2. \quad (9)$$

The transfer function Δ_{ℓ}^B contains the tensor evolution, the visibility function, and spin-weighted radial functions. Recombination produces a broad maximum near $\ell \approx 80$: modes must have evolved enough to create a quadrupole but not so long that their amplitude has decayed away. Reionization creates a second feature at $\ell \approx 10$ because renewed scattering at low redshift sees a large-scale tensor quadrupole. Its height depends on the optical depth τ , making large-scale E modes valuable for controlling the same reionization history.

Table 1. Physical stages connecting the primordial spectrum to CMB B modes

Stage	Scale or epoch	Tensor information	Principal limitation
Horizon exit	$k=aH$ during inflation	Sets $P_t(k)$, r , and n_t	Inflationary dynamics
Superhorizon era	$k \ll \mathcal{H}$	Nearly conserved h	Nonstandard stress/history
Recombination	$z \approx 1090$; $\ell \approx 30$ –150	Main tensor B-mode feature	Dust and lensing B
Reionization	z of order 6–10; $\ell \approx 10$	Large-angle B-mode feature	τ , $1/f$ noise, sky cuts
Late propagation	Structure at $z < \text{last scattering}$	E-to-B lensing conversion	Delensing residual

IV. OBSERVATIONAL INFERENCE

Inference begins with polarized frequency maps and ends with a posterior for the tensor amplitude; foreground separation and lensing removal are part of that estimator, not post-processing details.

A. Multifrequency sky model

A cleaned B-mode map is not directly observed. At each frequency ν , the polarization data contain a frequency-independent CMB field in thermodynamic units, low-frequency synchrotron, high-frequency thermal dust, and noise. A useful local model is

$$\mathbf{d}_{\nu}(\hat{\mathbf{n}}) = \mathbf{s}_{\text{CMB}}(\hat{\mathbf{n}}) + a_s(\nu, \hat{\mathbf{n}}) \mathbf{s}_s(\hat{\mathbf{n}}) + a_d(\nu, \hat{\mathbf{n}}) \mathbf{s}_d(\hat{\mathbf{n}}) + \mathbf{n}_{\nu}. \quad (10)$$

Synchrotron is often approximated by a power law, whereas dust follows a modified blackbody. After conversion from antenna to thermodynamic temperature, their leading spectral factors may be written

$$a_s(\nu) \propto \left(\frac{\nu}{\nu_s}\right)^{\beta_s}, \quad a_d(\nu) \propto \left(\frac{\nu}{\nu_d}\right)^{\beta_d} \frac{B_{\nu_d}(T_d)}{B_{\nu_d}(T_d)}. \quad (11)$$

The indices and polarization angles vary across the sky, so a single rigid scaling can bias r when sensitivity becomes high. Planck measured polarized dust power even in the faintest high-latitude regions and found no window in which foreground subtraction could be omitted [11]. The 2015 joint BICEP2/Keck–Planck analysis detected the 150×353 GHz B-mode correlation, found strong evidence for dust, no significant tensor signal, and $r_{0.05} < 0.12$ [12]. That result established multifrequency coherence—not apparent morphology at one band—as the decisive foreground test.

B. Lensing, delensing, and likelihood construction

Weak lensing remaps the primary polarization by the gradient of the projected potential ϕ . To first order, its flat-sky B mode is a convolution of scalar E modes with ϕ ,

$$B^{\text{lens}}(\ell) = \int \frac{d^2\mathbf{L}}{(2\pi)^2} W(\ell, \mathbf{L}) E(\mathbf{L}) \phi(\ell - \mathbf{L}). \quad (12)$$

Lensing therefore supplies both a cosmological observable and a variance floor for primordial B modes [8]. Quadratic combinations of temperature and polarization reconstruct ϕ [9], permitting a lensing template to be subtracted. SPTpol first detected lensing B modes by cross-correlating an E-mode map with a potential tracer and comparing the template with its measured B map [10]. Delensing efficiency must be propagated into the final covariance; subtracting a noisy template can otherwise exchange lensing variance for reconstruction noise.

Modern analyses fit all auto- and cross-frequency bandpowers. If $\hat{\mathbf{C}}$ is the data vector, $\mathbf{C}(\theta)$ the model containing r , lensing, dust, and synchrotron parameters, and Σ its bandpower covariance, a Gaussian approximation has

$$-2\ln \mathcal{L}(\theta) = [\hat{\mathbf{C}} - \mathbf{C}(\theta)]^T \Sigma^{-1} [\hat{\mathbf{C}} - \mathbf{C}(\theta)] + \ln |\Sigma| + \text{const.} \quad (13)$$

Cut-sky mode coupling, non-Gaussian lensing covariance, calibration, beam uncertainty, and foreground priors complicate this expression in production likelihoods. A robust result is tested against alternate masks, data splits, frequency selections, dust decorrelation models, and null spectra. Polarization-angle error is especially dangerous because it rotates the much larger E field into B; differential gain and beam mismatch can similarly leak temperature anisotropy into polarization [16].

C. Published constraints

Planck 2018 temperature and polarization data constrained inflationary parameter space and, when combined with the then-current BICEP/Keck likelihood, strongly disfavored several large-field monomial potentials [13]. The BK18 analysis combined BICEP/Keck 95, 150, and 220 GHz maps with WMAP and Planck bands spanning 23–353 GHz. Fitting lensed Λ CDM+r+dust+synchrotron gave $\sigma(r)=0.009$ and $r_{0.05}<0.036$ at 95% confidence [14]. A subsequent analysis combined BK18 with Planck PR4 while varying the six Λ CDM parameters; it obtained $r_{0.05}<0.034$, tightening to $r_{0.05}<0.032$ after adding BAO and CMB-lensing data [15]. The latter is a joint cosmological constraint and should not be labeled a B-mode-only bound.




Figure 3. Comparison of selected published 95% limits on $r_{0.05}$. The bars use different data combinations and likelihoods, as identified in the studies [12], [14], [15].

Table 2. Observational landmarks and their inference content

Analysis	Data lever	Published result	Interpretive role
SPTpol, 2013 [10]	E template $\times$ measured B	First detection of lensing B modes	Lensing is measurable and removable

Analysis	Data lever	Published result	Interpretive role
BKP, 2015 [12]	150 GHz $\times$ Planck 353 GHz	Dust detected; $r_{0.05} < 0.12$	Single-band excess was not primordial evidence
Planck 2018 [13]	Full-sky T, E, B and external BK	Restricted r - n_s space	Tensor limits test inflationary potentials
BK18, 2021 [14]	23–353 GHz; deep clean field	$r_{0.05} < 0.036$; $\sigma(r) = 0.009$	Foreground-marginalized B-mode limit
PR4+BK18, 2022 [15]	CMB; then +BAO+lensing	$r_{0.05} < 0.034$; combined < 0.032	τ and Λ CDM uncertainties propagated

No statistically established primordial tensor contribution appears in these likelihoods. The improving limits reflect more than lower detector noise: frequency leverage broke the dust degeneracy, lensing was measured rather than treated as an unknown excess, and joint parameter sampling propagated uncertainty in τ and the scalar sector. The detailed experimental and astrophysical hierarchy is reviewed in [17].

V. ANALYTICAL AND NUMERICAL INSIGHTS

Figure 1 evaluates (4) and (5) with $A_s = 2.10 \times 10^{-9}$, $k^* = 0.05 \text{ Mpc}^{-1}$, and the single-field consistency tilt. The three tensor spectra are separated mainly by normalization because $|n_t| \leq 0.004$ for the plotted values. Across four decades in k , the slow-roll tilt changes the amplitude far less than changing r from 0.032 to 0.001. A CMB measurement of r alone therefore determines an amplitude; testing $n_t = -r/8$ requires a much longer spectral lever arm or additional gravitational-wave observables.




Figure 1. Numerical evaluation of the slow-roll tensor spectrum and its energy-scale conversion. The vertical line in panel (a) marks $k^* = 0.05 \text{ Mpc}^{-1}$; panel (b) marks the combined $r < 0.032$ bound and the $\delta r = 0.001$ mission target.

Figure 2 isolates the propagation physics using the exact matter-era solution (7). The superhorizon plateau preserves the primordial normalization. After $x = k\eta$ crosses unity, the tensor amplitude oscillates with an envelope that decays approximately as x^{-2} , so its power decays approximately as x^{-4} . A visibility surface samples h' during this transition; modes far outside the horizon have too little time variation, while modes deep inside have already been damped. This competition is the analytical origin of the recombination-scale localization before the full spin-projection kernel is applied.




Figure 2. Exact matter-dominated tensor transfer function $T_h = 3j_1(x)/x$. The dashed vertical line marks $x=1$; the left panel shows the power loss across horizon entry and the right panel resolves the oscillatory phase.

Table 3. Inflationary scales implied by selected tensor amplitudes

r	$n_t = -r/8$	$H \square$ (GeV)	$V \square^{1/4}$ (GeV)
0.032	-4.000e-03	4.434e+13	1.368e+16
0.010	-1.250e-03	2.479e+13	1.022e+16
0.001	-1.250e-04	7.839e+12	5.750e+15

At the combined upper limit $r=0.032$, the slow-roll conversion gives $H^* \approx 4.4 \times 10^{13}$ GeV and $V^{*1/4} \approx 1.37 \times 10^{16}$ GeV. At $r=0.001$ the corresponding values are about 7.8×10^{12} GeV and 5.8×10^{15} GeV. These are upper scales under the assumptions entering (4) and (5); an independently tilted spectrum, noncanonical tensor normalization, or non-vacuum tensor source changes the translation even when the observed CMB bandpower is unchanged.

VI. DESIGN LOGIC FOR THE NEXT MEASUREMENTS

The Fisher information illustrates why survey design must balance sky area, angular resolution, and foreground control. For bandpower covariance approximated as Gaussian, a schematic tensor-amplitude information sum is

$$\sigma^{-2}(r) \simeq \sum_{\ell} \frac{(2\ell+1)f_{\text{sky}}}{2} \left[\frac{C_{\ell}^{BB,t}(r=1)}{r C_{\ell}^{BB,t}(r=1) + C_{\ell}^{BB,\text{res}} + N_{\ell}^{BB}} \right]^2. \quad (14)$$

Here $C_{\ell}^{BB,t}$ includes residual lensing and foreground uncertainty after cleaning. Increasing f_{sky} gains large-scale modes and controls sample variance, but it also admits brighter and more spatially complex Galactic emission. Finer resolution improves lensing reconstruction, whereas stable large-scale scanning and low $1/f$ noise preserve the reionization and recombination features. Multiple bands below and above the CMB intensity maximum constrain synchrotron and dust rather than merely increasing nominal CMB depth.

The published Simons Observatory design forecast targets $\sigma(r)=0.003$ with small-aperture observations in six bands from 27 to 280 GHz, aided by a large-aperture lensing map [18]. LiteBIRD is designed for full-sky polarization in 15 bands from 34 to 448 GHz with a total sensitivity near 2.2 $\mu\text{K-arcmin}$; its stated success criterion is $\delta r < 0.001$ for fiducial $r=0$, including systematic-error allocation [19]. These capabilities are complementary: space-based coverage controls the lowest multipoles and broad spectral separation, while high-resolution ground data sharpen lensing reconstruction and delensing.

Multitracer delensing combines internal CMB reconstruction with external tracers of the mass distribution. Forecasts for the Simons Observatory show how tracer correlation and overlap determine the residual lensing power and hence the attainable uncertainty on r . The practical target is not a visually empty B map. It is a likelihood whose residual lensing, foreground complexity, calibration transfer, and selection functions are all constrained tightly enough that a tensor-shaped component is stable under analysis changes.

VII. DISCUSSION

A primordial claim requires several mutually reinforcing signatures. The recovered component must follow the blackbody CMB spectrum in thermodynamic units, exhibit the tensor angular dependence in both the reionization and recombination ranges when accessible, remain stable across masks and foreground models, and pass instrumental null tests. A common r should be consistent with tensor contributions to TT and TE, although those spectra are limited by scalar cosmic variance. If sensitivity permits, TB and EB provide tests of parity violation or polarization-angle error rather than a standard parity-symmetric tensor amplitude.

The pivot scale and model space must accompany every numerical bound. Limits quoted at $k=0.002$ and 0.05 Mpc^{-1} coincide only after assuming a tensor tilt. Likewise, combining B modes with BAO or lensing tightens r by constraining correlated ΛCDM parameters, but the resulting number is conditional on that cosmological model. These distinctions explain the two BK18-related values used here: 0.036 is the collaboration's multifrequency B-mode result, whereas 0.032 is the broader PR4+BK18+BAO+lensing result.

A continuing null result at δr of order 10^{-3} would remove much of the parameter space of simple high-scale, large-field inflation, but it would not test every early-Universe tensor mechanism. Conversely, a detection would determine an amplitude before it determines an origin. Frequency coherence, angular shape, delensing consistency, and cross-experiment calibration would have to establish that origin. Once secured, the value of r would fix H^* through (5) for vacuum tensors and would convert the CMB from a record of density perturbations into evidence for quantized propagating geometry.

VIII. CONCLUSION


The link between primordial gravitational waves and the CMB is mathematically clean at its endpoints and observationally demanding in between. Inflation sets a tensor spectrum proportional to H^2 ; expansion and anisotropic stress determine its transfer; Thomson scattering projects its quadrupole into parity-odd B modes; and a multifrequency, delensed likelihood estimates r . Current published analyses find no significant primordial component and constrain $r_{0.05}$ below 0.032 in the stated combined data model. The next decisive advance will come from joining full-sky large-scale control, broad frequency leverage, precise polarization calibration, and high-resolution delensing. That combination tests both features of the tensor signal and makes the inferred inflationary scale auditable from maps to cosmological parameters.

REFERENCES

- [1] L. P. Grishchuk, "Amplification of gravitational waves in an isotropic universe," *Soviet Physics JETP*, vol. 40, no. 3, pp. 409–415, 1975.
- [2] A. A. Starobinsky, "Spectrum of relict gravitational radiation and the early state of the universe," *JETP Letters*, vol. 30, no. 11, pp. 682–685, 1979.
- [3] V. F. Mukhanov, H. A. Feldman, and R. H. Brandenberger, "Theory of cosmological perturbations," *Physics Reports*, vol. 215, nos. 5–6, pp. 203–333, 1992.
- [4] U. Seljak and M. Zaldarriaga, "Signature of gravity waves in the polarization of the microwave background," *Physical Review Letters*, vol. 78, no. 11, pp. 2054–2057, 1997.
- [5] M. Kamionkowski, A. Kosowsky, and A. Stebbins, "Statistics of cosmic microwave background polarization," *Physical Review D*, vol. 55, no. 12, pp. 7368–7388, 1997.
- [6] M. Zaldarriaga and U. Seljak, "An all-sky analysis of polarization in the microwave background," *Physical Review D*, vol. 55, no. 4, pp. 1830–1840, 1997.
- [7] S. Weinberg, "Damping of tensor modes in cosmology," *Physical Review D*, vol. 69, no. 2, art. no. 023503, 2004.
- [8] L. Knox and Y.-S. Song, "A limit on the detectability of the energy scale of inflation," *Physical Review Letters*, vol. 89, no. 1, art. no. 011303, 2002.
- [9] T. Okamoto and W. Hu, "CMB lensing reconstruction on the full sky," *Physical Review D*, vol. 67, no. 8, art. no. 083002, 2003.
- [10] D. Hanson et al. (SPTpol Collaboration), "Detection of B-mode polarization in the cosmic microwave background with data from the South Pole Telescope," *Physical Review Letters*, vol. 111, no. 14, art. no. 141301, 2013.
- [11] Planck Collaboration, "Planck intermediate results. XXX. The angular power spectrum of polarized dust emission at intermediate and high Galactic latitudes," *Astronomy & Astrophysics*, vol. 586, art. no. A133, 2016.
- [12] BICEP2/Keck and Planck Collaborations, "Joint analysis of BICEP2/Keck Array and Planck data," *Physical Review Letters*, vol. 114, no. 10, art. no. 101301, 2015.
- [13] Planck Collaboration, "Planck 2018 results. X. Constraints on inflation," *Astronomy & Astrophysics*, vol. 641, art. no. A10, 2020.
- [14] P. A. R. Ade et al. (BICEP/Keck Collaboration), "Improved constraints on primordial gravitational waves using Planck, WMAP, and BICEP/Keck observations through the 2018 observing season," *Physical Review Letters*, vol. 127, no. 15, art. no. 151301, 2021.
- [15] M. Tristram et al., "Improved limits on the tensor-to-scalar ratio using BICEP and Planck data," *Physical Review D*, vol. 105, no. 8, art. no. 083524, 2022.



- [16] W. Hu, M. M. Hedman, and M. Zaldarriaga, "Benchmark parameters for CMB polarization experiments," *Physical Review D*, vol. 67, no. 4, art. no. 043004, 2003..
- [17] M. Kamionkowski and E. D. Kovetz, "The quest for B modes from inflationary gravitational waves," *Annual Review of Astronomy and Astrophysics*, vol. 54, pp. 227–269, 2016.
- [18] Simons Observatory Collaboration, "The Simons Observatory: Science goals and forecasts," *Journal of Cosmology and Astroparticle Physics*, no. 02, art. no. 056, 2019.
- [19] LiteBIRD Collaboration, "Probing cosmic inflation with the LiteBIRD cosmic microwave background polarization survey," *Progress of Theoretical and Experimental Physics*, vol. 2023, no. 4, art. no. 042F01, 20230.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)