



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 12 **Issue:** XII **Month of publication:** December 2024

DOI: <https://doi.org/10.22214/ijraset.2024.65876>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

Detection of Carbohydrates, Fats, and Proteins in Foodstuffs

Falguni Bhavsar¹, Tanuja Babaleshwar², Divya Bobade³, Arvind Done⁴, Trushal Katke⁵, Vinay Gaikwad⁶

Department of Applied Sciences and Humanities, Pimpri Chinchwad college of Engineering, Pune

Abstract: The detection of carbohydrates, fats, and proteins in foodstuffs is crucial for understanding their nutritional content and quality. This analysis involves the use of simple chemical tests that produce observable changes, such as color changes, indicating the presence of these macronutrients. Carbohydrates are detected using tests like the Benedict's test for reducing sugars and the iodine test for starch. Fats are identified by the grease spot test or Sudan III dye test, while proteins are detected through the Biuret test. These methods are cost-effective, easy to perform, and essential in food science, health, and nutrition fields.

Keywords: Detection methods; Fats, proteins, carbohydrates; Nutritional composition; Food testing

I. INTRODUCTION

Food is one of the basic needs of all living organisms. Food contains various nutrients such as carbohydrates, proteins, fats etc. To detect these nutrients in food is crucial for understanding their nutritional value of food. The process of identifying these substances involves specific chemical tests. By performing these tests on food samples, we can confirm which nutrients are present and understand the food composition.

Carbohydrates are the primary source of energy. They can be found in various forms, such as simple sugars (Glucose, Sucrose) and complex carbohydrates (Starch). The presence of carbohydrates in food is typically detected through Iodine Tests or Benedict's Test. Fats play a crucial role in energy storage and insulation and cellular function. The detection of fats is by Emulsion Test.

Proteins are vital for growth, tissue repair, and enzyme function. Proteins in food can be detected by Biuret's Test.

By performing these tests, we can detect and analyze the composition of foodstuffs. This knowledge is important not only for dietary purposes but also for food manufacturing and quality control. In this project, we'll explore the chemical tests used for the detection of carbohydrates, proteins, and fats.

II. METHODOLOGY

A series of qualitative and quantitative tests were conducted to confirm the presence of carbohydrates, proteins, and fats. The reaction was observed for any colour changes or precipitate formation, indicating the presence of the compound.

1) Cheese (Detection of Fat):



Fig 1: Detection of fats in cheese

a) *Materials Required:*

- 10 g cheese sample
- Ethanol as a fat solvent
- Glass beaker or test tube
- Analytical balance
- Filter paper
- Hot water bath or Bunsen burner
- Measuring cylinder
- Funnel

b) *Procedure:*

- Preparation of Cheese Sample: Weigh exactly 10 g of cheese using a balance. Grind the cheese to a fine consistency.
- Adding Solvent: Transfer the ground cheese into a beaker or test tube. Add 20-30 mL of ethanol as the solvent. Stir the mixture thoroughly for 5-10 minutes to allow the fat to dissolve in the solvent.
- Separation of Fat-Solvent Solution: After stirring, let the mixture stand to allow the fat to separate from any undissolved material. Place filter paper in a funnel and pour the solution through it to separate the liquid from the solids. Collect the filtered solution in a clean, beaker.
- Evaporation of Solvent: Place the beaker with the fat-solvent solution in a hot water bath (around 70-80°C) to evaporate the solvent. Alternatively, a Bunsen burner with gentle heat can be used. Continue heating until the solvent has completely evaporated, leaving only the fat residue in the beaker. This may take 15-20 minutes.

c) *Result:*

The left-over residue in beaker is fat content in cheese.

2) *Egg Yolk (Detection of Protein):*

a) *Materials Required:*

- egg yolk (approximately 5-10 g)
- Ethanol
- Test tubes
- Pipette or dropper
- Distilled water
- Measuring cylinder
- Stirring rod or glass rod

b) *Procedure:*

- Preparation of Egg Yolk: Take 5-10 g of egg yolk. You can separate the yolk from the egg white and gently mash or homogenize the yolk. Weigh the sample precisely using an analytical balance.




Fig 2: Egg yolk

- Adding Ethanol: Add ethanol to prepared egg yolk and stir well then keep aside for a while.




Fig 3: Mixing of ethanol with egg yolk

- c) *Observation:* The egg yolk precipitate at bottom of beaker.
 d) *Result:* The precipitation of egg yolk indicates that protein is present in egg yolk.

3) Egg Yolk (Detection of Fat):

a) Materials Required:

- Egg (sample)
- Solvent: Ethanol
- Beaker
- Filter paper
- Rotary evaporator or water bath
- Analytical balance

b) Procedure:

Sample Preparation: Separate the egg yolk from the egg white, as fat is primarily present in the yolk. Weigh a specific amount of yolk (e.g 5 grams) using an analytical balance.

Fat Extraction: Add 20 mL of the solvent (ethanol) to the yolk in a beaker. Stir or shake the mixture thoroughly for 10-15 minutes to extract the fat. Filter the mixture through filter paper to separate the liquid fat extract.

Solvent Evaporation: Transfer the filtrate to a pre-weighed container. Use a rotary evaporator or place the container in a water bath at a low temperature to evaporate the solvent completely. Ensure only the fat residue remains in the container.

Observation and Result: The fat content in the egg yolk. This method allows the determination of fat in eggs.


4) Potato (Detection of Carbohydrates):

a) Materials Required:

- Fresh potato (sample)
- Distilled water
- Iodine solution
- Test tubes or small beakers
- Knife (to cut the potato)
- Dropper



a)



b)

Fig 4: a) Potato b) change in colour of potato to blue - violet colour it confirms that potato has carbohydrates (starch)

Conclusion: A blue-black color indicates the presence of starch (carbohydrate) in the potato sample.

This iodine test is a simple and reliable qualitative method for detecting starch in various plant tissues, including potatoes.

Detection of Carbohydrates using this method can be used for both Potato and Bread mention below in fig 4



Fig 5: Detection of carbohydrates in bread

5) Oil (Detection of Fat):

a) Materials Required:

- Oil (sample)
- Ethanol
- Distilled water
- Test tube or small beaker
- Pipette




Fig: 6 Cooking oil

b) *Procedure:* Sample Preparation: Take a small amount of the oil sample (about 2-3 mL) in a clean test tube or beaker.

Dissolving the Oil: Add 3-4 mL of ethanol to the test tube containing the oil sample. Shake the test tube gently to ensure that the oil dissolves completely in the ethanol. Ethanol is a solvent for oils and fats, so it should dissolve the fat easily.

Adding Water: Slowly add a few mL of distilled water to the ethanol-oil solution. Shake the mixture gently.

Observation: If fat is present in the oil, the mixture will turn cloudy or milky. This is due to the fact that the water causes the fat (which was dissolved in ethanol) to precipitate out as small droplets.

Conclusion: The appearance of a cloudy or milky solution upon adding water indicates the presence of fat in the oil sample. This simple ethanol test is often used for confirming the presence of lipids in oils and other fatty substances.

6) Gram flour (Detection of Protein):

a) Materials Required:

- Gram flour (approximately 5-10 g)
- Copper Sulphate (CuSO_4) solution
- Sodium hydroxide (NaOH) solution
- Test tubes
- Measuring cylinder
- Stirring rod or glass rod
- Distilled water
- Pipette or dropper
- Analytical balance (for precise measurements)



Fig 6: Colour change of gram flour indicates presence of proteins

b) Procedure:

- Preparation of Gram Flour Sample: Take 5-10 g of gram flour. Weigh the sample accurately using an analytical balance. Place the gram flour in a clean test tube.
- Dissolving Gram Flour in Water: Add 10-15 mL of distilled water to the test tube containing the gram flour. Stir the mixture gently to dissolve the flour. You might need to shake or stir it for a minute to form a consistent paste or suspension.

Adding Copper Sulphate (CuSO_4) Solution: After preparing the gram flour suspension, add 2-3 drops of copper Sulphate (CuSO_4) solution to the test tube.

Adding Sodium Hydroxide (NaOH) Solution: Add 2-3 mL of sodium hydroxide (NaOH) solution to the test tube. Sodium hydroxide will alkalize the solution, and if proteins are present, it will cause a color change when combined with copper ions.

Observation of Results: Mix the solution gently. If proteins are present in the gram flour, you will observe a violet or purple color forming in the solution. This color change is due to the formation of a complex between the copper ions and the peptide bonds of proteins.

c) Result:

In this research study, the detection and quantification of fat, protein, carbohydrates in potato, egg yolk, oil, gram flour, cheese, and bread were successfully carried out using Biuret test for protein, iodine test for starch, ethanol test for fat. The results observed overall are mentioned below in table 1.

Table 1

Food content	Chemical name	Change	Nutrient
Potato	Iodine	Colour changes to violet black	Carbohydrate
Egg yolk	Ethanol	It forms precipitate	Protein

Egg yolk	Ethanol	After heating it forms residue	Fat
Cheese	Ethanol	After heating it forms residue	Fat
Gram flour	CuSO ₄ (Copper sulphate) NaOH (Sodium Hydroxide)	Colour changes to violet	Protein
Oil	Ethanol	Solution turns milky or cloudy	Fat

III. CONCLUSIONS

The detection of carbohydrates, proteins, and fats in foodstuffs involves specific chemical tests that produce distinct reactions or colour changes. These tests, such as Molisch's for carbohydrates, Biuret for proteins, and Sudan III or grease spot test for fats, are simple yet effective in identifying the presence of these macronutrients. By employing these methods, we can qualitatively analyse food samples to better understand their nutritional composition, which is essential for dietary planning and scientific studies.

IV. ACKNOWLEDGMENT

We are thankful to Pimpri Chinchwad college of engineering nigdi Pune for providing chemistry lab with chemicals, apparatus for completion of experimental work. Also expressing our gratitude to Falguni S. Bhavsar mam for necessary guidance during the journey of research.

REFERENCES

- [1] G. It, WILSTER, W. V. PRICE, A. J. MORRIS, E. F. GOSS, G. P. SANDERS, Determination of fat, moisture and salt in soft cheese, JOURNAL OF DAIRY SCIENCE, VOLUME XXIII MARCH, 1940 NUMBER 3, June 22, 1939, DOI:[10.3168/jds.2017-13431](https://doi.org/10.3168/jds.2017-13431)
- [2] To Determine the Fat content of Chicken Egg Yolk Rida Amir, Hira Iftikhar , Aqsa Amjad, GSJ: Volume 9, Issue 8, August 2021, Online: ISSN 2320-9186 www.globalscientificjournal.com
- [3] SUNEETA PANICKER*, ANAM HAMDULE, ANALYSIS OF MEDICINAL PROPERTIES OF VIGNA ACONITIFOLIA (MATKI), Vol 14, Issue 5, 2021, DOI:[10.22159/ajpcr.2021.v14i5.41186](https://doi.org/10.22159/ajpcr.2021.v14i5.41186)
- [4] Guang C, Phillips R. Plant food-derived angiotensin I converting enzyme inhibitory peptides. J Agric Food Chem 2009;57:5113-20.
- [5] Study the Physical and Functional Properties of Chickpea and Black Gram Flours Nikhil D. Solanke1, Pradeep P. Thorat2* and Jayashri Ughade3, ISSN: 2319-7706 Volume 10 Number 10 (2021) Journal homepage: <http://www.ijcmas.com>



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)