



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 12 **Issue:** II **Month of publication:** February 2024

DOI: <https://doi.org/10.22214/ijraset.2024.58380>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

Determination of Residual Presence of Toxic Heavy Metal in the Liver of *LABEO BATA* (HAMILTON, 1822) Using Rain Water Harvest System

Sanjay Kumar Paul¹, Ipsita Jena², Kriti Dhawan³, Triparna Chatterjee⁴, Priti Mukherjee⁵, Samik Acharjee⁶, Bidisha Maitra Sen⁷, Santi Ranjan Dey⁸

^{1, 6, 8} Assistant Professor, Department of Zoology, Rammohan College, Kolkata

^{2, 3, 4, 5} Graduate Student, Department of Zoology, Rammohan College, Kolkata

⁷ Assistant Professor, Industrial Fish and Fisheries Department, Asutosh College, Kolkata

Abstract: Fish are special among vertebrates because they may absorb metal from the water through their gills and from their food through their guts (direct and trophic absorption methods). Since the fish's gills are the primary organ targeted by metal toxicity, the direct uptake route is more significant. Most research on the toxicity and accumulation of metals in fish focuses on individual metals. But in the natural world, fish are exposed to mixes of many metals, whose toxicity levels typically differ from those of individual metals because of their additive, sometimes less than additive (antagonistic) effects, or higher than additive (synergistic) effects. This article discusses the build-up of heavy metals in fish organs and tissues, particularly in relation to the intake of water-borne metals via the trophic pathway. The trophic route is important because it is related with "Biological Magnification". The fish we eat, usually comes from rivers or pond or bheri, in Kolkata. These are also contaminated with heavy metal toxicity. The rain water is contamination free from heavy metal. We collected live fish from market and kept them in "Rain Water Harvest System" for 24 hours. The accumulated heavy metal was expected to be eliminated from fish liver within 24 hours through the process of detoxification. We found that, even after 24 hours, fish liver contain certain heavy metal as residue. The model fish was *Labeo bata* (Hamilton, 1822). Far more studies on metal interactions should be performed to reach the level of predictability.

Keywords: *Labeo bata*, Heavy metal, Rain Water, Residual presence, Liver

I. INTRODUCTION

Because of their toxicity, buildup, and amplification in aquatic animals, heavy metals represent a significant concern to the aquatic environment (Komjarova and Blast, 2009). Due to their toxicity, persistence, and capacity for bioaccumulation, the majority of them (Zn, Cu, Ni, Cr, Pb, Cd, Hg, etc.) are listed as priority hazardous substances (pollutants) in several nations (Clearwater et al, 2002, Bury et al 2003, Chapman et al, 2003). It appears that strong binding to metallothioneins or other metal-binding proteins is how heavy metals accumulate in fish's gills, liver, kidneys, and muscle (Klavins et al 2009). Over the past several decades, researchers have focused a lot of emphasis on the bioaccumulation of heavy metals and other contaminants in fish. As a result, a considerable amount of field and experimental data have been compiled and reviewed (Kamunde and McPhile, 2011). Two pathways of uptake are involved in the bioaccumulation process in fish: the aqueous uptake of chemicals carried by the water by the gills (bioconcentration) and the dietary uptake of contaminated food and sediment by the intestinal mucosa (biomagnification). Therefore, intricate interactions between the aforementioned pathways and different elimination strategies (excretion, passive release, and metabolization) lead to bioaccumulation (Ergolu et al, 2005). However, under a technical definition of the word, metals themselves cannot be metabolised; instead, they can only be excreted from the body (Ghosh et al, 2007). Generally speaking, vertebrates only have two ways to acquire metal: directly from the food through trophic channels, such as gills, and indirectly from the water through the diet (Farkas et al, 2003).

II. MATERIALS AND METHODS

The "Rain Water Harvest System" has been introduced in 2020-21 at Rammohan College ((22.582952°N & 88.370997°E), used for this experiment. 5 batch of 10 fish each (Weight average 30 gm) of *Labeo bata* (Hamilton, 1822) were procured from "Rajendrapur Fish Seed Market" Naihati, 24 Parganas(N) and kept in the tank of "Rain water" for 24 Hours. No food was provided to the fish.

After 24 hours, the liver of fish were collected (as detoxification organ) and dissolved in HCl for ICPMS to Director of Quality Control (Food), Government of West Bengal.

Inductively coupled plasma (ICP) mass spectrometry (MS) finds frequent application in a wide range of research domains, including the food, material, chemical, semiconductor, and nuclear industries, as well as the earth, environmental, biological, and forensic sciences. All kinds of samples and matrices introduced by a range of specialised equipment find a suitable atomizer and element ioniser in the high ion density and high temperature of a plasma. Prominent characteristics include maximum sensitivity (ppt–ppq), relative salt resistance, element response independent of compound, and optimal quantitation precision contribute to ICP MS's unrivalled ability to effectively detect, identify, and consistently quantify trace elements. The molecular identification capabilities of ICP MS hyphenated to species-specific separation techniques are expanded by the growing availability of pertinent reference compounds and good separation selectivity. ICP MS is a productive and extremely sensitive technique for target-element oriented discoveries of relevant and unknown compounds, whereas molecular ion source MS is specialised in figuring out the structure of unknown molecules.



Rain Water Harvest System & Fish Tank


Labeo bata (Hamilton, 1822)

III. RESULT

Table 1. $\mu\text{g/gm}$ liver tissue sample

Cr	0.11 ± 0.02
Cu	2.91 ± 0.009
Zn	10.8 ± 0.07
As	0.07 ± 0.0003
Cd	0.09 ± 0.003
Sn	0.08 ± 0.003
Hg	0.04 ± 0.001
Pb	0.27 ± 0.008

IV. DISCUSSION

This is very important finding that, all industrial and non industrial heavy metal and metalloid are present in the liver of an edible fish. Liver is the detoxification organ, through which digested materials circulated in the body. So it is obvious that, heavy metal can be found in the muscles and bone of fish. Even after 24 hours of exposure of metal free water, these heavy metals were neither excreted nor diffused.

An extensive multifactorial experimental study must be carried out using representative multimetal mixtures in various combinations at fluctuating environmentally relevant metal concentrations, a wide variety of fish species (including commercially important and bioindicator species with varying ecological status), and meaningful organs and tissues for environmental risk assessment that may be used as a foundation for the creation of ecotoxicologically relevant water quality criteria for metals.

REFERENCES

- [1] Bury, N. R., P. A. Walker, and C. N. Glover. "Nutritive Metal Uptake in Teleost Fish." *Journal of Experimental Biology* 206: 11–23.2003.

- [2] Chapman, P. M., F. Wang, C. R. Janssen, R. R. Goulet, and C. N. Kamunde. "Conducting Ecological Risk Assess- ments of Inorganic Metals and Metalloids: Current Status." *Human and Ecological Risk Assessment: An International Journal* 9: 641–697.2003.
- [3] Clearwater, S. J., A. M. Farag, and J. S. Meyer. "Bio- availability and Toxicity of Dietborne Copper and Zinc to Fish." *Comparative Biochemistry and Physiology* 132: 269–313.2002.
- [4] Eroglu, K., G. Atli, and M. Canli. "Effects of Metal (Cd, Cu, Zn) Interactions on the Profiles of Metallothionein-like Proteins in the Nile Fish *Oreochromis niloticus*." *Bulletin of Environmental Contamination and Toxicology* 75: 390– 399. 2005.
- [5] Farkas, A., J. Salánki, and A. Specziár. 2003. "Age- and Size-specific Patterns of Heavy Metals in the Organs of Freshwater Fish *Abramis brama* L. Populating a Low- contaminated Site." *Water Research* 37 (5): 959–964.2003.
- [6] Ghosh, L., S. Adhikari, and S. Ayyappan. 2007. "Assessment of Toxic Interactions of Heavy Metals and Their Effects on Accumulation in Tissues of Freshwater Fish." *Research Journal of Environmental Toxicology* 1: 37–44.2007.
- [7] Kamunde, C., and R. MacPhail. "Subcellular Interac- tions of Dietary Cadmium, Copper and Zinc in Rainbow Trout (*Oncorhynchus mykiss*)." *Aquatic Toxicology* 105: 518–527.2011.
- [8] Kamunde, C., and R. MacPhail. "Metal–Metal Interac- tions of Dietary Cadmium, Copper and Zinc in Rainbow Trout *Oncorhynchus mykiss*." *Ecotoxicology and Environ- mental Safety* 74: 658–667.2011.
- [9] Klavins, M., O. Potapovics, and V. Rodinov. "Heavy Metals in Fish from Lakes in Latvia: Concentrations and Trends of Changes." *Bulletin of Environmental Contamina- tion and Toxicology* 82: 96–100. 2009.
- [10] Komjarova, I., and R. Blust. "Multimetal Interactions between Cd, Cu, Ni, Pb, and Zn Uptake from Water in the Zebrafish *Danio rerio*." *Environmental Science and Tech- nology* 43 (19): 7225–7229.2009.
- [11] Komjarova, I., and R. Blust. "Effects of Na, Ca, and PH on the Simultaneous Uptake of Cd, Cu, Ni, Pb, and Zn in the Zebrafish *Danio rerio*: A Stable Isotope Experi- ment." *Environmental Science and Technology* 43 (20): 7958–7963.2009.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)