



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 13 Issue: VI Month of publication: June 2025

DOI: https://doi.org/10.22214/ijraset.2025.72270

www.ijraset.com

Call: 🕥 08813907089 🔰 E-mail ID: ijraset@gmail.com



Development and Validation of UV-Visible Spectrophotometric Method for the Analysis of Cefepime in Pharmaceutical Formulations

Dr. P. Suguna

Lecturer in Chemistry, S.G.S Arts College (A) T.T.D, Tirupati, A.P., India

Abstract: A simple, sensitive, selective rapid spectrophotometric method has been developed for the determination of Cefepime in pure form and pharmaceutical formulations based on the oxidative coupling reaction with MBTH reagent, at P^{H} -4.0 which is extractable at 610 nm. Beer's law is obeyed in the concentration ranges 5-30 µg m Γ^{1} . The developed method was applied directly and easily for the analysis of the Pharmaceutical formulations. R.S.D was found to be 0.2604% and Recovery 98.65% respectively. The method was completely validated and proven to be rugged. The interferences of the other ingredients and excipients were not observed. The repeatability and the performance of the proved method were established by point and internal hypothesis and through recovery studies.

Keywords: Spectrophotometry, Cefepime, Oxidative coupling reaction, MBTH / FeCl₃

I. INTRODUCTION

Cefepime is chemically 6R,7R,Z)-7-(2-(2-aminothiazol-4yl(methoxyimino)acetamido)-3-((1-methylpyrrolidinium-1yl)methyle)-8oxo-5-thia-1-aza-bicyclo(4.2.0)oct-2-ene-2-carboxylate. Cefepime is a fourth-generation <u>cephalosporin</u> <u>antibiotic</u>. Cefepime has an extended spectrum of activity against <u>Gram-positive</u> and <u>Gram-negative</u> <u>bacteria</u>, with greater activity against both types of organism than third-generation agents. Cefepime injection is used to treat bacterial infections in many different parts of the body. It belongs to the class of medicines known as cephalosporin antibiotics. It works by killing bacteria or preventing their growth. However, this medicine will not work for colds, flu, or other virus infections. The empericala formula of Cefepime is $C_{19}H_{24}N_6O_5S_2$ and the molecular weight is 480.54grams. It has the following structure :





There is however no reported UV- Visible spectrophotometric method for the analysis of Cefepime in its technical grade and formulations. UV- Visible spectrophotometric method for the quantitative determination of Cefepime. Functional group used for color development of Cefepime was primary amine group. The results obtained in this method was based on complex formation reaction of Cefepime with Oxidative coupling reaction with MBTH .

In the present study an attempt has been made to develop simple UV-Visible spectrophotometric method for quantitative estimation of Cefepime in its technical grade, formulations and biological sample (Blood). The functional group used for the color development of Cefepime was primary amine. The result obtain in this method was based on coupling reaction formation reaction of Cefepime with MBTH/ Fecl₃. An attempt has been made to develop and validate to ensure their accuracy, precision, repeatability, reproducibility and other analytical method validation parameters as mentioned in various gradients.



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 13 Issue VI June 2025- Available at www.ijraset.com

II. MATERIALS AND METHODS

- A. Preparation Of Standard Calibration Curve Of Pure Drug
- 1) Solvent: Dimethyle Sulfoxide was used as Solvent.
- 2) Preparation of standard stock solution: Accurately weighed 100 mg of Cefexime was dissolved in 40 ml of Dimethyle Sulfoxide 1 in 100 ml volumetric flask and volume was made up to the mark with Dimethyle Sulfoxide . i.e. 1000 µg ml-1 (Stock solution A) From the above stock solution A 10 ml of solution was pipette out into 100 ml volumetric flask and the volume was made up to the mark with Dimethyle Sulfoxide to obtained the final concentration of 100 µg ml-1 (Stocksolution B)
- 3) Preparation of Calibration curve: Fresh aliquots of Cefepime ranging from 0.5 to 3ml were transferred into a series of 10 ml volumetric flasks to provide final concentration range of 5 to 30 μg ml⁻¹. To each flask 1ml of (0.01M) MBTH solution was added followed by 1ml of (0.7%) Ferric chloride solution and resulting solution was heated for 15 min and finally 1ml (0.5N) Hydrochloric acid solution was added. The solutions were cooled at room temperature and made up to mark with Dimethyle Sulfoxide. The absorbance of orange red colored chromogen was measured at 610 nm against the reagent blank. The color species was stable for 24 h. The amount of Cefepime present in the sample solution was computed from its calibration curve.
- 4) Procedure for formulations: Twenty tablets containing Cefepime were weighed and finely powdered. An accurately weighed portion of the powder equivalent to 100 mg of Cefepime was dissolved in a 100 ml of Dimethyle Sulfoxide and mixed for about 5 min and then filtered. The Dimethyle Sulfoxide was evaporated to dryness. The remaining portion of solution was diluted in a 100 ml volumetric flask to the volume with Dimethyle Sulfoxide up to 100 ml to get the stock solution A. 10 ml of aliquots was pipette out into 100 ml volumetric flask and the volume was made up to the mark with Dimethyle Sulfoxide to obtained the final concentration of $100 \mu g ml^{-1}$ (Stock solution). Subsequent dilutions of this solution were made with Dimethyle Sulfoxide to get concentration of 5 to 30 $\mu g ml^{-1}$ and were prepared as above and analyzed at the selected wavelength, 610 nm and the results were statistically validated.
- 5) Procedure for Blood sample: After collection of Blood sample it will be centrifuged. For isolation of Cefepime from plasma sample, Dimethyle Sulfoxide was used for protein precipitation. Liquid- Liquid extraction was performed with plasma by alkalinization with 1M NaOH, followed by extraction with 30% dichloromethane in Hexane. The upper organic layer was evaporated to dryness, the dry residue 100 mg was dissolved in 100 ml of Dimethyle Sulfoxide (1000 μ g ml⁻¹). From the above solution 10 ml is taken into a 100 ml of Volumetric flask and made up to the mark with Dimethyle Sulfoxide .(100 μ g ml⁻¹)

From the above solution ranging from 0.5-3 ml (5-30 μ g ml⁻¹) were transferred in to 10 ml Volumetric flask and to the each flask 1ml of (0.01M%) MBTH solution was added followed by 1ml of (0.7%) Ferric chloride solution and made up to the mark with Dimethyle Sulfoxide. Then the resulting solution was heated and finally 1ml (0.5N) Hydrochloric acid solution was added. The solutions were cooled at room temperature and made up to the mark with Dimethyle Sulfoxide. The absorbance of orange red colored chromogen was measured at 610 nm against the reagent blank. The color species was stable for 24 h. The amount of Cefepime present in the sample solution was computed from its calibration curve.









Fig-2: Beer's law plot of Cefepime With MBTH/FeCl3

Fig-3: Beer's law plot for MBTH in Blood sample



Fig-4: A Schematic reaction Mechanism of Cefepime With MBTH





ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538

Volume 13 Issue VI June 2025- Available at www.ijraset.com

Parameter	Visible method
Color	Green
Absorption maxima (nm)	610
Beer's law limits ($\mu g \text{ ml}^{-1}$)	5-30
Molar absorptivity (1 mol ⁻¹ cm ⁻¹)	5.263×10 ⁴
Sandell's Sensitivity (µg cm ⁻²)	0.0452
Regression equation (Y*)	
Slope (b)	0.022
Intercept(a)	0.001
Standard deviation(SD)	0.001
Correlation coefficient (r ²)	0.9999
%RSD (Relative Standard deviation)	0.2604
Range of errors	
Confidence limits with 0.05 level	0.0008
Confidence limits with 0.01 level	0.0010
Limits of detection (LOD)(µg ml ⁻¹)	0.1363
Limits of quantification (LOQ) (µg ml ⁻¹)	0.4545

Table-1.1: Optical characteristics a	and precision by MBTH
--------------------------------------	-----------------------

RSD of 6 independent determinations

Table-1.2: Assay results of Cefepime in formulations by visible Method

Name of the	Formulation in	Amount found by the	Amount found by the	% Recovery
Formulation	(mg)	proposed method (mg)	reference method (mg)	
CEFIXIME		249.56		
	250	t=0.0029	246.25	98.65
		F=8.5897		
		249.98		
MAXIPIME	250	t=0.0028	245.5	98.17
		F=8.5796		

t and F- values refer to comparison of the proposed method with reference method. Theoretical values at 95% confidence limits t= 0.00297 and F= 7.6177

Amount of CEF in	Amount of Standard	Total amount	%
formulation	CEF added (mg)	Found (mg)	Recovery
(mg)			
249.16	200	448.48	99.66
247.91	200	446.23	99.16
248.64	200	447.55	99.45
248.16	250	496.32	99.26
249.30	250	498.6	99.72
248.32	250	496.64	99.32
248.47	300	546.63	99.38
248.83	300	547.42	99.53
247.91	300	545.40	99.16

Table-1.3: Determination of accuracy of Cefepime



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538

Volume 13 Issue VI June 2025- Available at www.ijraset.com

	5	
Total amount found (mean)	Standard deviation	% RSD
248.57	0.6279	0.2526
248.59	0.6171	0.2482
248.40	0.4636	0.1866

Table-1.4: Statistical data for accuracy determination

The results are the mean of three readings at each level of recovery.

			r	.j		
Conc. (µg	Abs 1	Abs2	Abs3	Mean	Std. deviation	(%)RSD
ml^{-1})						
5	0.108	0.109	0.105	0.107	0.0002	0.1869
10	0.219	0.218	0.212	0.216	0.0003	0.1388
15	0.329	0.326	0.327	0.327	0.0001	0.0305
20	0.439	0.437	0.439	0.438	0.0001	0.0228
25	0.549	0.546	0.543	0.546	0.0003	0.0549
30	0.659	0.654	0.653	0.655	0.0003	0.0458
			<u> </u>			

Table-1.5: Repeatability data for CEF at 610 nm

Average of six determinations.

Table-1.6: Color stability data for MBTH Method

Conc. in µg ml ⁻¹				Time ir	n Hours			
20	4	8	12	16	20	24	28	32
	0.439	0.439	0.439	0.440	0.441	0.441	0.398	0.295

Table-1.7. Assay results of Celephine in blood sample				
Name of the	Formulation	Amount found by	Amount found by	% of
Formulation	in (mg)	the proposed	the reference	Recovery
		method in (mg)	method (mg)	
CEFIXIME		2.9		
	5	t=0.0029	2.85	95.78
		F=0.0004		
		2.99		
MAXIPIME	5	t=0.0027	2.86	95.45
		F=0.00039		

Table-1.7: Assay results of Cefepime in Blood sample

tand F values refer to comparison of the proposed method with reference method. Theoretical values at 95% confidence limits t=0.00796 and F=0.0029.

Table-1.8: Determination of accuracy of Celeph	nme
--	-----

		-	1	
Name of the Formulation in	Amount of Drug	Amount of	Total amount	%
(mg)	in Blood sample	Standard Drug	found (mg)	Recovery
	(mg)	added in (mg)		
5	2.99	5	5.98	59.8
5	2.98	5	5.96	59.6

The results are the mean of two readings at each level of recovery.



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 13 Issue VI June 2025- Available at www.ijraset.com

Concentration in Abs1 Abs2 Abs3 Mean Std. Deviation (%) RSD µg ml⁻¹ 0.4 0.052 0.051 0.052 0.051 0.0005 0.9803 0.0015 0.8 0.105 0.104 0.102 0.103 0.9708 1.2 0.158 0.156 0.156 0.156 0.0011 0.7051 1.6 0.211 0.213 0.219 0.214 0.0004 0.1869 2.0 0.264 0.267 0.263 0.264 0.0020 0.7575 2.4 0.0015 0.316 0.319 0.317 0.317 0.4731

Table-1.9: Repeatability data for Cefepime at 610nm

Average of six determinations

III. RESULTS AND DISCUSSION

A. Optical Parameters

In order to ascertain the optimum wavelength of maximum absorption (λ_{max}) formed in UV spectrophotometric method and of the colored species formed in each so the visible spetrophotometric method, specified amount of Cefepime in solution 5-30 µg ml⁻¹ were taken and the colors were developed following the above mentioned procedures individually. The absorption spectra were scanned on spectrophotometer in the wavelength region of 380-800 nm against corresponding reagent blank. The regent blank absorption spectrum of each method was also recorded against distilled water / Dimethyle Sulfoxide. The results are graphically represented in (fig- 1).

B. Parameters fixation

In developing these methods, a systematic study of the effects of various relevant parameters in the methods concerned were under taken by verifying one parameter at a time and controlling all other parameter to get the maximum color development MBTH Method reproducibility and reasonable period of stability of final colored species formed. The following studies were conducted.

Method: The results obtained in this method were based on oxidation followed by coupling reaction of Cefepime with MBTH, Ferric chloride and Orthophosphoric acid to form an green colored chromogen that exhibited maximum absorption at 610 nm against the corresponding reagent blank. The functional group used for the color development for this method was primary amine group. A schematic reaction mechanism of Cefepime with MBTH reagent was shown in (fig-4). The effect of various parameters such as concentration and volume of MBTH and strength of acid order of addition of reagents, solvent for final dilution were studied by means of control experiments varying one parameters at a time.

C. Optical Characteristics

The reference method adhere to beer's law the absorbance at appropriate wave length of a set of solutions contains different amounts of Cefepime and specified amount of reagents (as described in the recommended procedure) were noted against appropriate reagent blank. The beers law plot of the system illustrated graphically (fig: 2) least square regression analysis was carried out for the slope. Intercept and Correlation Coefficient. Beer's law limits, Molar absorptivity & Sandells sensitivity for Cefepime with each of mentioned reagents was calculated. The optical characteristics were present in the table- 1.1. In order to test whether the colored species formed in the method adhere the beer's law the absorbance at appropriate wavelength of a set of solutions contain different amounts of Cefepime and specified amount of reagents (as described in the recommended procedure) were noted against appropriate reagent blanks or distilled water. The beers law plots of the system illustrated graphically (fig -2 & 3) least square regression analysis was carried out for the slope, intercept and correlation coefficient, beer's law limits molar absorptivity Sandells sensitivity for Cefepime with each of mentioned reagents (as described in the recommended procedure) were noted against appropriate reagent blanks or distilled water. The beers law plots of the system illustrated graphically (fig -2 & 3) least square regression analysis was carried out for the slope, intercept and correlation coefficient, beer's law limits molar absorptivity Sandells sensitivity for Cefepime with each of mentioned reagents were calculated. The optical characteristics are presented in the table – 1.1.

D. Precision

The precision of each one among the five proposed spectrophotometric methods were ascertained separately from the absorbance values obtain by actual determination of a fixed amount of Cefepime n (10, 5 μ g ml⁻¹ respectively) in final solution. The percent relative standard deviation and percent range of error (at 0.05 and 0.01 confidence limits) were calculated for the proposed methods and presented in tables – 1.1.



E. Analysis of formulations

Commercial formulations of Cefepime were successfully analyzed by the proposed methods. The values obtained from the proposed and reference methods were compared statistically by the t and F tests and were found that those proposed methods do not differ significantly from the reported methods and they were presented in table-1.2. The proposed methods also applied for Biological Samples (Blood) for good recoveries are obtained which were recorded in table-1.7.

F. Accuracy

Recovery studies were carried by applying the Standard addition method to Drugs sample present in formulations for the known amount of Cefepime the recovery studies were carried .By applying the same method to Biological sample (Blood) to which known amount of Cefepime correspond to 2 mg Formulations taken by the patient. By the follow of Standard addition method 2 mg of label claim was added. After the addition of these standards the contents were transferred to 100 ml volumetric flash and dissolved in solvent. Finally the volume was made up to the mark with solvent. The solution was filtered through Whitman No. 41filter paper. The mixed sample solutions were analyzed and their absorbance value was determined. At each level of recovery five determinations were performed and present in Table – 1.3. The results obtain were compared with expected results and were statistically validated in Table – 1.4.

G. Linearity and Range

The linearity of analytical method is its ability to elicit test results that are directly proportional to the concentration of analyze in sample with in a given range. The range of analytical method is the interval between the upper and lower levels of analyze that have been demonstrated within a suitable level of precision, accuracy and linearity.

H. Specificity and Selectivity

Specificity is a procedure to detect quantitatively the analyze in the presence of components that may be expected to the present in the sample matrix. While selectivity is a procedure to detect the analyze qualitatively in presence of components that may be expected to present in the sample matrix. The excipient in formulations was spiked in a pre-weighed quantity of Drugs and then absorbance was measured and calculations were done to determine the quantity of the Drugs.

I. Repeatability

Standard solutions of Cefepime were prepared and absorbance was measured against the solvent as the blank. The observance of the same concentration solution was measure six times and standard deviation was calculated and presented in table -1.5.

J. Interferences Studies

The effect of wide range of inactive, ingredients usually present in the formulations for the assay of Cefepime under optimum conditions was investigated. None of them interfered in the proposed methods even when they are present in excess fold than anticipated in formulations.

K. Stability of the color for MBTH Method

The color was developed using 5 μ g/ml of Cefepime solution and 1ml of (0.01M) MBTH solution was added followed by 1ml of (0.2%) Ferric chloride solution and resulting solution was heated at 100 ⁰ c for 15 min. The solutions were cooled to room temperature and added 1ml (0.2M) of orthophosphoric acid solution and made up to mark with distilled water in 10 ml volumetric flask. The absorbance of Bluish green colored species was measured at 610 nm against the reagent blank.

IV. CONCLUSION

The proposed method can be used for determination of Cefepime in Formulations. The method is rapid, simple and has great sensitivity and accuracy. Proposed method makes use of simple reagents, which an ordinary analytical laboratory can afford. Method is sufficiently sensitive to permit determination even down to $10\mu g ml^{-1}$. The proposed method is suitable for routine determination of Cefepime in its formulation and Blood. The commonly used additives such as Starch, Lactose, Titanium dioxide, and Magnesium state do not interfere with the assay procedures.



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538

Volume 13 Issue VI June 2025- Available at www.ijraset.com

REFERENCES

- [1] Hawraa Jumaa Hashim, Nisreen Kais Abood* and Olfat A. Nief "Spectroscopic Estimation of Cefepime by using Batch, Cloud Point extraction and Flow Injection Analysis methods" Egyptian Journal of Chemistry, DOI: 10.21608/EJCHEM.2021.62847.3440Egypt. J. Chem. Vol. 64, No. 12 pp. 689(2021)
- [2] Dalia M. Jamil'New spectrophotometric method for determination of cefepime in pure and pharmaceutical formulation by cloud point extraction in trace" Elsevier, https://doi.org/10.1016/j.rechem.2024.101429
- [3] Sathyanarayana L, Samatha K^{*}, Susmitha K, Sanath kumar Development of UV Visible Spectrophotometric Method for Estimation of Cefipime Hydrochloride in Bulk and Dosage Form Journal of Chemical and Pharmaceutical Research (ISSN : 0975-7384) DOI: 10.37532/0975-7384.2022.14(3).019.
- [4] Imad Tarek Hanoon "A New Spectrophotometric Method for the determination of Cefepime in pharmaceutical formulations" Systematic Review Pharmacy Vol 11, Issue 6, 2020 DOI: 10.31838/srp.2020.6.52Sys Rev Pharm 2020; 11(6): 330 336
- [5] roopa kothathi papanna*, jayanna bidarur krishnegowda^a, padmarajaiah nagaraja^b spectrophotometric method for the determination of cefepime, cefazolin sodium and cefalothin sodium in pure and pharmaceutical dosage forms by using ninhydrin" int j pharm pharm sci, vol 7, issue 5, 194-1990 riginal article, received: 09 feb 2015 revised and accepted: 05 mar 2015
- [6] Shazalia Mahmoud Ali¹, Abdalla A. Elbashir² , Hassan Y. Aboul-Enein³ Spectroscopic Methods for Analysis of Cephalosporins in Pharmaceutical Formulations' World Journal of Analytical Chemistry, Vol. 3, No. 1A, 2015, pp 21-32. doi: 10.12691/wjac-3-1A-5 | Research Article
- [7] Monika Dąbrowska, Małgorzata Starek, Jan Krzek, Elżbieta Papp, Piotr Król "A degradation study of cefepime hydrochloride in solutions under various stress conditions by TLC-densitometry" Biomedical chromatography, 07 July 2014. https://doi.org/10.1002/bmc.3288
- [8] Farida El-Dars 1Nahla S. Elshater2 and Safaa M. Abd Elaziz1 "Analytical Determination of Cefepime Residues in Rabbit' Muscles, Liver and Kidney Using HPLC"Current Science International EISSN:2706-7920 ISSN: 2077-4435, DOI: 10.36632/csi/2019.8.4.10
- [9] MARWA S. ELAZAZY* AND ABDALLA SHALABY "Validated Spectrophotometric Assay of Cefepime Hydrochloride and Cefuroxime Sodium Using a Tetrazolium Salt, ISSN: 0973-4945; E-Journal of Chemistry" http://www.ejchem.net 2012, 9(4), 2261-2267
- [10] Rabindra K. Nanda*, Dipak A. Navathar, Amol A. Kulkarni, Subodh S. Patil, "Simultaneous Spectrophotometric Estimation of Cefepime and Tazobactam in Pharmaceutical Dosage Form" International Journal of ChemTech Research CODEN (USA): IJCRGG ISSN : 0974-4290 Vol.4, No.1, pp 152-156, Jan-Mar 2012
- [11] Lella Kalyani, Stability indicating RP-HPLC method development and validation of cefepime and amikacin in pure and pharmaceutical dosage forms. J. Pharm. Sci. 54 (03) • 2018 • <u>https://doi.org/10.1590/s2175-97902018000317258</u>
- [12] R,Sagar,RP-HPLC Method for Simultaneous Estimation of Cefepime Hydrochloride and Tazobactam Sodium in Bulk and Pharmaceuticals ,Journal of chemistry,https://doi.org/10.1155/2013/208057
- [13] Cherti N, Kinowski JM, Lefrant JY, Bressolle F. J Chromatogr B ,<u>High-performance liquid chromatographic determination of cefepime in human plasma and in urine and dialysis fluid using a column-switching technique.</u> Biomed Sci Appl. 2001 Apr 25;754(2):377-86. doi: 10.1016/s0378-4347(00)00630-7. PMID: 11339281
- [14] Elkhaüli H, Linger L, Monteil H, Jehl F, <u>High-performance liquid chromatographic assay for cefepime in serum.</u> J Chromatogr B Biomed Sci Appl. 1997 Mar 7;690(1-2):181-8. doi: 10.1016/s0378-4347(96)00406-9. PMID: 9106042
- [15] Calahorra B, Campanero MA, Sádaba B, Azanza JR. Biomed Chromatogr. 1999 Jun;13(4):272-5. Rapid high-performance liquid chromatographic determination of cefepime in human plasma. doi: 10.1002/(SICI)1099-0801(199906)13:4<272::AID-BMC842>3.0.CO;2-0. PMID: 10416059
- [16] Modai J. <u>Diffusion of 3-quaternary ammonium cephem antibiotics into cerebrospinal fluid of patients with bacterial meningitis.</u> J Chemother. 1996 Feb;8 Suppl 2:83-90. PMID: 8738850.
- [17] Dafale NA, Semwal UP, Rajput RK, Singh GN ,<u>Selection of appropriate analytical tools to determine the potency and bioactivity of antibiotics and antibiotic resistance</u>. J Pharm Anal. 2016 Aug;6(4):207-213. doi: 10.1016/j.jpha.2016.05.006. Epub 2016 May 24. PMID: 29403984
- [18] Emirhan Nemutlu¹, Sedef Kir, Doruk Katlan, M Sinan Beksaç Simultaneous multiresponse optimization of an HPLC method to separate seven cephalosporins in plasma and amniotic fluid: application to validation and quantification of cefepime, cefixime and cefoperazone PMID: 19782200 DOI: 10.1016/j.talanta.2009.06.034
- [19] Behin Sundara Raj I S R Punitha S Krishnan ,Stability Studies of Cefepime Hydrochloride by Stability Indicating RP-HPLC Method, DOI: https://doi.org/10.37285/ijpsn.2013.6.3.10
- [20] 1 ella Kalyani1, Chava Venkata Nageswara Rao2, Stability indicating RP-HPLC method development and validation of cefepime and amikacin in pure and pharmaceutical dosage forms, Brazilian Journal of Pharmaceutical Sciences http://dx.doi.org/10.1590/s2175-97902018000317258











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24*7 Support on Whatsapp)