iJRASET
International Journal For Research in Applied Science and Engineering Technology



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

**Volume:** 14 **Issue:** III **Month of publication:** March 2026

**DOI:** https://doi.org/10.22214/ijraset.2026.78091

www.ijraset.com

Call: 08813907089 | E-mail ID: ijraset@gmail.com

International Journal for Research in Applied Science & Engineering Technology (IJRASET)
*ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538*
*Volume 14 Issue III Mar 2026- Available at www.ijraset.com*

# Drought Vulnerability Assesment of Vadakarapathy

Ahammed Yasim VP[1], Fathima Ridha C[2], Mohammed Sinan A P[3], Rishana Sherin MP N F[4], Prof.Anagha[5]
*Civil engineering Department, APJ KTU*

***Abstract:*** ***Drought is a recurring natural hazard that significantly impacts water resources, agriculture, ecosystems, and socio-economic stability. This study focuses on the analysis of drought characteristics using meteorological indices to evaluate temporal and spatial variations. Standardized indices such as the Standardized Precipitation Index (SPI) and the Standardized Precipitation Evapotranspiration Index (SPEI) were applied to long-term rainfall and climatic data to assess drought intensity, duration, and frequency. The results reveal distinct seasonal and inter-annual fluctuations, with severe droughts occurring predominantly during monsoon deficits. Spatial mapping highlights vulnerable zones that are consistently prone to water stress. The findings provide valuable insights for drought monitoring and management, emphasizing the need for effective early warning systems and adaptive water resource planning. This analysis contributes to strengthening regional drought resilience and supporting sustainable development strategies.***

***Keywords: Drought Analysis, Standardized Precipitation Index (SPI), Water Resource Management, Meteorological Drought, Drought Intensity, Drought Frequency, Drought Vulnerability.***

## I. INTRODUCTION

Drought is often called a “silent disaster” because it doesn’t announce itself like a flood or storm, yet it can disrupt lives for months or even years. It starts quietly—with less rainfall than usual, drying soil, and shrinking water sources—but soon its effects spread across every aspect of daily life. For communities that rely on farming and local water resources, drought can mean crop failure, water shortages, reduced income, and increased hardship for families.

In India, drought has always been a challenge because so much of the population depends on rainfall for agriculture. Even with modern irrigation and farming methods, when the monsoon fails or is uneven, millions of people feel the impact. Crops fail, farmers face economic stress, and sometimes families have no choice but to migrate temporarily in search of work.

Kerala is often thought of as a state blessed with plenty of water, thanks to its rivers, backwaters, and high rainfall. Yet, the reality is more complex. Rainfall is not always evenly distributed, population and water demand are increasing, and groundwater is being used faster than it can recharge. This means that even here, certain areas can face serious water stress.

Palakkad district, located on the eastern side of Kerala, is one such area. It lies in the rain shadow of the Western Ghats, which makes it drier than the rest of the state. Even though it is called the “rice bowl of Kerala,” its agriculture is highly vulnerable to rainfall shortages. Farmers depend on monsoon rains and groundwater, and when these fail, crops and livelihoods are at risk.

Vadakarapathy, a village in Chittur Taluk of Palakkad, is a clear example of these challenges. Most families here rely on agriculture and use wells or borewells for both irrigation and daily water needs. In years of good rainfall, life runs smooth ly. But when rains are weak or delayed, water shortages, crop losses, and livelihood struggles quickly become apparent. Over time, repeated drought events make the community more vulnerable and less able to cope with the next dry season.

Understanding drought in a place like Vadakarapathy is not just an academic exercise—it has real implications for the people who live there. By studying the causes and impacts, we can identify who is most at risk, what resources or strategies can help them cope, and how to plan better for the future. This knowledge can guide farmers, local authorities, and the community to take practical steps to reduce the impacts of drought and make the village more resilient

## II. CAUSES AND CONSEQUENCES

### *A. Casuses*

*1)* Unpredictable Rainfall: One of the main causes is the village’s location in the rain shadow region of the Western Ghats. This means that Vadakarapathy receives less rainfall than the western parts of Kerala. Even when the monsoon arrives, rainfall is often uneven and irregular, with long dry spells in between. Such unpredictability makes it difficult for farmers to plan their sowing and harvesting schedules, directly affecting crop health and yields.


©IJRASET: All Rights are Reserved 

2) Dependence on Groundwater: Most households and agricultural fields in Vadakarapathy rely on wells and borewells for water. During years of low rainfall, these sources fail to recharge, leading to declining water tables. Over time, continuous extraction without adequate recharge structures increases the village's exposure to drought.
3) Agricultural Practices: The local economy is heavily agrarian, with crops like paddy, banana, and coconut dominating the landscape. These crops require consistent water supply, and reliance on rainfed cultivation makes farming highly sensitive to rainfall variability. The lack of alternative irrigation techniques, such as drip systems or rainwater harvesting, further limits the community's ability to cope with water shortages.
4) Socio-economic Constraints: Many farmers in Vadakarapathy are smallholders with limited financial resources. They often lack access to modern technology, credit, or government support to invest in water-saving measures. Additionally, limited awareness about drought-resilient crops or sustainable farming practices reduces their adaptive capacity.
5) Infrastructure Limitations: The village has few large-scale water storage or irrigation structures. Ponds, check dams, and canals are insufficient to meet the needs during dry periods, which increases both exposure and sensitivity to drought events.

### B. *Consequences*

1) Agricultural Losses: Reduced rainfall and limited water availability often result in partial or total crop failure. Farmers lose income, and the shortage of local produce can affect food security in the village. Repeated losses may push families into debt and reduce their capacity to invest in future crops.
2) Water Scarcity: Wells, ponds, and other water sources dry up during prolonged dry periods, creating shortages not only for irrigation but also for drinking and daily use. Families may need to travel long distances or depend on water tankers, which adds financial and physical strain.
3) Livelihood Impacts: Agriculture provides employment for both farmers and laborers. When crops fail, many lose their primary source of income. Some families may be forced to migrate temporarily for work, which disrupts community life and social stability.
4) Environmental Stress: Repeated droughts lead to soil degradation, reduced vegetation cover, and declining groundwater levels. These changes reduce the natural resilience of the environment, making the village more prone to future droughts and other climate- related stresses.
5) Social Challenges: Water scarcity and economic stress can lead to conflicts over shared resources. Vulnerable groups, including women, children, and landless laborers, are disproportionately affected. Health and sanitation may also decline due to lack of adequate water, further impacting quality of life.

### C. *Area Of Studty*




## III. DROUGHT VULNERABILITY CALCULATION

Drought Vulnerability=f(Exposure, Sensitivity, Adaptive Capacity)

### A. *Exposure*

The degree to which a region, population, or system is exposed to drought-related climatic conditions**,** mainly how often, how long, and how severe drought occurs in that area.

Exposure = w1(normal rainfall deviation)+w2(normal standard precipitation index)

| Exposure value | Exposure level |
|---|---|
| 0 to 0.33 | Low exposure |
| 0.34 to 0.66 | Moderate exposue |
| 0.67 to 1 | High exposure |

- Rainfall Deviation (%)=Actual Rainfall−Normal Rainfall×100
  Normal rainfall

| Rainfall deviation | Condition |
|---|---|
| ≥ 40% | Extremely wet |
| 20% to 39% | Very wet |
| 10% to 19% | Moderately wet |
| -9% to +9% | Near normal |
| -10% to -19% | Moderate drought |
| -20% to -39% | Severe drought |
| ≤ -40% | Extreme drought |

- Standard precipitation index (SPI)=(Yearly Rainfall-Mean rainfall)/Standerd deviation
  SPI expresses how much the observed precipitation deviates from the long-term mean, in units of standard deviation.

| SPI | Condition |
|---|---|
| ≥ 2.0 | Extremely wet |
| 1.5 to 1.99 | Very wet |
| 1.0 to 1.49 | Moderately wet |
| -0.99 to +0.99 | Near normal |
| -1.0 to -1.49 | Moderate drought |
| -1.5 to -1.99 | Severe drought |
| ≤ -2.0 | Extreme drought |

| | january | feb | mar | apr | may | jun | jul | aug | sep | oct | nov | dec |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 1994 | 5.6 | 11.94 | 0.94 | 129.89 | 46.3 | 318.55 | 530.85 | 140.65 | 152.5 | 280.74 | 95.06 | 0 |
| 1995 | 7.25 | 1.52 | 1.66 | 44.25 | 106.26 | 147.46 | 347.12 | 121.87 | 305.81 | 96.97 | 107.43 | 0 |
| 1996 | 0.01 | 0.11 | 5.36 | 137.35 | 50.48 | 239.58 | 409.45 | 127.89 | 162.95 | 301.59 | 49.88 | 86.45 |
| 1997 | 0 | 0 | 48.48 | 5.55 | 96.94 | 121.51 | 516.25 | 330.15 | 42.03 | 256.32 | 367.56 | 43.76 |
| 1998 | 0 | 0 | 0.84 | 35.1 | 71.25 | 350.53 | 465.63 | 166.34 | 125.41 | 107.18 | 170.73 | 107.05 |
| 1999 | 0 | 4.77 | 0.09 | 28.23 | 123.44 | 269.63 | 397.47 | 153.87 | 11.53 | 216.59 | 55.34 | 3 |
| 2000 | 2.59 | 28.68 | 8.2 | 55.31 | 15.6 | 227.72 | 177.49 | 390.19 | 81.41 | 41.49 | 76.12 | 16.78 |
| 2001 | 0.28 | 9.59 | 1.24 | 44.82 | 42.69 | 283.11 | 192.83 | 151.37 | 34.01 | 121.69 | 228.57 | 13.17 |
| 2002 | 0 | 0 | 17.62 | 4.23 | 51.34 | 219.12 | 88.26 | 236.51 | 75.74 | 125.61 | 91.3 | 2.28 |
| 2003 | 0 | 25.53 | 101.17 | 47.56 | 6.37 | 133.87 | 188.37 | 99.09 | 13.6 | 259.74 | 27.92 | 12.44 |
| 2004 | 6.66 | 2.3 | 13.75 | 55.59 | 256.12 | 435.85 | 245.33 | 268.37 | 82.15 | 146.76 | 39.82 | 0 |
| 2005 | 9.05 | 0.49 | 2.8 | 124.77 | 79.92 | 232.4 | 697.13 | 215.94 | 243.16 | 95.74 | 85.83 | 98.67 |
| 2006 | 30.19 | 0 | 150.72 | 42.41 | 215.24 | 254.26 | 350.44 | 140.52 | 133.52 | 56.31 | 185.63 | 0 |
| 2007 | 0.82 | 0.4 | 0 | 79.82 | 54.37 | 437.4 | 469.04 | 228.41 | 259.88 | 258.32 | 28.91 | 49.47 |
| 2008 | 3.44 | 25.24 | 113.03 | 20.96 | 29.31 | 150.01 | 168.16 | 101.07 | 136.37 | 333.9 | 9.61 | 1.55 |
| 2009 | 0 | 0 | 41.79 | 2.37 | 45.85 | 98.85 | 519.05 | 114.16 | 137.75 | 164.77 | 233.78 | 0.34 |
| 2010 | 0.68 | 0 | 3.36 | 53.07 | 44.16 | 163.53 | 208.54 | 174.6 | 124.89 | 155.16 | 223.9 | 27.7 |

©IJRASET: All Rights are Reserved | SJ Impact Factor 7.538 | ISRA Journal Impact Factor 7.894 |

| | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 2011 | 0 | 75.36 | 10.25 | 103.58 | 22.02 | 428.34 | 297.21 | 269.89 | 161.66 | 156.51 | 222.41 | 14.02 |
| 2012 | 0 | 0 | 0 | 85 | 17.02 | 195.94 | 211.77 | 216.69 | 100.47 | 172.77 | 51.19 | 1.1 |
| 2013 | 0 | 61.58 | 10.17 | 75.39 | 23.68 | 443.99 | 539.95 | 248.69 | 154.82 | 140.74 | 82.22 | 13.69 |
| 2014 | 0.12 | 0 | 11.38 | 7.95 | 148.23 | 176.15 | 506.96 | 332.21 | 175.04 | 216.81 | 8.79 | 7.35 |
| 2015 | 0 | 0.08 | 4.89 | 148.26 | 234.16 | 413.44 | 144.36 | 127.73 | 114.69 | 125.3 | 127.66 | 8.02 |
| 2016 | 0 | 0 | 0 | 0.61 | 108.33 | 242.37 | 201.81 | 119.39 | 38.68 | 80.68 | 21.53 | 27.75 |
| 2017 | 3.2 | 0 | 32.23 | 22.5 | 81.9 | 214.98 | 192.5 | 227.07 | 253.06 | 41.18 | 34.96 | 20.93 |
| 2018 | 0 | 18.27 | 36.8 | 70.97 | 274.35 | 440.41 | 594.26 | 614.21 | 55.3 | 109.35 | 29.95 | 3.17 |
| 2019 | 0 | 0 | 2.94 | 57.62 | 24.19 | 118.46 | 267.28 | 610.79 | 246.02 | 289.46 | 87.38 | 5.25 |
| 2020 | 0 | 0 | 31.77 | 79.72 | 70.08 | 88.12 | 236.04 | 434.99 | 265.13 | 216.58 | 43.79 | 0 |
| 2021 | 21.02 | 1.33 | 51.09 | 127.62 | 190.7 | 159.39 | 323.97 | 129.3 | 185.73 | 342.49 | 212.62 | 23.4 |
| 2022 | 4.02 | 0.42 | 25.24 | 58.14 | 132.91 | 700.4 | 512.67 | 592.47 | 198.87 | 90.83 | 173.77 | 121.63 |
| 2023 | 0.26 | 0.42 | 0.03 | 41.16 | 5 | 24.96 | 367.97 | 21.47 | 112.01 | 165.21 | 179.38 | 37.19 |
| 2024 | 72.24 | 0 | 5.03 | 13.15 | 233.79 | 252.75 | 640.22 | 204.16 | 78.81 | 151.96 | 103.23 | 44.07 |

This is the 30-year rainfall data collected From this data
Exposure = 0.5361 (Moderate Exposure)

*B. Sensitivity*

In drought vulnerability assessment**,** Sensitivity means how much a system (such as agriculture, water resources, soil, or people) is affected when drought occurs. If a system is highly sensitive, even a small decrease in rainfall can cause large damage (crop failure, water shortage, etc.). If a system is less sensitive, it can tolerate drought better.

Sensitivity = s = (s1+ s2+ s3) / 3 GROUND WATER DATA

| year | Pre monsoon | Post monsoon | Fluctuation | avg | GW normal |
|---|---|---|---|---|---|
| 2015 | 7.43 | 4.91 | 2.52 | 6.17 | 0.205835 |
| 2016 | 8.22 | 5.84 | 2.38 | 7.03 | 0.321988 |
| 2017 | 9.62 | 6.77 | 2.85 | 8.195 | 0.479335 |
| 2018 | 9.62 | 5.51 | 4.11 | 7.565 | 0.394246 |
| 2019 | 8.29 | 4.72 | 3.57 | 6.505 | 0.25108 |
| 2020 | 6.034 | 5.25 | 0.874 | 5.642 | 0.134522 |
| 2021 | 5.262 | 4.03 | 1.232 | 4.646 | 0 |
| 2022 | 6.7 | 5.41 | 1.29 | 6.055 | 0.190303 |
| 2023 | 12.07 | 12.03 | 0.04 | 12.05 | 1 |
| 2024 | 12.95 | 10.69 | 2.26 | 11.82 | 0.968936 |

Mean = 0.394625
Range of ground water depth

| Index value | Vulnerability |
|---|---|
| 0-0.25 | Low |
| 0.25-0.5 | Moderate |
| 0.5-0.75 | High |
| 0.75-1 | Very high |

Moderate vulnerability Soil depth

| soil depth class | Area (Ha) | % area | Vulnerability score | Soil sensitivity index |
|---|---|---|---|---|
| Deep to very deep | 2583 | 53.11536 | 2 | 106.2307218 |
| Very deep | 2188 | 44.9928 | 1 | 44.9928028 |
| Rock out crops | 92 | 1.891836 | 4 | 7.56734526 |

Total soil sensitivity % = 1.587908698 Range of soil depth

| range | Vulnerability |
|---|---|
| 1-1.5 | Very Low |
| 1.5-2.5 | Low |
| 2.5-3.5 | Moderate |
| 3.5-4 | High |

Low Vulnerability

| Crops | Area | % | Sensitivity score | LULC sensitivity index |
|---|---|---|---|---|
| Coconut | 1678.42 | 34.20921 | 3 | 102.6276206 |
| Paddy Land (current Fallow) | 1093 | 22.2773 | 5 | 111.3864918 |
| Residential | 403 | 8.213862 | 2 | 16.42772413 |
| Seasonal Crop | 306 | 6.236828 | 4 | 24.94731307 |
| Paddy Land (Long Fallow) | 277 | 5.645756 | 5 | 28.22878154 |
| Virippu | 205 | 4.178267 | 4 | 16.71306921 |
| Barren Rocky Land | 145 | 2.95536 | 1 | 2.9553598 |
| Other Perinial Crops | 122 | 2.486579 | 3 | 7.45973577 |
| River / Stream | 29 | 0.591072 | 1 | 0.59107196 |
| Road | 24 | 0.489163 | 1 | 0.489163001 |
| Agricultural (Current Fallow) | 67 | 1.36558 | 4 | 5.462320182 |
| Agricultural (Long Fallow) | 46 | 0.937562 | 4 | 3.750249677 |
| Other Plantaion Crop | 70 | 1.426725 | 3 | 4.280176262 |
| Other Built Ups | 60 | 1.222908 | 2 | 2.445815007 |
| Mixed Crop | 356.92 | 7.274669 | 4 | 29.09867641 |
| Ponds | 24 | 0.489163 | 1 | 0.489163001 |

LULC sensitivity index % = 3.573527314
Range of LULC

| range | Vulnerability |
|---|---|
| 1-2 | Low |
| 2-3 | Moderate |
| 3-4 | High |
| 4-5 | Very high |

High vulnerability

*C. Adaptive Capacity*

Adaptive Capacity in drought vulnerability refers to the ability of a system, community, or region to adjust, cope with, and recover from drought conditions. It indicates how well an area can reduce the negative impacts of drought through available resources, infrastructure, technology, and management practices. Factors such as irrigation facilities, water storage systems, groundwater availability, socio-economic conditions, and effective water management policies influence adaptive capacity. Areas with better water management and stronger socio-economic support generally have higher adaptive capacity.

ADAPTIVE CAPACITY = (irrigation + population + drainage)/3

*1) Irrigation Index*

Irrigation index = Irrigated Area / Cultivable Area Total Area = 4951 Ha
Cultivable Area = 2300 Ha Irrigated Area = 2070 Ha
Irrigation Index = 2070/2300 = 0.9

*2) Population Density*

Total Population = 26459 (2011 Census) Population Density = Total Population / Area
Population Density = 26459 / 49.51 = 534 people/km

*3) Drainage Density*

| Drains | Length |
|---|---|
| Order 1 | 8.24 km |
| Order 2 | 5.90 km |
| Order 3 | 2.46 km |
| Order 4 | 5.08 km |
| Order 5 | 2.34 km |

Total length of drain = 24.02 km
Drainage density = total stream length / Area Drainage density = 24.02 / 49.51 = 0.48 km/km²

*4) Normalise These Values*

Scoring of irrigation

| Irrigation ratio | Score |
|---|---|
| <0.2 | 1 |
| 0.2-0.4 | 2 |
| 0.4-0.6 | 3 |
| 0.6-0.8 | 4 |
| >0.8 | 5 |

Scoring of drainage density

| Drainage density | Score |
|---|---|
| <1 | 1 |
| 1-2 | 2 |
| 2-3 | 3 |
| 3-4 | 4 |
| >4 | 5 |

Scoring of population density

| Population density | Score |
|---|---|
| <200 | 1 |
| 200-400 | 2 |
| 400-600 | 3 |
| 600-800 | 4 |
| >800 | 5 |

From above

| Parameter | Value | Score |
|---|---|---|
| Irrigation index | 0.90 | 5 |
| Population density | 534 people / km | 3 |
| Drainage density | 0.48 km/km² | 1 |

©IJRASET: All Rights are Reserved | SJ Impact Factor 7.538 | ISRA Journal Impact Factor 7.894 |

Adaptive capacity = (5+3+1)/3 = 3 Range of adaptive capacity

| AC Value | Vulnerability |
|---|---|
| 1-2 | Low |
| 2-3 | Moderate |
| 3-4 | High |
| 4-5 | Very high |

*D. Final Drounght Vulnerability Index (DVI)*

DVI = (EXPOSURE + SESITIVITY -ADAPTIVE CAPAITY)/3 DVI = (0.5361+1.8520-3)/3

DVI = -0.6199

Vulnerability classification

| DVI Value | Vulnerability |
|---|---|
| 0-0.25 | Low |
| 0.25-0.5 | Moderate |
| 0.5-0.75 | High |
| 0.75-1 | Very high |

DVI = -0.6199 (low vulnerability)

## III. CONCLUSIONS


This study evaluated drought vulnerability in the study area by analyzing several environmental and socio-economic factors such as soil characteristics, land use/land cover, groundwater depth, drainage density, irrigation ratio, and population density. These parameters were used to calculate the sensitivity and adaptive capacity of the area, which were then combined to obtain the Drought Vulnerability Index (DVI). The results show that the study area falls under low drought vulnerability. One of the main reasons for this may be the high irrigation of agricultural crops, which helps farmers manage water shortages during dry periods. The availability of irrigati on facilities improves the adaptive capacity of the region and reduces the overall impact of drought conditions. Overall, the study shows that effective water use and irrigation practices play an important role in reducing drought risk. The findings of this study can be useful for planners and local authorities in promoting better water resource management and sustainable agricultural practices in the future.

## IV. ACKNOWLEDGMENT

We would remember with grateful appreciation, the encouragement and support rendered by the authority of Eranad Knowledge City Technical Campus, especially Mr. ADARSH.T.K, Principal, Eranad Knowledge City Technical Campus, Manjeri, to successfully complete this project and Michael Shell and other contributors for developing and maintaining the IEEE LaTeX style files which have been used in the preparation of this template. We express our deepest sense of gratitude to Mrs. Fathima Fayiza Elachola, Head of the Department of Civil Engineering and project coordinator Mrs. Safiya.N.N, Assistant Professor, Department of CE for their keen interest and constant encouragement with our work during all stages. We greatly acknowledge all other staff members of the department and all our friends and well- wishers, who directly or indirectly contributed in this work and and Michael Shell and other contributors for developing and maintaining the IEEE LaTeX style files which have been used in the preparation of this template.Our heartfelt thanks to my family members for their kind cooperation in completing this project and last but not least, we are indebted to God Almighty for being the guiding light throughout this project and helped us to complete the same within the stipulated time.

## REFERENCES

[1] Sruthi D. Sivan & S. K. Pramada et al., (2024); "Spatiotemporal analysis of historic and future drought characteristics over a monsoon dominated humid region (Kerala) in India", Springer Nature.
[2] Sruthi D. Sivan et al., (2024); "Comparative Analysis of SPI and SPEI as Meteorological Drought Indices in the Highland Region of Kerala", ResearchGate.
[3] Nivya Sekharan et al., (2022); "Drought Intensity Mapping of Kannur District, Kerala, India", ResearchGate.
[4] R. Vigneshwari et al., (2024); "Analysis of Meteorological Drought for Selected Districts of Saurashtra Region of Gujarat", International Journal of Environment and Climate Change.
[5] Lal Thompsan et al., (2021); "Precepts for Groundwater Drought Index in Kerala: A Case Study from Chaliyar River Basin", Special Publication of the Geological Society of India.



10.22214/IJRASET

45.98

IMPACT FACTOR: 7.129

IMPACT FACTOR: 7.429

# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)