iJRASET
International Journal For Research in Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 13 **Issue:** VI **Month of publication:** June 2025

DOI: https://doi.org/10.22214/ijraset.2025.72741

www.ijraset.com

Call: 08813907089 | E-mail ID: ijraset@gmail.com

International Journal for Research in Applied Science & Engineering Technology (IJRASET)
ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538
Volume 13 Issue VI June 2025- Available at www.ijraset.com

Effect of Mass Bathing on Water Quality of the Ganga River at Haridwar, Uttarakhand

Abhishek Ranjan[1], Abhinav Sahay[2], Umesh Kumar[3], Kundan Kumar[4]
[1]*FRI Deemed University, Dehradun, Uttarakhand, India*
[2,3,4]*Amity University Patna, Bihar, India*

Abstract: ***The Ganga River, revered as one of the holiest rivers in the world, is facing significant degradation due to urbanization, industrialization, and population growth. Despite its sacred status and origin from the Gangotri glacier in the Himalayas, it accumulates pollutants from various sources including domestic, industrial, and agricultural. Over-abstraction for irrigation has further reduced its dry weather flow. Mass bathing during religious rituals contributes to pollution load. This study analyses the changes in the water quality of the Ganga after Purnima Mass Bathing in 2013. The focus is on assessing how mass bathing impacts water quality specifically during this event at Haridwar, Uttarakhand.***
Keywords: Ganga River, Water quality, Mass bathing, Haridwar, Physiochemical characteristics.

I. INTRODUCTION

Water is undeniably essential for life, with the human body containing a significant percentage of it [1,2]. However, the availability of clean water is a growing concern, with water shortages and pollution posing significant challenges [3,4]. The importance of water purification and conscious use is emphasized [5]. The control and ownership of water is also becoming increasingly politicized [6]. In ancient Sanskrit literature, the importance of water is highlighted, with a call for its protection and proper management [7]. Water is the most vital and fundamental requirement for sustaining life. It is impossible to survive without water. Rivers play a crucial role in supporting human settlements, but various natural and human-induced factors can impact the quality of river water.

India possesses 14 major river systems, 55 minor rivers, and numerous small rivers. The Ganges boasts the largest river basin in the country, ranking 15th in Asia and 29th globally. It covers over a quarter (26.2%) of India's geographical area with a drainage capacity of as much as 8,61404 Km2 [8]. The Ganges enters the plains at Haridwar (29°58'N, 78°10'E), renowned for its religious significance. This year witnessed the organization of the world's largest religious congregation here, during which studies were conducted to assess the water quality of the River Ganges before and after ceremonial baths by examining selected water quality parameters.

A series of studies have consistently shown that mass bathing at Haridwar has a significant negative impact on the water quality of the Ganga River. Several researchers found that mass bathing led to increased levels of pollutants such as biochemical oxygen demand, chemical oxygen demand, total suspended solids, ammonia nitrogen, and coliforms, as well as a decrease in dissolved oxygen. These changes make the water unsuitable for drinking or bathing and can lead to waterborne diseases [9-14]. Srivastava specifically noted that the water was not fit for either purpose [15].

II. STUDY AREA

A. Haridwar

Haridwar is located at a latitude of 29˚ 58˙N and a longitude of 78˚10˙E, with an elevation of 285.56 meters above sea level (Fig.1). The study was conducted along a stretch of the river spanning from Bhooma Niketan to Pul Jatwara, which has a width ranging from 15-40 meters and a depth between 2-10 meters. The total surface area for this section was measured at 220000 m^2, with a volume of water recorded at 1760000 m^3.

Haridwar, India, is characterized by a temperate climate with dry winters and warm to hot summers, and an annual precipitation of over 31 centimeters [16, 17]. The region has experienced significant urban development, with a random pattern of urbanization. The spatiotemporal analysis of water resources in the area has revealed an increase in surface runoff from 1995 to 2010, followed by a decrease from 2010 to 2018 [18]. The region's groundwater quality is generally good, but requires continuous monitoring due to the presence of certain ions [19]. The area has also experienced enhanced pre-monsoon warming, which has been attributed to increased atmospheric loading of absorbing aerosols [20]. The study of land surface temperature and NDVI in the region has shown a gradual increase in maximum land surface temperature, likely due to global warming and climatic changes [21].

©IJRASET: All Rights are Reserved




Fig1: Haridwar district

III. METHODOLOGY

The methodology adopted for the present study is described below:

A. Materials and Methods

A comprehensive analysis was conducted on 26th March 2013 to evaluate the influence of large-scale bathing on the condition of Ganga River water, focusing specifically on two locations: cable bridge (upstream) and Gau ghat (downstream) along the River Ganga was selected as study and sampling sites. Upstream point selected at Cable Bridge where river Ganga enters Har-ki-pauri and its water is relatively cleaner as compared to the second point (Downstream) where it gets polluted by the practice of mass bathing on different ghats along its journey through Har-ki-pauri.

The two points are taken on at upstream and other at downstream in order to compare the changes that water of Ganga goes through when it passes through the bathing ghats. In order to get the precise result, firstly the speed of flow of river ganga was calculated along with the distance between the two sampling points so that we could calculate the exact time that Ganga water takes to reach the second sampling point from its entrance.

The speed of ganga water was calculated by using a meter tape, a hollow plastic ball and a stopwatch. The time that the hollow plastic ball took to cover 1 meter was calculated and then the time that would be taken by the ball to travel 1000 meters was calculated, which in our case was 9.01 minutes, the distance between the two-sampling point was 1000 meters. After these 12 samples, six samples each from both the points were collected at a regular interval of 1 hour, from 10 am to 3 pm. Since the water took 9.01 minutes to reach Gau Ghat (Down Stream) from Cable Bridge entrance (Upstream) sampling point, two persons had to be present at these two sights and need to communicate to each other through Cell phone to remind each other about the time when the samples have to collected. Some specific physicochemical characteristics such as temperature, pH, dissolved oxygen, total dissolved solids, conductivity, alkalinity, chloride, magnesium, sodium, potassium, nitrate, and calcium were examined. All the collected samples are analyzed in the water quality laboratory of “National Institute of Hydrology” of Roorkee. The physical and chemical parameters of the analytical results of Ganga water were compared with the standard guideline values recommended by WHO and Indian standard specifications for drinking water is: 10500 for drinking and public health purposes.

B. Laboratory Analysis

Water Samples were collected from Har ki Pauri ghat on bathing days. Some selected Physio-chemical parameters viz. Temperature, pH, Conductivity, TDS, DO, Turbidity, BOD, Chloride, Hardness, TSS, TS were analyzed. Out of these parameters, Temperature, pH, DO, TDS, Conductivity and Turbidity were analyzed on spot at the time of sampling using Deluxe water and soil analysis kit Model 191 E and Transparency was also measured on the spot by Sacchi disc method. While DO, BOD, Chlorides, Hardness, TSS, TS were determined in the Research laboratory using given method of APHA and Trivedi and Goel [22, 23]

©IJRASET: All Rights are Reserved

IV. REULTS AND DISCUSSION

The sample denotations U and D means Upstream and Downstream, collected from Cable bridge and Gau ghat on the occasion of mass bathing of Purnima in Haridwar at the interval of 1 hour from 10 am to 3 pm (at the interval of 1hour). (Table 1)

Table: 1 Time interval between samples collected.

SAMPLE	TIME
U-1	10:00am
U-2	11:00am
U-3	12:00pm
U-4	1:00pm
U-5	2:00pm
U-6	3:00pm
D-7	10:09am
D-8	11:09am
D-9	12:09pm
D-10	1:09pm
D-11	2:09pm
D-12	3:09pm

1) *pH:* The pH values was recorded between 6.7-8.1(Table 2, Fig. 2). pH is most important in determining the corrosive nature of water. Lower the pH value higher is the corrosive nature of water. pH was positively correlated with electrical conductance and total alkalinity [24]. The minimum pH value was noted 6.7 (sample D6) while maximum pH value was noted 8.1. (sample U2) The pH values of all samples are well within the limits prescribed by BIS and WHO [25,26]. A range of factors contribute to the change in pH of the Ganga River water, including climate change, population growth, land use changes, and pollution control [27]. These changes have led to an increase in nutrient loading, particularly nitrogen and phosphorus, which has altered the N:P:Si stoichiometry and led to a decline in water quality [28]. The presence of heavy metals, such as Zn, Pb, Mn, Fe, Cu, Si, Al, Ni, Cd, Mg, and Co, has also been identified, although these are within acceptable limits [29,30]. The interaction of water quality with alkaline phosphatase has been found to be a useful indicator of river health [31]. However, the overall water quality of the Ganga River has declined, with higher levels of turbidity, total dissolved solids, and suspended solids, particularly in the monsoon season [32-34].

Table: 2 pH values of all the samples collected.

S. No.	***SAMPLES**	**pH Concentration**	**Guidelines**
1	U1	7.2	pH values well within Indian standards and WHO guidelines.
2	U2	8.1	
3	U3	7.3	
4	U4	7.6	
5	U5	7.3	
6	U6	6.7	
7	D1	7.1	
8	D2	7.1	
9	D3	7.1	
10	D4	7.2	
11	D5	7.2	
12	D6	6.7	

©IJRASET: All Rights are Reserved | SJ Impact Factor 7.538 | ISRA Journal Impact Factor 7.894 |




Figure: 2 Bar diagram of different values of pH

2) *Electrical Conductivity (EC):* Electrical Conductivity of the 12 samples ranges from 164-193 (µs/cm) (Table 3, Fig. 3). EC shows significant correlation with ten parameters such as temperature, pH value, alkalinity, total hardness, calcium, total solids, total dissolved solids, chemical oxygen demand, and chloride and iron concentration of water. Most of the salts in water are present in their ionic forms and capable of conducting current and conductivity is a good indicator to assess river water quality. The minimum EC value noted was 164 µs/cm (sample U3) and the maximum value noted was 193 µs/cm (sample U2).
The electrical conductivity of river water is influenced by various factors, including physicochemical parameters such as color, total dissolved solids, chloride, fluoride, total phosphorus, total alkalinity, calcium, magnesium, sodium, and dissolved oxygen [35]. Diurnal fluctuations in electrical conductivity can be attributed to evapotranspiration and earth tide effects [36]. Urbanization, particularly the expansion of impervious surfaces, can lead to increased specific conductance in streams [37]. The intrusion of river water into aquifers can also affect electrical conductivity [38]. Electrical conductivity has been proposed as a single indicator of water quality deterioration in tropical rivers [39]. The effect of pressure on the electrical conductivity of sea water has also been studied [40]. Lastly, fluctuations in electrical conductivity have been used as a natural tracer for bank filtration in a losing stream [41].

Table: 3 EC of the samples collected.

S. No.	*SAMPLES	Electrical Conductivity (µs/cm)	Guidelines
1	U1	169	No limits mentioned in Indian standards and WHO guidelines.
2	U2	193	
3	U3	164	
4	U4	166	
5	U5	168	
6	U6	165	
7	D1	168	
8	D2	166	
9	D3	166	
10	D4	165	
11	D5	165	
12	D6	168	

©IJRASET: All Rights are Reserved | SJ Impact Factor 7.538 | ISRA Journal Impact Factor 7.894 |




Figure: 3 showing bar diagram of EC for all the samples collected.

3) *Total Hardness:* The total hardness of river water can be influenced by a variety of factors, including seasonal changes, the construction of diversion hydropower dams [42], and the application of a magnetic field [43]. The use of silver nanoparticles and partial least squares modeling has also been proposed as a method for predicting total hardness [44]. Other factors such as pH, temperature, and buffer solution preparation can also impact the determination of total hardness [45]. Hardness of the Ganga River water at Haridwar during mass bathing ranged between 2 -92 mg/L. The minimum value noted was 62 mg/L (sample U4) and the maximum value noted was 92 mg/L (sample D1) (Table 4, Fig. 4). The Total Hardness values of all samples are well within the limits prescribed by BIS and WHO [25,26] .

Table: 4 Total Hardness of all the samples collected.

S. No.	*SAMPLES	Total Hardness (mg/L)	Guidelines
1	U1	72	Limits within the range of Indian standards and WHO guidelines.
2	U2	86	
3	U3	64	
4	U4	62	
5	U5	70	
6	U6	70	
7	D1	92	
8	D2	70	
9	D3	82	
10	D4	86	
11	D5	72	
12	D6	86	

©IJRASET: All Rights are Reserved | SJ Impact Factor 7.538 | ISRA Journal Impact Factor 7.894 |




Figure: 4 Total Hardness of all the samples collected

4) *Alkalinity:* Alkalinity is the measure of the capacity of the water to neutralize a strong acid. The Alkalinity in the water is generally imparted by the salts of carbonates, silicates, etc. together with the hydroxyl ions in free State. The bicarbonate alkalinity of the Ganga river water at Haridwar during mass bathing ranged between 60 to 92 mg/L(Table 5,Fig. 5). Most of the natural waters contain substantial amounts of dissolved carbon dioxide, which is the principal source of alkalinity. The minimum value of alkalinity noted was 56 mg/L (sample D3) and the maximum value noted was 92 mg/L (sample U2). The values of Alkalinity of all the samples are well within the limits prescribed by BIS and WHO [25,26]. The alkalinity of river water can be influenced by a variety of factors, including human activities, watershed geology, and seasonal variations. For example, the presence of high concentrations of nitrogen and phosphorus can lead to algae bloom, which in turn can increase the pH value of the water [46]. Similarly, human-accelerated chemical weathering, acid deposition, and topography can contribute to increased river alkalinization [47]. In urban areas, the discharge from industries, including untreated wastewater, can also lead to high alkalinity levels in river water [48]. The reduction of nutrient loads in rivers can result in changes in carbon and alkalinity dynamics [49]. Other factors, such as the addition of oxidizing bactericide and the equilibria of carbonic acid, can also affect the pH and alkalinity of river water [50,51].

Table: 5 showing alkalinity of all the samples collected

S. No.	SAMPLES	*Alkalinity (mg/L)	Guidelines
1	U1	60	Values well within Indian standards, there are no guidelines mentioned in WHO.
2	U2	92	
3	U3	60	
4	U4	64	
5	U5	66	
6	U6	60	
7	D1	64	
8	D2	64	
9	D3	56	
10	D4	72	
11	D5	62	
12	D6	66	




Figure: 5 Bar diagram for Alkalinity of all the samples collected.

5) Chlorides: Chlorides of the Ganga River water at Haridwar during mass bathing ranged between 1-2 mg/L (Table 6, Fig. 6). In majority of the sample's chlorides are not determined. The minimum value of chloride noted was 1mg/L (samples U4, U6 and D1) and the maximum value noted was 2 mg/L (sample D3). The values of Chlorides in all the samples are well within the limits prescribed by BIS and WHO [25,26]. The Ganga River's chloride levels are influenced by a variety of factors, including heavy metal contamination [52], climate change and socio-economic changes [27], and urban influences [28]. These factors have led to an increase in chloride levels, as seen in the case of Japan's rivers [53]. The presence of heavy metals in the river's sediments [54] and the seasonal variation in hydro-chemical parameters [29] further contribute to the changes in chloride levels. The decline in water quality due to untreated sewage and industrial effluents [32] and the presence of heavy metals in the river's surface water also play a role in these changes [55].

Table: 6 values of Chlorides of all the samples collected

S. No.	*SAMPLES	Chlorides (mg/L)	Guidelines
1	U1	0	Limits are within The Indian standards and WHO guidelines.
2	U2	0	
3	U3	0	
4	U4	1	
5	U5	0	
6	U6	1	
7	D1	1	
8	D2	0	
9	D3	2	
10	D4	0	
11	D5	0	
12	D6	0	

©IJRASET: All Rights are Reserved | SJ Impact Factor 7.538 | ISRA Journal Impact Factor 7.894 |




Figure: 6 Bar diagram for Chloride of all the samples collected.

6) Calcium (Ca): Magnesium of the Ganga River water at Haridwar during mass bathing ranged between 16-26 mg/L (Table 7, Fig. 7). The minimum value noted was 16 mg/L (samples D1) and the maximum value noted was 26 mg/L (sample D3). The values of Calcium for all the samples are well within the limits prescribed by BIS and WHO [25,26]. A range of factors contribute to changes in calcium levels in the Ganga River, with potential impacts on water quality and ecosystem health. These include water-rock interaction, vegetation, atmospheric deposition, and urban influences.
[55- 61]. For example, Pandey highlight the role of changing atmospheric deposition chemistry and urban influences in increasing nutrient and heavy metal concentrations in the river. These changes can lead to imbalances in nutrient ratios, potentially affecting river ecology [62]. The presence of Cryptomeria japonica in headwater streams can also alter calcium dynamics [59].

Table: 7 Values for Calcium of all the samples collected

S. No.	SAMPLES	*Calcium (mg/L)	Guidelines
1	U1	23	Values within Indian standards, there are no WHO guidelines mentioned.
2	U2	24	
3	U3	22	
4	U4	23	
5	U5	20	
6	U6	20	
7	D1	16	
8	D2	22	
9	D3	26	
10	D4	22	
11	D5	17	
12	D6	21	

©IJRASET: All Rights are Reserved | SJ Impact Factor 7.538 | ISRA Journal Impact Factor 7.894 |




Figure: 7 Bar diagram for Calcium of all the samples collected.

7) Magnesium: Magnesium of the Ganga river water at Haridwar during mass bathing ranged between 1-13 mg/L (Table 8,Fig. 8). The minimum value noted for Magnesium was 1mg/L (sample U4) and the maximum value noted was 13mg/L (sample D1). The values of Magnesium for all the samples are well within the limits prescribed by BIS and WHO [25,26]. A range of studies have identified various factors contributing to changes in the magnesium levels of the Ganga River. Pandey [57,63]and Khan [64] allude to the impact of changing atmospheric deposition chemistry, with increased levels of nitrogen, phosphorus, and heavy metals being significant contributors. This is further exacerbated by urban influences, as evidenced by Pandey and Pandey, which have led to imbalances in nutrient limitation and shifts in the N:P: Si stoichiometry [28,58]. The presence of heavy metals in the river, particularly downstream of urban areas, has been highlighted by Matta (2018) and Pandey (2010), with the latter also noting a correlation between heavy metal concentration and atmospheric deposition. These changes have led to a slight to high level of pollution in the river, as indicated by Matta and Matta [29,30].

Table: 8 Values for Magnesium of all the samples collected.

S. No.	*SAMPLES	Magnesium (mg/L)	Guidelines
1	U1	3	Value are within Indian standards and WHO guide lines.
2	U2	6	
3	U3	2	
4	U4	1	
5	U5	5	
6	U6	5	
7	D1	13	
8	D2	3	
9	D3	4	
10	D4	8	
11	D5	7	
12	D6	8	




Figure: 8 Bar diagram for Magnesium of all the samples collected**.**

8) Sodium (Na): Magnesium of the Ganga river water at Haridwar during mass bathing ranged between 3.14-3.41 mg/L (Table 9,Fig. 9). The minimum value noted for sodium was 3.12 mg/L (sample D3) and the maximum value noted was 3.43 mg/L (sample D1). The values of Sodium for all samples are well within the limits prescribed by BIS and WHO [25,26].

A range of factors have been identified as causes of change in the sodium levels of the Ganga River. Pandey [58,63,] and Chatterjee [65] all point to the impact of changing atmospheric deposition chemistry, with increased levels of nitrogen, phosphorus, and sodium from this source. This is further exacerbated by the dissolution of saline-alkaline soils, which contributes to the river's sodium levels [65]. The changing stoichiometry of nitrogen, phosphorus, and silicon in the river, as well as the impact of carbon and nutrient loading, also play a role in altering the river's sodium levels [28,57]. These changes have significant implications for the river's ecology and water quality, as highlighted by Tiwari [66].




Figure: 9 Bar diagram for Sodium of all the samples collected.

Table: 9 values for Sodium of all the samples collected.

S. No.	***SAMPLES**	**Sodium (mg/L)**	**Guidelines**
1	U1	3.15	Values are well within Indian standards and WHO guidelines.
2	U2	3.22	
3	U3	3.4	
4	U4	3.41	
5	U5	3.3	
6	U6	3.21	
7	D1	3.43	
8	D2	3.14	

9	D3	3.12	
10	D4	3.21	
11	D5	3.2	
12	D6	3.29	

9) Potassium (K+): Potassium of the Ganga river water at Haridwar during mass bathing ranged between 2.12-2.29 mg/L(Table 10,Fig. 10). The minimum value noted was 2.12 mg/L (sample D2) and maximum value noted was 2.29 mg/L (sample U4). A range of factors have been identified as causes of change in potassium (K+) levels in the Ganga River, including atmospheric deposition of nitrogen and phosphorus [63], climate change and socio-economic factors [27,67], and anthropogenic activities such as urban-industrial development [58]. These changes have significant impacts on the river's water quality, with increased nutrient and carbon loads from both point and non-point sources [31] and a decline in water quality due to untreated sewage and industrial effluents [52,68]. The specific impact of these changes on K+ levels in the river, however, is not explicitly addressed in the literature.

Table: 10 Values for Potassium of all the samples collected.

S. No.	***SAMPLES**	**Potassium (mg/L)**	**Guidelines**
1	U1	2.17	No limits mentioned in Indian standards as well as in WHO guidelines.
2	U2	2.2	
3	U3	2.2	
4	U4	2.29	
5	U5	2.19	
6	U6	2.18	
7	D1	2.21	
8	D2	2.12	
9	D3	2.13	
10	D4	2.17	
11	D5	2.14	
12	D6	2.15	




Figure: 10 Bar diagram for Potassium of all the samples collected

10) *Nitrate:* Sample analysis of Ganga river during mass bathing the level of nitrate ranged between 1.76-6.6 mg/L(Table 11,Fig. 11). The minimum value noted for nitrate was 1.76 mg/L (sample U2) and the maximum value noted was 6.6 mg/L (sample U3 and U5). All the 12 samples of upstream and downstream are well within the prescribed by the BIS [25] but the sample U3 and U5 values are marginally higher than WHO [26] prescribed value. A range of factors contribute to the changes in nitrate levels in the Ganga River, including atmospheric deposition of nitrogen and phosphorus [63], future climate and socio-economic changes [67], and non-point source-driven carbon and nutrient loading [69]. These changes have significant impacts on the river's water quality, with implications for aquatic life and human health [66,70]. The use of multivariate statistical techniques can help in evaluating and managing these changes [71].

Table: 11 Values for Nitrate of all the samples collected.

S. No.	*SAMPLES	Nitrate (mg/L)	Guidelines
1	U1	3.08	Values within Indian standards But maximum value above WHO guidelines.
2	U2	1.76	
3	U3	6.6	
4	U4	2.64	
5	U5	6.6	
6	U6	5.72	
7	D1	2.64	
8	D2	3.08	
9	D3	3.08	
10	D4	0	
11	D5	0	
12	D6	4.84	




Figure: 11 Bar diagram for Nitrate of all the samples collected.

11) *BOD:* Bio-Chemical Oxygen Demand (BOD) of the Ganga river water at Haridwar during mass bathing ranged between 0.80-9.20 mg/L (Table 12, Fig. 12). The minimum value noted for BOD was 0.8 mg/L (sample D3 and D5) and the maximum value noted was 9.2 mg/L (sample U2). The values of BOD for all samples are well within the limits prescribed by BIS [25] but the value of BOD of sample U1 is well above prescribed WHO [26] values A range of factors have been identified as causes of change in the BOD of the Ganga River. These include increased atmospheric deposition of nitrogen and phosphorus [63], future climate and socio-economic changes [67], and the impact of sewage treatment plants' effluent discharge [72]. The COVID-19 lockdown also had a significant impact, leading to a decrease in dissolved oxygen and an increase in BOD [73]. The river's water quality has been further affected by non-point source-driven carbon and nutrient loading [69]. These changes have implications for the river's suitability for aquatic life [66] and the effectiveness of management strategies [27].

Table: 12 showing values for BOD of all the samples collected.

S. No.	*SAMPLES	BOD (mg/L)	Guidelines
1	U1	9.2	Values within Indian standards but maximum value above WHO guidelines.
2	U2	8.4	
3	U3	0	
4	U4	0	
5	U5	5.6	
6	U6	2.8	
7	D1	0	
8	D2	6.4	
9	D3	0.8	
10	D4	2.8	
11	D5	0.8	
12	D6	2	




Figure: 12 Bar diagram for BOD of all the samples collected

Table: 13 Maximum and minimum values of all the parameters along with Standards.

S. No	Parameter	Unit	Maximum	Minimum	Average	Indian standards	WHO standards
1	pH		8.1	6.7	7.4	6.5-9.5	6.5-9.5
2	EC	µs/cm	193	164	178.5	-	-
3	TH	mg/L	92	62	77	300	200
4	Alkalinity	mg/L	92	56	74	200	-
5	Cl^-	mg/L	1	2	1.5	250	250
6	Ca^{++}	mg/L	26	16	21	75	-
7	Mg^{++}	mg/L	13	1	7	30	150
8	Na^+	mg/L	3.43	3.12	3.27	180	200
9	K^+	mg/L	2.29	2.12	2.20	-	-
10	NO_3^-	mg/L	6.6	1.76	4.18	45	3
11	BOD	mg/L	9.2	0.8	5	30	6

©IJRASET: All Rights are Reserved | SJ Impact Factor 7.538 | ISRA Journal Impact Factor 7.894 |

V. CONCLUSION

The systematic study conducted on the Ganga River during the Purnima Mass Bathing in Haridwar on 26th March 2013 provides valuable insights into the impact of mass bathing on water quality. Two key sampling points, Cable Bridge (Upstream) and Gau Ghat (Downstream), were chosen to assess changes in water quality as the river passes through bathing ghats in Har-ki-pauri.

The analysis of physico-chemical parameters, including pH, Electrical Conductivity (EC), Total Hardness (TH), Alkalinity, Chlorides, Calcium (Ca), Magnesium, Sodium (Na), Potassium (K+), Nitrate, and Bio-Chemical Oxygen Demand (BOD), revealed several noteworthy findings. The results indicate that the water quality of the Ganga River during mass bathing remains within acceptable limits set by Indian standards and World Health Organization (WHO) [25,26] guidelines for most parameters.

pH values ranged from 6.7 to 8.1, demonstrating the water's neutral to slightly alkaline nature. EC, indicative of water salinity, fluctuated between 164 and 193 µs/cm. The TH values, an important aspect of water hardness, fell within the range of 62 to 92 mg/L. Alkalinity ranged from 60 to 92 mg/L, while chloride levels remained minimal, between 1 and 2 mg/L.

The concentrations of calcium, magnesium, sodium, and potassium were within acceptable limits, indicating the absence of excessive mineral content. Nitrate levels, though generally within Indian standards, slightly exceeded WHO guidelines in a few samples. BOD values, representing organic pollution, ranged from 0.8 to 9.2 mg/L. While most values were within Indian standards, one sample exceeded the WHO guideline.

In conclusion, despite the practice of mass bathing, the water quality of the Ganga River at Haridwar during the Purnima Mass Bathing event remains relatively stable and within acceptable limits for the majority of parameters. Continuous monitoring and proactive measures are essential to ensure the preservation of this sacred river's ecological integrity and safeguard the health of those participating in religious rituals.


REFERENCES

[1] Kumar, M. and Avinash Puri. “A review of permissible limits of drinking water.” Indian Journal of Occupational and Environmental Medicine 16 (2012): 40 - 44.https://doi.org/10.4103/0019-5278.99696
[2] Murray, MargaretM. and Dagmar Curjel Wilson. “The Importance of Water.” Health Education Journal 4 (1946): 42 - 46. https://doi.org/10.1177/001789694600400110
[3] Kilic, Zeyneb. “The importance of water and conscious use of water.” (2020).https://doi.org/10.15406/IJH.2020.04.00250
[4] Mititelu, Magdalena et al. “Water, the indispensable component for the health and functioning of the human body.” Farmacist.ro (2023): n. pag.https://doi.org/10.26416/farm.210.1.2023.7760
[5] Sanghavi, Yash and A. Balaji. “Water: An imperative human resource.” 2013 IEEE Integrated STEM Education Conference (ISEC) (2013): 1-3. https://doi.org/10.1109/ISECon.2013.6525227
[6] Black, Maggie. “The No-Nonsense Guide to Water.” (2004).
[7] Nerkar, Sanjivani Shripad. “The Importance of Water in Ancient Sanskrit Literature ‘पथयां ृ ीण रनान जलमनंसभाषतम ु ’.” (2020).
[8] Saini, P., Sharma, V., and Joshi, B. D.(2009) A Random survey report about rise in water borne diseases in Haridwar city during Kanwarmela (2008). J. Environ. Bio-Sci., 23 (2): 215–219.
[9] Sinha, Atul Kumar et al. “Impact of mass bathing on the water quality of the Ganga River at Haudeshwarnath (Pratapgarh), India--a case study.” The Science of the total environment 101 3 (1991): 275-80 .https://doi.org/10.1016/0048-9697(91)90041-C
[10] Tyagi, Vinay Kumar et al. “Impairment in water quality of Ganges River and consequential health risks on account of mass ritualistic bathing.” Desalination and Water Treatment 51 (2013): 2121-2129.https://doi.org/10.1080/19443994.2013.734677
[11] Kulshrestha, H K and S. Sharma. “Impact of mass bathing during Ardhkumbh on water quality status of river Ganga.” Journal of environmental biology 27 2 Suppl (2006): 437-40 .
[12] Bhadula, Sushil and Bishambhar Datt Joshi. “Impact of Religio-Touristic Activities on the Water Quality of Ganga River and Solid Waste Generation within Haridwar City, India.” The International Journal of Plant, Animal and Environmental Sciences 2014 (2014): n. pag.
[13] Kumar, V. et al. “Water Quality of River Ganga with Reference to Physico-Chemical and Microbiological Characteristics during Kanwar Mela 2017, at Haridwar, India: A Case Study.” (2018).
[14] Kar, Sayanti. “Determination of Water Quality Index (WQI) During Mass Bathing in Different Ghats of River Ganga in Howrah and North 24 Parganas District, West Bengal, India.” International Journal for Research in Applied Science and Engineering Technology (2017): 1097-1104.
[15] Srivastava, R.K., Sinha A.K., Pandey, D.P., Singh, K.P. and Chandra, h.(1996). Water quality of the river Ganga at Phaphamau (Allahabad) :Effect of mass bathing during Maha Kumbh, Environ.Toxicol. Water Qual., 11(1): 1-5..
[16] Jha, Ramakar et al. “Analysis of urban development of Haridwar, India, using entropy approach.” KSCE Journal of Civil Engineering 12 (2008): 281-288.https://doi.org/10.1007/S12205-008-0281-Z
[17] Pranuthi, G. et al. “Trend and Change Point Detection of Precipitation in Urbanizing Districts of Uttarakhand in India.” Indian journal of science and technology 7 (2014): 1573-1582.
[18] Pathak, Shray et al. “Spatiotemporal Analysis of Water Resources in the Haridwar Region of Uttarakhand, India.” Sustainability (2020): n. pag.https://doi.org/10.3390/su12208449
[19] Krishan, Gopal et al. “Current Groundwater Conditions in Alluvial Gangetic Plains of Haridwar District, Uttarakhand, India.” Current World Environment 11 (2016): 737-742.https://doi.org/10.12944/CWE.11.3.07

[20] Gautam, Ritesh et al. "Enhanced pre-monsoon warming over the Himalayan-Gangetic region from 1979 to 2007." Geophysical Research Letters 36 (2009): n. pag.https://doi.org/10.1029/2009GL037641
[21] Rehman, Zia ur et al. "Analysis of Land Surface Temperature and NDVI Using Geo-Spatial Technique: A Case Study of Keti Bunder, Sindh, Pakistan." Journal of Basic and Applied Sciences 11 (2015): 514-527.https://doi.org/10.6000/1927-5129.2015.11.69
[22] APHA (American Public Health Association). (1998). Standard methods for the examination of water and waste water. 20th edition. American Public Health Association Inc., New York, USA. 1998.
[23] Trivedi, R. K and P. K. Goel. (1986). Chemical and Biological Methods for Water Pollution Studies. Karad: Environmental Publications.173-184.
[24] Gupta, D. P., Sunita and Saharan, J. P.(2009) Physiochemical Analysis of Ground Water of Selected Area of Kaithal City (Haryana) India, Researcher, 1(2): 1-5. 4.
[25] Bureau of Indian Standards (1991) Indian standard specification for drinking water IS-10500. Indian Standard Institute, New Delhi 5.
[26] World Health Organization (2014) Guidelines for drinking water quality, 4th edn. World Health Organization
[27] Jin, Li et al. "Assessing the impacts of climate change and socio-economic changes on flow and phosphorus flux in the Ganga river system." Environmental science. Processes & impacts 17 6 (2015): 1098-110 .https://doi.org/10.1039/c5em00092k
[28] Pandey, Jitendra et al. "The Changing State of N : P : Si Stoichiometry in Ganga River : Possible Implications for Production and Fate of Phytoplankton Biomass." (2015).https://doi.org/10.14294/water.2015.1
[29] Matta, Gagan et al. "Assessment of Heavy Metals Toxicity and Ecological Impact on Surface Water Quality Using HPI in Ganga River." INAE Letters 3 (2018): 123-129.https://doi.org/10.1007/S41403-018-0041-4
[30] Matta, Gagan et al. "Comparative study on seasonal variation in hydro-chemical parameters of Ganga River water using comprehensive pollution index (CPI) at Rishikesh (Uttarakhand) India." Desalination and Water Treatment 118 (2018): 87-95.https://doi.org/10.5004/DWT.2018.22487
[31] Yadav, Amita and Jitendra Pandey. "Contribution of point sources and non-point sources to nutrient and carbon loads and their influence on the trophic status of the Ganga River at Varanasi, India." Environmental Monitoring and Assessment 189 (2017): 1-19.https://doi.org/10.1007/s10661-017-6188-8
[32] Singh, Leena. "PHYSICO-CHEMICAL CHARACTERISTICS OF RIVER WATER OF GANGA IN MIDDLE GANGA PLAINS." (2013).
[33] Tyagi, Shivani et al. "Multivariate Statistical Analysis of River Ganga Water at Rishikesh and Haridwar, India." Analytical Chemistry Letters 10 (2020): 195 - 213.https://doi.org/10.1080/22297928.2020.1756405
[34] Khan, Mohd Yawar Ali et al. "Assessment of surface water quality and its spatial variation. A case study of Ramganga River, Ganga Basin, India." Arabian Journal of Geosciences 9 (2015): 1-9.https://doi.org/10.1007/s12517-015-2134-7
[35] Shrestha, Arun Kumar and Nabin Bahadur Basnet. "The Correlation and Regression Analysis of Physicochemical Parameters of River Water for the Evaluation of Percentage Contribution to Electrical Conductivity." Journal of Chemistry (2018): n. pag.https://doi.org/10.1155/2018/8369613
[36] Calles, Ulla Maria. "Diurnal Variations in Electrical Conductivity of Water in a Small Stream." Hydrology Research 13 (1982): 157-164.https://doi.org/10.2166/NH.1982.0013
[37] Baker, Matthew E. et al. "Impacts of Expanding Impervious Surface on Specific Conductance in Urbanizing Streams." Water Resources Research 55 (2019): 6482 - 6498.https://doi.org/10.1029/2019WR025014
[38] Johnson, Tim et al. "Four-dimensional electrical conductivity monitoring of stage-driven river water intrusion: Accounting for water table effects using a transient mesh boundary and conditional inversion constraints." Water Resources Research 51 (2015): 6177 - 6196.https://doi.org/10.1002/2014WR016129
[39] Yap, Chee Kong. "Variations of electrical conductivity between upstream and downstream of Langat River, Malaysia: its significance as a single indicator of water quality deterioration." Pertanika journal of tropical agricultural science 36 (2013): 299-309.
[40] Horne, Ralph Albert and Galen R. Frysinger. "The effect of pressure on the electrical conductivity of sea water." Journal of Geophysical Research 68 (1963): 1967-1973.https://doi.org/10.1029/JZ068I007P01967
[41] Vogt, Tobias et al. "Fluctuations of electrical conductivity as a natural tracer for bank filtration in a losing stream." Advances in Water Resources 33 (2010): 1296-1308.https://doi.org/10.1016/J.ADVWATRES.2010.02.007
[42] Yalaletdinova, A V et al. "Probabilistic assessment of water quality by total hardness using gamma distribution." Journal of Physics: Conference Series 2388 (2022): n. pag.https://doi.org/10.1088/1742-6596/2388/1/012162
[43] Fantin-Cruz, Ibraim et al. "Changes in river water quality caused by a diversion hydropower dam bordering the Pantanal floodplain." Hydrobiologia 768 (2015): 223 - 238.https://doi.org/10.1007/s10750-015-2550-4
[44] Banejad, Hossein and E. Abdosalehi. "THE EFFECT OF MAGNETIC FIELD ON WATER HARDNESS REDUCING." (2009).
[45] Shariati-Rad, Masoud and Saeid Heidari. "Partial least squares and silver nanoparticles in spectrophotometric prediction of total hardness of water." Journal of Chemometrics 35 (2021): n. pag.https://doi.org/10.1002/cem.3345
[46] Jun-rong, Yin. "Discuss On the Determination of Total Hardness in Water." Pollution Control Technology (2011): n. pag.
[47] Baoping, Qin. "Study on the Causes of Water Bias Alkalinity in Luanhe River Channel." Research of Environmental Sciences (1999): n. pag.
[48] Kaushal, Sujay S. et al. "Increased River Alkalinization in the Eastern U." (2013).
[49] Karobi, Sadia Nazneen et al. "SEASONAL VARIATIONS IN TURAG RIVER WATER ALKALINITY." Pollution Research (2022): n. pag.https://doi.org/10.53550/pr.2022.v41i03.006
[50] Paetsch, Johannes and Helmuth Thomas. "Consequences of nutrient load reductions for carbon and alkalinity dynamics within the region of freshwater influence of River Elbe." (2021).https://doi.org/10.5194/EGUSPHERE-EGU21-1716
[51] Jinling, Sinopec. "ANALYSIS OF CAUSES FOR MEASURED ABNORMAL ALKALINITY OF CIRCULATION WATER." Large Scale Nitrogenous Fertilizer Industry (2011): n. pag.
[52] Luklema, L.. "Factors affecting pH change in alkaline waste water treatment - I." Water Research 3 (1969): 913-930.https://doi.org/10.1016/0043-1354(69)90075-X
[53] Pandey, Jitendra et al. "Heavy metal contamination of Ganga River at Varanasi in relation to atmospheric deposition." (2010).
[54] Tabayashi, Yu and Masumi Yamamuro. "Changes in the impact of anthropogenic effects on river water quality during the last 50 years in Japan." Wetlands Ecology and Management 17 (2009): 409-415.https://doi.org/10.1007/s11273-008-9117-3
[55] Leena, Singh et al. "Status of Heavy Metal Concentration in Water and Sediment of River Ganga at Selected Sites in the Middle Ganga Plain." (2012).

[56] Aktar, Md. Wasim et al. “Assessment and occurrence of various heavy metals in surface water of Ganga river around Kolkata: a study for toxicity and ecological impact.” Environmental Monitoring and Assessment 160 (2010): 207-213.https://doi.org/10.1007/s10661-008-0688-5
[57] Cenki-Tok, Bénédicte et al. “The impact of water-rock interaction and vegetation on calcium isotope fractionation in soil- and stream waters of a small, forested catchment (the Strengbach case).” Geochimica et Cosmochimica Acta 73 (2009): 2215-2228.https://doi.org/10.1016/J.GCA.2009.01.023
[58] Pandey, Jitendra et al. “Impact of changing atmospheric deposition chemistry on carbon and nutrient loading to Ganga River: integrating land–atmosphere–water components to uncover cross-domain carbon linkages.” Biogeochemistry 119 (2014): 179-198.https://doi.org/10.1007/s10533-014-9957-2
[59] Pandey, Jitendra et al. “Anthropogenically Induced Shifts in N:P:Si Stoichiometry and Implications in Ganga River.” Air, Soil and Water Research 9 (2016): n. pag.https://doi.org/10.4137/ASWR.S32780
[60] Ohta, Tamihisa et al. “The Effects of Differences in Vegetation on Calcium Dynamics in Headwater Streams.” Ecosystems 21 (2018): 1390-1403.https://doi.org/10.1007/s10021-018-0229-1
[61] Bhutiani, Rakesh et al. “QUALITY ASSESSMENT OF GANGA RIVER AT HARIDWAR WITH REFERENCE TO VARIOUS PHYSICO-CHEMICAL PARAMETERS.” (2016).
[62] Pandey, Jitendra et al. “Metal Contamination of Ganga River (India) as Influenced by Atmospheric Deposition.” Bulletin of Environmental Contamination and Toxicology 83 (2009): 204-209.https://doi.org/10.1007/s00128-009-9744-2
[63] Pandey, Jitendra et al. “Anthropogenically Induced Shifts in N:P:Si Stoichiometry and Implications in Ganga River.” Air, Soil and Water Research 9 (2016): n. pag.https://doi.org/10.4137/ASWR.S32780
[64] Pandey, Jitendra et al. “Impacts of Changing Atmospheric Deposition Chemistry on Nitrogen and Phosphorus Loading to Ganga River (India).” Bulletin of Environmental Contamination and Toxicology 91 (2013): 184-190.https://doi.org/10.1007/s00128-013-1016-5
[65] Khan, Mohd Yawar Ali et al. “Geochemistry of the Dissolved Load of the Ramganga River, Ganga Basin, India: Anthropogenic Impacts and Chemical Weathering.” Frontiers in Environmental Science (2022).https://doi.org/10.3389/fenvs.2022.823385
[66] Chatterjee, Jayati and Sunil Kumar Singh. “Impact of Dissolution of Saline-Alkaline Soils on the Hydrochemistry and Erosion Rates of the Ganga River System.” Geochemistry 23 (2022): n. pag.https://doi.org/10.1029/2021GC009914
[67] Tiwari, Ashish Kumar et al. “Time Scale Changes in the Water Quality of the Ganga River, India and Estimation of Suitability for Exotic and Hardy Fishes.” Journal of Waste Water Treatment and Analysis 7 (2016): 1-8.https://doi.org/10.4172/2157-7587.1000254
[68] Whitehead, Paul W. et al. “Dynamic modeling of the Ganga river system: impacts of future climate and socio-economic change on flows and nitrogen fluxes in India and Bangladesh.” Environmental science. Processes & impacts 17 6 (2015): 1082-97 .https://doi.org/10.1039/c4em00616j
[69] Choudhary, Leena. “PHYSICO-CHEMICAL CHARACTERISTICS OF RIVER WATEROF GANGA IN MIDDLE GANGA PLAINS.” International Journal of Innovative Research in Science, Engineering and Technology 2 (2013): 4349-4357.
[70] Singh, Rachna and Jitendra Pandey. “Non-point source-driven carbon and nutrient loading to Ganga River (India).” Chemistry and Ecology 35 (2018): 344 - 360.https://doi.org/10.1080/02757540.2018.1554061
[71] Kumar, Deepak et al. “Source characterization and human health risk assessment of nitrate in groundwater of middle Gangetic Plain, India.” Arabian Journal of Geosciences 12 (2019): 1-12.https://doi.org/10.1007/s12517-019-4519-5
[72] Bagla, Piyush et al. “Multivariate Analysis of Water Quality of Ganga River.” Journal of The Institution of Engineers (India): Series B 102 (2021): 539 - 549.https://doi.org/10.1007/S40031-021-00555-Z
[73] Bhutiani, Rakesh et al. “IMPACT ASSESSMENT OF SEWAGE TREATMENT PLANTS’ EFFLUENT DISCHARGE ON THE QUALITY OF GANGA RIVER AT HARIDWAR, UTTARAKHAND.” Journal of Mountain Research (2019): n. pag.https://doi.org/10.51220/jmr.v14i2.10
[74] Vidyarthi, Ajit Kumar et al. “Impact of COVID-19 lockdown on real-time DO–BOD variation of river Ganga.” Environmental Resilience and Transformation in Times of COVID-19 (2021): n. pag.https://doi.org/10.1016/B978-0-323-85512-9.00002-4

©IJRASET: All Rights are Reserved | SJ Impact Factor 7.538 | ISRA Journal Impact Factor 7.894 |



10.22214/IJRASET

45.98

IMPACT FACTOR: 7.129

IMPACT FACTOR: 7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)