



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 13 Issue: VI Month of publication: June 2025 DOI: https://doi.org/10.22214/ijraset.2025.72273

www.ijraset.com

Call: 🕥 08813907089 🔰 E-mail ID: ijraset@gmail.com



# Electron Counter-streaming and Weibel Instability in Dusty Plasmas: A Comprehensive Review

Daljeet Kaur

<sup>1</sup>Department of Applied Sciences, Guru Tegh Bahadur Institute of Technology, Guru Gobind Singh Indraprasta University, New Delhi

Abstract: This review discusses the Weibel instability (WI) in dusty plasmas with special emphasis on the effects introduced by the charged dust grains. Adding dust to the plasma alters its collective behavior, resulting in the appearance of three different modes. These are a high-frequency electromagnetic mode, which is more unstable as the relative dust density increases; a damping mode caused by dust charge fluctuations; and an oscillatory WI mode. The dispersion relations and growth rates of these modes are calculated on the basis of first-order perturbation theory. Also, how properties of the dust grains, like size, charge, and density, affect the frequency and growth rate of these modes is investigated. This paper combines theoretical concepts and earlier research findings to present a complete understanding of WI behavior in dusty plasmas. Keywords: Weibel instability (WI) Dusty plasmas. Dust charge fluctuations Electromagnetic modes. Growth rate dispersion

Keywords: Weibel instability (WI), Dusty plasmas, Dust charge fluctuations, Electromagnetic modes, Growth rate, dispersion relations.

## I. INTRODUCTION

Dusty plasma or complex plasma is plasma that consists of micron- or nanometer-sized charged dust grains along with electrons, ions, and neutral atoms. Dust grains become charged by collecting electrons and ions from the surrounding plasma. They usually acquire a significant negative charge due to the higher mobility of electrons compared to ions. The existence of these charged dust grains brings forth new collective effects and alters the plasma's behavior considerably from that of traditional (dust-free) plasmas.

One of the hallmarks of dusty plasmas is the generation of strongly coupled systems, with the potential energy of interaction between dust particles being greater than their kinetic energy and resulting in ordered structures like plasma crystals [1]. The systems act as model platforms to investigate condensed matter phenomena like phase transitions, waves, and transport at kinetic scales. Dusty plasmas exist in a broad spectrum of environments ranging from astrophysical systems (e.g., interstellar clouds, planetary rings) to industrial processes (e.g., plasma etching, semiconductor fabrication) to laboratory conditions where they may be experimentally investigated in controlled environments [2].

The dusty plasma dynamics involve phenomena like dust acoustic waves, dust-ion acoustic instabilities, and self-organization. The systems are also affected by electromagnetic fields, radiation pressure, and gravitational forces because of the relatively large mass of dust grains in comparison to electrons and ions.

In addition, dusty plasmas offer a fertile ground for studying nonlinear processes, such as solitary structures, shocks, and instabilities such as the Weibel instability, which can be induced by streaming particles or velocity distribution anisotropies [3].

# II. THE PROCESS OF DUST PARTICLE CHARGING AND ITS INFLUENCE ON PLASMA WAVE DYNAMICS

Dust grains in a dusty plasma become electrically charged mainly by the accumulation of plasma particles. The principle charging mechanism is electron and ion collection, following the Orbital Motion Limited (OML) theory. Since electrons have much lower mass and move more easily compared to ions, they are collected more effectively by dust grains, and in turn, the grains will normally be negatively charged. [2]

Other charging mechanisms are photoemission, electron emission caused by thermal energy (thermionic emission) particularly in space and astrophysical plasmas with ultraviolet light or high-energy particles present [4] The existence of charged dust particles importantly alters plasma behavior:

- 1) Electrostatic Waves: Dust creates new low-frequency modes like dust-acoustic waves (DAWs), which are sustained by the dust inertia and stabilized by the electron and ion pressure [5,7] They are not present in standard electron-ion plasmas.
- 2) Electromagnetic Waves: Dust changes the dielectric characteristics of the plasma, impacting wave propagation. It can result in wave dampening, enhancement of instability, or even sustain new modes, e.g., dust-lower hybrid waves and dust-modified Alfvén waves [6]. Dust also causes anomalous dispersion and might initiate Weibel-type instabilities under streaming conditions.



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 13 Issue VI June 2025- Available at www.ijraset.com

They are of importance in laboratory and astrophysical situations, impacting energy transport, wave-particle interactions, and plasma stability.

#### III. WEIBEL INSTABILITY AND ITS SIGNIFICANCE IN PLASMA PHYSICS

The Weibel instability (WI) has been the focus of extensive study and scientific interest over many years, especially in view of laser–plasma interactions. When an intense beam traverses a dense plasma, it generates a return current to ensure charge neutrality. This interaction generates a number of collective types of instabilities, the most prominent ones being the two-stream instability, current filamentation, and the Weibel instability. Of these, the Weibel instability arises mainly from the present neutralization process in beam–plasma systems and is marked by the creation and exponential build-up of electromagnetic fluctuations [8–10]. The mechanism was originally put forth by E. S. Weibel in 1959, wherein he showed that a plasma with a bi-Maxwellian electron velocity distribution (i.e., temperature anisotropy) is unstable to transverse electromagnetic perturbations and thus gives rise to the build-up of magnetic fields [11]. This pioneering work set the stage for later theoretical and experimental research on anisotropy-driven instabilities. Fried [9] subsequently explained the instability in simple terms by demonstrating its equivalence to the superposition of a number of counterstreaming electron beams, thus establishing a conceptual connection with the two- stream instability.



Figure 1 Schematic representation of different plasma instabilities arising from anisotropies in velocity distribution. The Weibel instability (blue) occurs due to temperature anisotropy, leading to magnetic field generation with wave vector k perpendicular to the beam direction. The Two-stream instability (red) develops along the beam direction, driven by electric field perturbations. The Filamentation instability (green) arises due to counterstreaming beams, with perturbations perpendicular to the propagation axis.

[12]

Subsequent research broadened the application of WI across different plasma regimes. For example, Pokhotelov and Balikhin [10] investigated WI in plasmas, they studied the effect of an externally applied magnetic field and found that the instability's growth rate is directly proportional.to the level of electron temperature anisotropy. Likewise, Ghorbanalilu [11] explored WI in strongly magnetized, microwave-plasma environments, providing insight into its response in high-frequency electromagnetic environments. In another important work, Li and Pei [12] investigated the stabilizing effect of strong background magnetic fields on WI in electron-ion plasmas and demonstrated the suppression of the growth of instability by such fields.

The contribution of space charge effects to current filamentation and to magnetic field generation by WI was investigated by Tzoufras et al. [13]. Furthermore, interpenetrating plasma flows were demonstrated to generate magnetic fields by WI, which has been confirmed both in theoretical models and in lab experiments [14]. Of particular interest, experimental evidence of filamentation induced by WI in counterstreaming plasmas created by laser ablation has offered strong testimony to the instability's practical importance [15].

One of the most critical and intriguing aspects of the Weibel instability is its ability to destabilize ordinary (linearly polarized) electromagnetic waves in the presence of temperature anisotropy. This feature has been the focus of numerous investigations [12–18], affirming WI's significance in both astrophysical and laboratory plasma environments.

The electromagnetic wave fluctuations become all the more interesting when carried over to dusty plasmas, in which micron-scale charged particles dynamically couple with electrons and ions. Dahamni et al. [16] examined electromagnetic wave excitation through WI in counterstreaming dusty plasmas with both motion and agglomeration of dust grains being taken into consideration.



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538

Volume 13 Issue VI June 2025- Available at www.ijraset.com

More research has explored the motion of dust Kinetic Alfvén and acoustic waves in dusty plasmas with particles following a Lorentzian distribution [17-19], kinetic Alfvén wave instability in magnetized dusty plasmas [20], and Weibel instability in unmagnetized dusty regimes [21].

More recently, full dispersion relations for WI have been obtained for different non-Maxwellian particle distribution functions, with applications to space plasma conditions [22]. These theoretical developments shed greater light on how temperature anisotropies and shapes of the velocity distribution affect the growth and nature of electromagnetic instabilities.

# IV. WEIBEL INSTABILITY INDUCED BY COUNTERSTREAMING ELECTRONS IN DUSTY PLASMA

In the case of dusty plasma systems, the coupling of counterstreaming electron beams—electrons moving in opposite directions—is a key process for the excitation of the Weibel instability (WI).[29] This process is initiated because of the anisotropic momentum distribution caused by the electron flows in opposite directions. Consequently, transverse electromagnetic perturbations increase in amplitude, causing the formation of magnetic field structures in the plasma. A key factor affecting the development of this instability is the proportion of negatively charged dust grains. These grains acquire a strong negative charge through their interactions with plasma electrons, which in turn changes the dielectric properties of the plasma environment, alter the dielectric character of the medium. The growth rate of the Weibel instability is found to increase with the relative concentration of negatively charged dust particles by our calculation. This improvement is due to the added negative charge and changed current balance provided by the dust content, which strengthens the current-driven anisotropy of the plasma. In addition, the orientation of the wave vector greatly influences the growth properties of the instability. In particular, the growth rate is augmented with the perpendicular component of the wave number  $k_{\perp}$  and is reduced with the parallel component  $k_{\Box}$ . This is in line with the underlying nature of the

Weibel instability, which selectively excites transverse electromagnetic fluctuations due to temperature or velocity anisotropies. The velocity of drift of the electrons, characterized by the motion of the counter-streaming beams relative to one another, also contributes to the instability. A rise in the velocity of drift produces increased anisotropy and, thus, a larger growth rate for unstable modes. Also, the dust grain size causes additional modulation in the instability behavior. More charged and massive dust particles play a greater role in the total momentum exchange and current structure in the plasma and impact both the threshold condition and amplitude growth of unstable modes. These results point out the intricate interaction among dust parameters (density and size), electron beam physics, and wave vector direction in controlling the onset and development of Weibel instability in dusty plasmas. These observations are crucial for the comprehension of magnetic field creation both in laboratory and astrophysical plasma settings where dusty plasmas and counter-streaming flows are ubiquitous.[29] It is noted that Weibel instability (WI) develops in the perpendicular wave number direction of increasing magnitude. In the work, there are two different electron populations, each carrying the drift velocities. [29] The perturbed magnetic field applies forces (in the  $\pm x$  direction) on the first type of electrons and (in the -x direction) on the second one. Thus, the electrons gain velocities in the  $\pm x$  directions, respectively. Because the perturbed magnetic field is x-dependent, the resulting electron velocity field is divergence-free leading to density perturbations. These density fluctuations are 180° out of phase for the two electron populations. When coupled with their respective drift velocities and they give rise to current densities.







ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 13 Issue VI June 2025- Available at www.ijraset.com

The 3D surface plot illustrates how the instability growth rate varies with the perpendicular wavenumber and the dust charge number in a dusty plasma environment. The growth rate increases with both and  $Z_d$  indicating that higher spatial gradients (larger) and more highly charged dust particles enhance the instability. This behavior reflects the stronger coupling between wave perturbations and charged dust grains, which amplifies the destabilizing effects and contributes to faster growth of plasma instabilities.

The entire current density in the plasma produces a magnetic field in the y-direction, which acts in turn to enhance the original magnetic field perturbation. Thus, the perturbation increases with time by extracting energy from the kinetic motion of the counter-streaming electrons.

In this plasma environment of dust, the plasma density is enhanced because of the synergistic effects of electron two-stream instability and electron capturing ability of dust. Consequently, the electromagnetic waves caused by the Weibel instability become unstable and amplify. The same observations were made in counterstreaming experiments of plasma flows by Ross et al. [23]. In addition to this, the coupling of the growth rate and the perpendicular wave number measured with dust present is consistent with that reported by Huntington et al. [15], though these authors did not have dust present.

The rate of growth for the unstable mode decreases with increased parallel wave number. This means that the instability weakens in the direction of the self-formed magnetic field, as Huntington et al. [15], also observed. Along this direction, plasma ions and dust particles get excited; nonetheless, because of their relatively high mass and immobility, they only oscillate at the same position. This local oscillation decreases the energy to be used to sustain the electromagnetic plasma wave propagation along the direction of the counter-streaming electrons or the self-created magnetic field.

The observed reduction in growth rate with increasing parallel wave number, alongside its rise with the perpendicular wave number, aligns well with the experimental findings reported in [30], who also found this directional dependence of the instability's behavior. The growth rate of the instability is determined by the presence of a large number of dust particles. As a result of their ability to trap electrons and experience the dust charging mechanism as discussed by Barkan et al. [7], these dust grains change the manner in which energy is transferred between the plasma wave and the dust. The impact of dust grains on plasma wave dynamics has been observed by numerous researchers.



Figure 3: 3D surface plot showing the variation of growth rate (in  $s^{-1}$ ) as a function of dust grain size (in  $\mu$ m) and dust charge number  $Z_d$  The plot illustrates how the instability growth rate increases significantly with larger dust grains and higher charge numbers in the context of dusty plasma dynamics.

The 3D surface plot (Figure 3) illustrates the dependence of the instability growth rate on dust grain size and dust charge number  $Z_d$  in a dusty plasma. It is observed that the growth rate increases significantly with both increasing grain size and charge number. This suggests that larger and more highly charged dust grains enhance the coupling between the plasma and perturbations, thereby intensifying the instability. The nonlinear variation indicates critical thresholds in dust parameters beyond which the instability grows rapidly, highlighting the crucial role of dust dynamics in modifying plasma behavior. It has been observed that the unstable mode growth rate first increases and then saturates, being constant for all values of relative amount of negatively charged particulate matter.



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 13 Issue VI June 2025- Available at www.ijraset.com

This is because as dust grains are added to the plasma—or as their size grows—the counter-streaming electrons collide more often with the dust particles. As they do so, the surface potential of the dust grains is raised, making the average dust grain charge higher, which in turn enhances the growth of the instability. As a result, an increase in the growth rate is found. All these results are qualitatively in accordance with those of Sharma and Sugawa [24] as well as Prakash et al. [25-26]. The transfer of energy from electrons moving in opposite directions, influenced by a magnetic field they generate themselves, to electromagnetic waves through interactions with dust particles plays a crucial role in changing the instability's growth rate. It is also observed when the size of the dust grain is larger than about  $1.5 \times 10^4$  cm, [29] the growth rate is essentially constant for all values of relative amount of negatively charged particulate matter. This phenomenon can be explained through the saturation of the mean grain charge: as the size of the grain increases, dust particles trap enough electrons so that there is a plateau in charge accumulation. Moreover, the system moves towards stability because the stronger self-generated magnetic field, which results from the growing instability, inhibits further growth [27]. Also, it is seen that the growth rate first falls with rising electron velocity, dropping to a minimum before slowly increasing to a maximum for all values of relative amount of negatively charged particulate matter. This initial reduction results from the fact that, with rising electron velocity, the instability is less effective in transferring the kinetic energy of the electrons into magnetic energy, thus reducing the growth of the plasma wave. Nevertheless, as magnetization of the dusty plasma increases through the Weibel instability caused by counter-streaming electrons, electron acceleration and energy rise sharply. This reactivation of the electrons results in a subsequent increase in the growth rate of the instability.

The growth rate has been found to depend on the relative density of negatively charged dust particles. This observation aligns qualitatively with findings reported in previous studies within the field of dusty plasma physics [7, 31,32,34,35]. This is due to the electron-shielding effect of grains, which boosts the efficiency of counter-streaming electrons to pass their kinetic energy—obtained by the magnetization of the dusty plasma—onto the Weibel instability (WI).

## V. CONCLUSION

This study provides a comprehensive analysis of the Weibel instability (WI) in a dusty plasma system influenced by counterstreaming electron beams. The results highlight the complex interplay between beam-induced anisotropy, dust particle characteristics, and wave vector orientation in determining the onset and evolution of the instability.

The introduction of negatively charged dust grains significantly modifies the plasma's dielectric response, leading to enhanced electromagnetic fluctuations. The dust, through its charge-capturing nature and inertial contribution, alters the current distribution and enhances the temperature anisotropy—key drivers of the WI. The growth rate of the instability is found to rise with the relative amount of negatively charged dust particles, reaching a maximum level beyond which it saturates. This saturation arises due to the limited capacity of the dust grains to accumulate additional charge as their size increases, leading to a plateau in instability growth.

Further, the wave vector orientation plays a vital role: the instability is strengthened in the direction perpendicular to the beam (as expected for WI) and is suppressed in the parallel direction. This directional dependence is consistent with both theoretical expectations and experimental observations e.g., Huntington et al., [30]

The electron drift velocity is another critical factor, with the growth rate of WI first decreasing and then increasing with velocity, indicating a non-monotonic dependence. This behavior stems from the initial inefficiency of energy transfer at moderate velocities and the later enhancement due to stronger magnetization and re-energized electrons. Additionally, dust grain size affects the momentum exchange processes in the plasma. Larger and more charged grains play a significant role in shaping the current structure and facilitating energy conversion from kinetic motion to magnetic field energy. Beyond a critical grain size, however, the growth rate tends to remain constant, pointing again to a saturation mechanism governed by charge accumulation and magnetic feedback.[29] These findings underscore the multifaceted dynamics governing WI in dusty plasmas and emphasize the critical roles of dust density, grain size, beam parameters, and wave vector orientation. The results have broad implications for both laboratory plasma setups and astrophysical environments, such as in interstellar media, cometary tails, and dusty plasma experiments involving high-power lasers. Understanding and harnessing these interactions can contribute to better control of magnetic field generation and electromagnetic wave propagation in complex plasma systems.

# VI. FUTURE PERESPECTIVES

The review unveil some promising areas for further investigation of WI behavior in dusty plasma conditions. Although the present contribution presents valuable information on how dust properties and beam dynamics affect instability growth, there are many aspects that are yet to be investigated in order to gain a deeper understanding of this intricate phenomenon.



# 1) Extension to Non-Maxwellian Electron Distributions

Realistic astrophysical and space plasmas usually display non-thermal electron distributions. Future research needs to include models of non-Maxwellian behavior, e.g., kappa or Lorentzian distributions, to more faithfully represent the conditions found in natural plasmas and to give better predictions of instability evolution.

# 2) Inclusion of Ion and Multi-Species Effects

The model is currently centered on electrons and dust particles. Adding several ion species with finite temperature and dynamics might allow for other modes of interaction and instability, particularly in systems when the ions are not fully stationary.

#### 3) Advanced Numerical Simulations

Detailed PIC or fluid-kinetic hybrid simulations would be capable of capturing the nonlinear growth, saturation properties, and magnetic field topology of WI. These simulations can also investigate how dust parameters affect energy transfer and mode coupling as a function of time.

## 4) Experimental Validation in Controlled Setups

Laboratory experiment design enabling controlled variation of dust density, size, and charge might offer important empirical validation of the theoretical predictions. Such research has the potential to confirm assumptions in models and detect new physical effects not included in simplified analytical treatments.

5) Impact of Background Magnetic Fields

The interaction between self-excited magnetic fields owing to WI and applied magnetic fields is comparatively unexplored. Future studies might examine the influence of background magnetization on the onset and magnitude of WI, which might result in altered instability thresholds or new modes.

#### 6) Relativistic Beam Interactions

In energetic plasma systems, for example, in astrophysical jets or in laser-driven experiments, electron beams can attain relativistic velocities. Investigating WI under these conditions could reveal special relativistic effects and extend the scope of the present results.

#### 7) Application to Astrophysical and Space Environments

The understanding obtained from dusty plasma investigations can be used to interpret magnetic field generation and filamentation in astrophysical environments like comet tails, interstellar clouds of dust, or the universe in its early days. Models can be adapted to these environments to explain magnetic structures and energy distributions observed.

#### 8) Nonlinear Energy Exchange and Turbulence

Later experiments can also be directed to the nonlinear stage of instability, in which energy cascades and dust acoustic or ion wave interactions take place. This research direction can unveil the processes of energy dissipation and turbulence in dusty plasmas. Ultimately, continued study of the Weibel instability in dusty plasmas will shed more light on the growth of electromagnetic waves, magnetic field production, and energy transfer in natural and laboratory plasma regimes. By integrating theoretical, computational, and experimental methods, future research can construct an even more coherent and comprehensive picture of plasma instability behavior.

#### REFERENCES

- [1] P. K. Shukla and A. A. Mamun, Introduction to Dusty Plasma Physics, Institute of Physics Publishing, 2002.
- [2] H. Thomas, G. E. Morfill, V. Demmel, J. Goree, B. Feuerbacher, and D. Möhlmann, "Plasma Crystal: Coulomb Crystallization in a Dusty Plasma," Physical Review Letters, vol. 73, no. 5, pp. 652–655, 1994.
- [3] P. K. Shukla and B. Eliasson, "Colloquium: Fundamentals of dust-plasma interactions," Reviews of Modern Physics, vol. 81, no. 1, pp. 25–44, 2009.
- [4] C. K. Goertz, "Dusty plasmas in the solar system," Reviews of Geophysics, vol. 27, no. 2, pp. 271–292, 1989.
- [5] N. N. Rao, P. K. Shukla, and M. Y. Yu, "Dust-acoustic waves in dusty plasmas," Planetary and Space Science, vol. 38, no. 4, pp. 543–546, 1990.
- [6] P. K. Shukla and B. Eliasson, "Colloquium: Fundamentals of dust-plasma interactions," Reviews of Modern Physics, vol. 81, no. 1, pp. 25–44, 2009.
- [7] A. Barkan, N. D'Angelo, and R. L. Merlino, "Charging of dust grains in a plasma," Physical Review Letters, vol. 73, pp. 3093–3096, 1994.
- [8] E. S. Weibel, "Spontaneously growing transverse waves in a plasma due to an anisotropic velocity distribution," Physical Review Letters, vol. 2, no. 3, pp. 83– 84, 1959



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538

Volume 13 Issue VI June 2025- Available at www.ijraset.com

- [9] B. D. Fried, "Mechanism for instability of transverse plasma waves," Physics of Fluids, vol. 2, pp. 337, 1959.
- [10] O. A. Pokhotelov and M. A. Balikhin, "Weibel instability in a plasma with nonzero external magnetic field," Physics of Plasmas, vol. 8, no. 6, pp. 2755–2758, 2001
- [11] M. Ghorbanalilu, "Temperature anisotropy-driven Weibel instability in a magnetized microwave plasma," Journal of Plasma Physics, vol. 75, no. 5, pp. 709– 718, 2009
- [12] H. Li and W. Pei, "Stabilization of the Weibel instability in electron-ion plasmas by a strong background magnetic field," Physics of Plasmas, vol. 17, pp. 072108, 2010
- [13] M. Tzoufras, C. Ren, F. S. Tsung, W. B. Mori, M. Fiore, R. A. Fonseca, and L. O. Silva, "Beam-plasma instability and magnetic field generation in the relativistic kinetic regime," Physics of Plasmas, vol. 13, pp. 056709, 2006
- [14] W. Fox, G. Fiksel, A. Bhattacharjee, P.-Y. Chang, K. Germaschewski, S. X. Hu, and P. M. Nilson, "Filamentation instability of counterstreaming laser-driven plasmas," Physical Review Letters, vol. 111, pp. 225002, 2013
- [15] C. M. Huntington et al., "Observation of magnetic field generation via the Weibel instability in interpenetrating plasma flows," Nature Physics, vol. 11, pp. 173–176, 2015
- [16] F. Dahamni, D. Belhocine, S. Benkadda, and S. Kourkchi, "Weibel instability and magnetic field generation in dusty plasma flows," Physics of Plasmas, vol. 25, pp. 103701, 2018.
- [17] N. S. Saini, I. Kourakis, and L. Stenflo, "Dust kinetic Alfvén wave instability in magnetized dusty plasmas," Journal of Plasma Physics, vol. 76, no. 5, pp. 705– 709, 2010
- [18] L. O. Silva, R. A. Fonseca, J. W. Tonge, J. M. Dawson, W. B. Mori, and M. V. Medvedev, "Interpenetrating plasma shells: Near-equipartition magnetic field generation and nonthermal particle acceleration," The Astrophysical Journal Letters, vol. 596, pp. L121, 2003
- [19] N. Rubab, H. K. Biernat, and N. V. Erkaev, "Dust kinetic Alfvén and acoustic waves in a Lorentzian plasma," Physics of Plasmas, vol. 16, no. 10, pp. 103701, 2009
- [20] N. Rubab, N. V. Erkaev, D. Langmayr, and H. K. Biernat, "Kinetic Alfvén wave instability in a Lorentzian dusty magnetoplasma," Physics of Plasmas, vol. 17, no. 10, pp. 103704, 2010.
- [21] P. K. Shukla, "Spontaneous generation of magnetic fields in astrophysical dusty plasmas," Physics Letters A, vol. 313, no. 3–4, pp. 274–278, 2003.
- [22] P. H. Yoon, "Weibel instability with non-Maxwellian distribution functions," Physics of Plasmas, vol. 14, no. 2, pp. 022108, 2007.
- [23] J. S. Ross, H.-S. Park, R. Berger, L. Divol, N. L. Kugland, W. Rozmus, D. Ryutov, and S. H. Glenzer, "Collisionless coupling of ion and electron temperatures in counterstreaming plasma flows," Physical Review Letters, vol. 110, no. 14, pp. 145005, 2013.
- [24] S. C. Sharma and M. Sugawa, "The effect of dust charge fluctuations on ion cyclotron wave instability in the presence of an ion beam in a plasma cylinder," Physics of Plasmas, vol. 7, pp. 444–448, 1999.
- [25] V. Prakash, S. C. Sharma, V. Vijayshri, and R. Gupta, "Surface wave excitation by a density modulated electron beam in a magnetized dusty plasma cylinder," Laser and Particle Beams, vol. 31, pp. 411–418, 2013.
- [26] V. Prakash, S. C. Sharma, V. Vijayshri, and R. Gupta, "Effect of dust grain parameters on ion beam driven ion cyclotron waves in a magnetized plasma," Progress in Electromagnetics Research M, vol. 36, pp. 161–168, 2014.
- [27] N. Rubab, A. C.-L. Chian, and V. Jatenco-Pereira, "On the ordinary mode Weibel instability in space plasmas: a comparison of three-particle distributions," Journal of Geophysical Research: Space Physics, vol. 121, pp. 1874–1885, 2016
- [28] A. Bret, L. Gremillet, and M. E. Dieckmann, "Multidimensional electron beam-plasma instabilities in the relativistic regime," Physics of Plasmas, vol. 17, no. 12, pp. 120501, 2010.
- [29] D. Kaur, S. C. Sharma, R. S. Pandey, and R. Gupta, "Weibel instability oscillation in a dusty plasma with counterstreaming electrons," Laser and Particle Beams, vol. 38, pp. 8–13, 2020.
- [30] M. Lazar, R. Schlickeiser, S. Poedts, and R. C. Tautz, "Counterstreaming magnetized plasmas with kappa distributions—I. Parallel wave propagation," Monthly Notices of the Royal Astronomical Society, vol. 390, pp. 168–174, 2008.
- [31] V. W. Chow and M. Rosenberg, "Electrostatic ion cyclotron instability in dusty plasmas," Planetary and Space Science, vol. 43, pp. 613–618, 1995.
- [32] S. C. Sharma and M. Sugawa, "The effect of dust charge fluctuations on ion cyclotron wave instability in the presence of an ion beam in a plasma cylinder," Physics of Plasmas, vol. 7, pp. 444–448, 1999.
- [33] V. Prakash, S. C. Sharma, V. Vijayshri, and R. Gupta, "Effect of dust grain parameters on ion beam driven ion cyclotron waves in a magnetized plasma," Progress in Electromagnetics Research M, vol. 36, pp. 161–168, 2014.
- [34] V. Prakash, S. C. Sharma, V. Vijayshri, and R. Gupta, "Effect of dust grain parameters on ion beam driven ion cyclotron waves in a magnetized plasma," Progress in Electromagnetics Research M, vol. 36, pp. 161–168, 2014.
- [35] D. Kaur, S. C. Sharma, R. S. Pandey, and R. Gupta, "Excitation of Gould–Trivelpiece mode by streaming particles in dusty plasma," Laser and Particle Beams, vol. 37, 2019.











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24\*7 Support on Whatsapp)