



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 12 Issue: V Month of publication: May 2024

DOI: https://doi.org/10.22214/ijraset.2024.61935

www.ijraset.com

Call: © 08813907089 E-mail ID: ijraset@gmail.com



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538

Volume 12 Issue V May 2024- Available at www.ijraset.com

Elevating Forestry Prediction: A Study on Machine Learning Model for Plantations Survival Rate Analysis

Akshaya Rathore

DFO Guna Forest Division

Abstract: This paper details the development and preliminary findings of a machine learning model designed to predict the survival rate of plantations. Drawing data from official sources, various vegetation indices were used as features for the predictive model. Initial results show potential, despite certain limitations, suggesting avenues for further enhancement and application.

I. INTRODUCTION

Plantations play a significant role in environmental conservation and economic sustenance. Predicting their survival rates becomes essential for sustainable development and forest management. With advancements in remote sensing and machine learning, this research aims to develop a predictive model using satellite imagery indices and the Gradient Boosted Trees algorithm to determine the survival rate of plantations.

II. STUDY AREA

The study area includes plantations done by MP Forest Department in East Chhindwara Division from 2015 to 2018.

III. METHODOLOGY

A. Data Collection

The primary source for the research data was the Madhya Pradesh Forest Department's official portal www.mpforest.gov.in, from which the Plantation Survival Report and KMLs of plantations were extracted.

Plt ID Ci	rcle	Division	Range	Beat	Compt	t Category	Plt.Year	Scheme	2015	2015	2016	2016	2017	2017	2018	2018	2019	2019	2020	2020	2021
110 10	1010	DIVIDION	Honge	<u>5000</u>	Compt	COTCHOLL	110.1001	<u>serienie</u>	Pre	Post	Pre	Post	Pre	Post	Pre	Post	Pre	Post	Pre	Post	Pre
19282 Chhi	indwara l	East Chhindwara	Amarwara	Devangao	1208 RF			Working Plan Implementation		90	86.06	82.03	80.92	76.47	60	58	56.92	59.23	58.85		
		East Chhindwara		Ghatsaliwara		Misc		Working Plan Implementation		82	74.53	71.45	68.76	67.15	65.02	64.28	57.17	64.27	61.03		
		East Chhindwara		Putra		Bamboo		Working Plan Implementation		85	80.46	78.39	76.21	75.44	78			65	65		
		East Chhindwara		Bagla	1216 RF	Teak		Others				92.47	91.04	87.62	79	93.71	90	72.86	69.86	65.43	60
		East Chhindwara		Surlakhapa	1158 PF	Teak		Others				91.51	87.46	83.87	77.57	82	70.29	62.93	62.18	48.39	45.36
		East Chhindwara		Sarsdol	1230 RF	Teak		Others				94.9	88.17	81.99	85		82	79.54	79.4	77.1	59.42
		East Chhindwara		Karapatha		Teak		Others				88.57	82.22	78.6	75.14	84	74	70	69.14	67.71	63.86
		East Chhindwara		Amarwara		Teak		Others				91.26	81.55	74.87	80			68	65.4	60.4	55.7
		East Chhindwara		Dungariya	1189 RF	Teak		Others				82.64	75.6	66.42	80			79	78.09	71.71	70.13
		East Chhindwara		Medki	1131 PF	Bamboo		Others				92.64	89.98	86.78	85		74	70.08	64	58	54
		East Chhindwara				Misc		Environment Forestry				95.81	91.95	74.77	85		75	75	72.33	66.84	60.7
		East Chhindwara		Medki		Bamboo		Others				93.18	86.53	80.51	75			85.12	80	63	60
		East Chhindwara		Gourpani	1126 PF	Bamboo		Others				91.21	84.85	78.45	87			69.12	62.98	53	50
		East Chhindwara		Bhajipani	1175 RF	Bamboo		Others						94.5	80			90	85	81	71
		East Chhindwara		Dulara		Misc		Others						95.6	80		88	89.52	85	83	75
		East Chhindwara		Dhasanwara		Misc		Compensatory Afforestation						93.5	78		91.01	95.01	93.26	90.25	89.73
		East Chhindwara		Devangao		Misc		Compensatory Afforestation						95	76.16	95	90.24	94.33	94.24	94.46	94.15
		East Chhindwara		Sarsdol		Misc		Others						95.02	75.02	75.02	72	72.25	72	70.46	59.08
		East Chhindwara		Kubri		Misc		Others						89	90	95	80	78	77	76	70
		East Chhindwara		Gadadaryaw		Misc		Others						90.13	70.13	90.13	88	84	73.87	68	66.67
		East Chhindwara		Putra		Misc		Others						92	80	90	87.75	88	88	88	88
		East Chhindwara		Karapatha		Misc		Compensatory Afforestation						98.8	78	90	84	89	87.61	83.94	82.7
		East Chhindwara		Sariyapani		Misc		Working Plan Implementation								95		93.33	90	80	70
		East Chhindwara		Sariyapani		Misc		FDA (NAP)								95	90	94.02	78.05	65	55
		East Chhindwara		Devangao		Misc		FDA (NAP)								95	91.7	93.12	90.7	87	73.95
		East Chhindwara		Chimouaa		Misc		Compensatory Afforestation										100	90.18	86.4	86.24
		East Chhindwara		Karapatha		Misc		Compensatory Afforestation										96	88	89.98	77.76
		East Chhindwara		Chimouaa		Misc		Compensatory Afforestation										100	90.02	86.4	86.08
		East Chhindwara		Chimouaa		Misc		Compensatory Afforestation										100	95.2	83.2	79.2
		East Chhindwara		Tendani		Misc		Compensatory Afforestation										100	80	77.6	83.2
		East Chhindwara		Bagla		Misc		Compensatory Afforestation										100	80	80	80
		East Chhindwara		Thavari		Misc		Compensatory Afforestation										98	95.2	92.4	81.6
		East Chhindwara		Devangao		Misc		Compensatory Afforestation										99.53	99	98.64	97.84
		East Chhindwara		Sejwara		Misc		Working Plan Implementation										98.75	92.5	97	92.65
		East Chhindwara		Medki		Misc		Working Plan Implementation										97.54	94	86	83
		East Chhindwara		Baratmari		Misc		Working Plan Implementation										96.48	80	85	80
105772 Chhi	indwara l	East Chhindwara	Amarwara	Tinsai	1147 PF	Misc	2019	Working Plan Implementation										99	94	89.38	86
				•		•															

Fig.1 Survival Report of Plantations



International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 12 Issue V May 2024- Available at www.ijraset.com

Planted Divison: East Chhindwara Scheme: Working Plan Plantation Teak Implementation Category: Teak 30000 12 2 X 2 Range: Chaura Bamboo Sanction Cost: 3560451 2000 4 X 4 Maintenance Agency: Area Type : Forest 12 Kala Siras/ 500 3 X 3 107100 Actual Soil Type: Forest Kala Shirish Working Circle: Plantation Mgt Expenditure: Soils 500 12 3 X 3 12 Karanj/Kanji 500 3 X 3 Khamher/ 500 12 3 X 3 Seevan Arjun 1000 12 3 X 3 **Evaluation Details** Status Evaluated By Period Survival Type

Fig.2 KML file of Plantations download

Date

6-May-2017

Mr. Vinay Kumar Meshram 15-Oct-2015

vinay kumar meshram ro 2-Jun-2016

Vinay Kumar Meshram RO 2-Oct-2016

(%) 91.80

81.31

84.19

80.74

B. Data Processing

Oct-2015

May-

2016

2016

May-

Done

Done

Done

RO Chourai

chourai

Chourai

Lalii Uikev

Post-

Monsoon

Monsoon

Monsoon

Pre-

The KMLs are then checked for their geometrical validity and then KMLs of plantations of same year are merged to have shapefile containing geometries of all plantations of same year.





Fig.3 KML files Processing in QGIS



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 12 Issue V May 2024- Available at www.ijraset.com



Satellite imagery indices, including NDVI (Normalized Difference Vegetation Index), MCARI (Modified Chlorophyll Absorption Ratio Index), SAVI (Soil-Adjusted Vegetation Index), MSI (Moisture Stress Index), and NDWI (Normalized Difference Water Index), were considered. These indices provide crucial insights into vegetation health, soil moisture, chlorophyll content, and water stress, making them imperative for analyzing plantation survival.

Using Google Earth Engine, the slope of these indices was determined over a five-year span to track and understand growth trends. The code takes input as shapefile containing geometries of all plantations of particular year and output as CSV file containing slope trends of all indices and plantation ID.

```
// Improved cloud and shadow masking using the SCL band.
var maskCloudsAndShadows = function(image) {
 var SCL = image.select('SCL');
 var mask = SCL.neq(3).and(SCL.neq(8)).and(SCL.neq(9)).and(SCL.neq(10)).and(SCL.neq(2));
 return image.updateMask(mask);
};
// Function to convert system:time_start metadata to a band.
var addTimeBand = function(image) {
// Convert milliseconds from Unix epoch to years since 2000 for improved numerical stability
 var yearsSince2000 = image.metadata('system:time_start').divide(1000 * 60 * 60 * 24 * 365.25).subtract(2000);
 return image.addBands(yearsSince2000.rename('time'));
};
// Load Sentinel-2 Surface Reflectance data.
var collection = ee.ImageCollection('COPERNICUS/S2 SR')
  .filterDate('2016-01-01', '2022-01-01')
  .filterBounds(regions) // Adjust as per your requirements
  .map(maskCloudsAndShadows);
// Compute NDVI, SAVI, NDWI, MCARI, MSI for each image in the collection.
var computeIndices = function(image) {
 var ndvi = image.normalizedDifference(['B8', 'B4']).rename('NDVI');
 var savi = image.expression(
  '((B8 - B4) / (B8 + B4 + 0.5)) * 1.5',
  { 'B8': image.select('B8'), 'B4': image.select('B4') }
 ).rename('SAVI');
 var ndwi = image.normalizedDifference(['B3', 'B8']).rename('NDWI');
 var mcari = image.expression(
  '0.2 * (2.5 * (NIR - RED) - 1.3 * (NIR - BLUE))',
   'NIR': image.select('B8'),
   'RED': image.select('B4'),
```



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 12 Issue V May 2024- Available at www.ijraset.com

```
'BLUE': image.select('B2')
 ).rename('MCARI');
 var msi = image.select('B11').divide(image.select('B8')).rename('MSI');
 return image.addBands([ndvi, savi, ndwi, mcari, msi]);
};
var withIndices = collection.map(computeIndices).map(addTimeBand);
// Compute the linear trend over time for each index.
var trendNDVI = withIndices.select(['time', 'NDVI']).reduce(ee.Reducer.linearFit());
var trendSAVI = withIndices.select(['time', 'SAVI']).reduce(ee.Reducer.linearFit());
var trendNDWI = withIndices.select(['time', 'NDWI']).reduce(ee.Reducer.linearFit());
var trendMCARI = withIndices.select(['time', 'MCARI']).reduce(ee.Reducer.linearFit());
var trendMSI = withIndices.select(['time', 'MSI']).reduce(ee.Reducer.linearFit());
// Compute the slope for each region for each index
var computeSlopesForRegion = function(feature) {
 var slopes = {
  'slope_NDVI': trendNDVI.reduceRegion({ reducer: ee.Reducer.mean(), geometry: feature.geometry(), scale: 10, maxPixels: 1e9
}).get('scale'),
  'slope_SAVI': trendSAVI.reduceRegion({ reducer: ee.Reducer.mean(), geometry: feature.geometry(), scale: 10, maxPixels: 1e9
}).get('scale'),
  'slope NDWI': trendNDWI.reduceRegion({ reducer: ee.Reducer.mean(), geometry: feature.geometry(), scale: 10, maxPixels:
1e9 }).get('scale'),
  'slope_MCARI': trendMCARI.reduceRegion({ reducer: ee.Reducer.mean(), geometry: feature.geometry(), scale: 10, maxPixels:
1e9 }).get('scale'),
  'slope_MSI': trendMSI.reduceRegion({ reducer: ee.Reducer.mean(), geometry: feature.geometry(), scale: 10, maxPixels: 1e9
}).get('scale')
 };
 return feature.set(slopes);
};
var results = regions.map(computeSlopesForRegion);
// Export the results to a CSV
Export.table.toDrive({
 collection: results.select(['layer', 'slope_NDVI', 'slope_SAVI', 'slope_NDWI', 'slope_MCARI', 'slope_MSI']),
 description: 'index trend slopes',
 folder: 'YOUR_GOOGLE_DRIVE_FOLDER_NAME',
 fileNamePrefix: 'index_slopes',
 fileFormat: 'CSV'
});
```

Fig.4 Code Snippet Used



International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 12 Issue V May 2024- Available at www.ijraset.com


A		С	D				н
system:index	Name	slope_MCARI	slope_MSI	slope_NDVI	slope_NDWI	slope_SAVI	.geo
2 0	19282	154.5343461	-0.2164081668	0.1377919358	-0.07101912856	0.2066520858	{"type":"Polygon","co
3 1	19393	156.9190414	-0.2331009133	0.1429581158	-0.08724809969	0.2144030191	{"type":"Polygon","cc
1 2	19395	121.9599274	-0.08549358721	0.09398345108	-0.04240114477	0.1409526939	{"type":"Polygon","cc
5 4	19268	118.025751	-0.09823957037	0.0960210239	-0.05663429632	0.1440093	{"type":"MultiPolygor
5	19269	155.0995777	-0.1784348111	0.1150790121	-0.07415531469	0.1725999481	{"type":"Polygon","cc
6	19272	118.4652155	-0.2057260546	0.1049743431	-0.07113253121	0.1574418222	{"type":"MultiPolygor
7	19283	127.4521552	-0.1920846053	0.09566752602	-0.0601483986	0.1434838067	("type":"MultiPolygor
8	19318	155.3514357	0.002409663849	0.1435519647	-0.08526188324	0.2152928245	{"type":"Polygon","co
0 9	19390	177.7067534	-0.1923126285	0.1486470523	-0.08115190413	0.2229379396	{"type":"Polygon","cc
1 00000000000000000	19391	113.1864937	-0.1587692604	0.09479481155	-0.04193772272	0.1421676767	{"type":"Polygon","cc
2 00000000000000000	19392	61.75641322	-0.05044091273	0.04086791998	-0.01983820533	0.06129367215	{"type":"Polygon","co
3 < 00000000000000000	19398	87.96906089	-0.1347571117	0.07816838872	-0.040547038	0.1172323646	{"type":"Polygon","co
4 00000000000000000	19400	124.7826303	-0.1357711431	0.1119769377	-0.05095998663	0.1679373111	{"type":"Polygon","cc
5 0000000000000000	19401	151.0403436	-0.1629308578	0.1140908065	-0.05349684247	0.1711134672	{"type":"Polygon","co
6 0000000000000000	19444	131.9218121	-0.130157687	0.1021965295	-0.05026357723	0.1532714987	{"type":"Polygon","co
7 10	20045	122.7043572	-0.1588435103	0.1102083205	-0.06802105224	0.1652883545	{"type":"Polygon","cc
8 3	19399	139.2572438	-0.1347879132	0.1106562813	-0.05257724714	0.1659584235	{"type":"Polygon","cc

Fig.5 Output CSV file

D. Model Creation

Our dataset, housed within Google Sheets, encompassed 112 unique plantation records, each denoted by a Plantation ID. Each record detailed the slopes of satellite-derived indices, serving as predictive features for plantation survival rates. Using the "Simple ML for Sheets" extension, we streamlined machine learning directly within the spreadsheet, bypassing intricate coding processes. For the modelling:

- 1) Data Labelling: Plantation survival status after 5 years, extracted from official reports, was our target variable.
- Feature Selection: Continuous slope values from indices like NDVI and SAVI became our independent variables, suitable for regression models.
- 3) Model Training & Evaluation: We employed the Gradient Boosted Trees algorithm for its robustness in handling vast datasets. The extension facilitated automatic data partitioning for training and validation, subsequently evaluating the model's accuracy on unseen data.





International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538

Volume 12 Issue V May 2024- Available at www.ijraset.com




Fig.6 Model Evaluation Report

IV. RESULTS AND INTERPRETATION

Of the 112 plantations analysed, the model accurately predicted the survival of 91. The overall accuracy stood at 81.25%. However, the research encountered a limitation in the form of the database's restricted scope, sourced from just one forest division spanning four years only 2015 to 2018. Consequently, predictions for plantations with reduced survival rates showed significant errors.

V. DISCUSSION

The model, in its present iteration, holds potential, even if the accuracy isn't at an optimal level. Its primary value lies in assisting field officers in identifying plantations at risk. By flagging potential failures, proactive measures can be initiated to mitigate issues. For future iterations, it's imperative to diversify and expand the dataset. Incorporating additional indices and geometric features could further enhance the model's predictive capabilities.

VI. **FUTURE WORK**

- Augmenting the dataset is a priority, ensuring diverse data sources to refine the model further.
- New indices and geometric parameters, especially features like land surface temperature, will be considered in the updated
- Plans to automate the entire model are underway using platforms such as Google Colab, making the process more user-friendly.

REFERENCES

- Liu, Pei. Machine-Learning-Based-Survival-Analysis. github.com/liupei101
- Elith, J., & Leathwick, J.R. (2008). A working guide to boosted regression trees. besjournals.onlinelibrary.wiley.com
- Arain, A. (Year not specified). Machine Learning Approach to Quantify Leaf Depletion. github.com/Arain23









45.98



IMPACT FACTOR: 7.129



IMPACT FACTOR: 7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call: 08813907089 🕓 (24*7 Support on Whatsapp)