iJRASET
International Journal For Research in Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 10 **Issue:** I **Month of publication:** January 2022

DOI: https://doi.org/10.22214/ijraset.2022.39772

www.ijraset.com

Call: 08813907089 | E-mail ID: ijraset@gmail.com

International Journal for Research in Applied Science & Engineering Technology (IJRASET)
ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538
Volume 10 Issue I Jan 2022- Available at www.ijraset.com

Eulers Totient Function (Number Theoretic Functions), Right Angle Triangle and Their Applications

Chirag Gupta
Department of Mathematics, C.H.M College

Abstract: In this paper, we introduce some new theorem and results(section II,III and VI) on Euler's Totient Function , Right angle triangle and their applications

Keywords: Euler's totient function, right angle triangle, Golden ratio, Hardy – Ramanujan number, student's pencil compass and etc.


I. INTRODUCTION(PRELIMINARY)


1) *Definition 1.1 :* for $m \geq 1$, $\varphi(m)$ denote the number of positive integers not exceeding m that are relatively prime to m. Euler's totient function is also known as Euler's phi function.

If $m = p_1.p_2$ where p_1, p_2 are the primes $(p_2 > p_1)$, Let divisors of $m\ are\ 1, p_1, p_2\ and\ p_1.p_2$.

Then , $\varphi(m)$= (fourth divisors of m – second divisors of m) - (Third divisor of m – first divisor of m).

$$\text{Or, } \varphi(m) = (p_1.p_2 - p_1) - (p_2 - 1)$$
$$= p_1(p_2 - 1) - (p_2 - 1)$$
$$= (p_1 - 1)(p_2 - 1)$$
$$\varphi(m) = (p_1 - 1)(p_2 - 1)$$
$$If\ E\ [HCF(1, p_1), LCM(1, p_1)],$$
$$F\ [HCF(p_1, p_2), LCM\ (p_1, p_2)], and$$
$$G\ [HCF(p_2, p_1.p_2), LCM(p_2, p_1.p_2)]$$
$$Are\ the\ vertices\ of\ triangle\ EFG$$





©IJRASET: All Rights are Reserved

Then triangle EFG is inverted right angle triangle

$$And\ \varphi(m) = |A - B|$$

Where A,B are the Altitude and Base of triangle EFG

and $(i)\ If\ A > B$

then, $A = p_1.p_2 - p_1\ and\ B = p_2 - 1$

$(ii)\ If\ B > A$

then , $B = p_1.p_2 - p_1\ and\ A = p_2 - 1$

$$\varphi(m) = |A - B|$$

Squaring the both side,

$$(\varphi(m))^2 = A^2 + B^2 - 2.A.B$$

Since, $A^2 + B^2 = H^2$ where H is the hypotenuse

Then , $(\varphi(m))^2 = H^2 - 2.A.B$

divide both side of above equation by 4,

$$\left(\frac{\varphi(m)}{2}\right)^2 = \left(\frac{H}{2}\right)^2 - \frac{1}{2}.A.B$$

Or , $T = \left(\frac{H}{2}\right)^2 - \left(\frac{\varphi(m)}{2}\right)^2$

Where T is the area of triangle EFG = $\frac{1}{2}.A.B$.

$H^n - (\varphi(m))^n$ is perfectly divisible by 2^n , $n \geq 1$.

If $m = p_1.p_2 \ldots . p_z, Z \geq 2$ then total pair of primes = $\frac{z(z-1)}{2}$ Or $(Z-1)^{th}$ triangular number

Therefore, total number of inverted right angle triangle whose area is $\left(\frac{H}{2}\right)^2 - \left(\frac{\varphi(S)}{2}\right)^2$ is (z-1)th Triangular number.

Theorem and Results

II. INEQUALITY RELATION

1) Theorem 2.1: If $= p_1.p_2 \ldots . p_z$,
Where $p_1, p_2, \ldots . p_z$ are the primes

$(p_1 < p_2 < \ldots . < p_z)$, $Z \geq 3$.

Let divisors of m are $d_1, d_2, \ldots . d_{\tau(m)}$, Consider m have $^{\tau(m)}c_4$ [$^{\tau(m)}c_4 = \sum_{i=i}^{n} i[\{\tau(m)\text{-}2\text{-}i\}^{th}$ triangular number] , where n= $\tau(m) - 3$] groups , each group consists of four elements (at least one element of each group should be distinct) . let groups be ($a^{'}, b^{'}, c^{'}, d^{'}$) , ($a^{''}, b^{''}, c^{''}, d^{''}$) , ….. ($a^{''\ldots e\ times}, b^{''\ldots e\ times}, c^{''\ldots e\ times}, d^{''\ldots e\ times}$) where e = $^{\tau(m)}c_4$

And every four elements of each group belongs to the divisors of m that is $d_1, d_2, \ldots\ldots d_{\tau(m)}$. Let E_1[HCF($a^{'}, b^{'}$), LCM ($a^{'}, b^{'}$)],

F_1[HCF($b^{'}, c^{'}$), LCM($b^{'}, c^{'}$)], G_1[HCF($c^{'}, d^{'}$), LCM ($c^{'}, d^{'}$)] , E_2[HCF($a^{''}, b^{''}$), LCM ($a^{''}, b^{''}$)],


F_2[HCF($b^{''}$, $c^{''}$), LCM($b^{''}$, $c^{''}$)], G_2[HCF($c^{''}$, $d^{''}$), LCM ($c^{''}$, $d^{''}$)] , …….. E_e[HCF($a^{''\ldots e\ times}$, $b^{''\ldots e\ times}$), LCM ($a^{''\ldots e\ times}$, $b^{''\ldots e\ times}$)], F_e[HCF($b^{''\ldots e\ times}$,$c^{''\ldots e\ times}$), LCM($b^{''\ldots e\ times}$,$c^{''\ldots e\ times}$)], G_e[HCF($c^{''\ldots e\ times}$,$d^{''\ldots e\ times}$), LCM ($c^{''..e\ times}$, $d^{''..e\ times}$)] are the vertices (or points) of $E_1F_1G_1$,$E_2F_2G_2$, ….$E_eF_eG_e$ and


W_1 = Total number of groups for which E_Ψ,F_Ψ,G_Ψ form Collinear line.

Where Ψ belongs to [1,e]

W_2 = Total number of groups for which E_Ψ,F_Ψ,G_Ψ form triangles (other than right angle triangle).


W_3 = Total number of groups for which E_Ψ,F_Ψ,G_Ψ form non-inverted right angle triangles





W_4 = Total number of groups for which E_Ψ,F_Ψ,G_Ψ form inverted right angle triangles whose area is not equal to $\left(\frac{H}{2}\right)^2 - \left(\frac{\varphi(S)}{2}\right)^2$ and

W_5 = Total number of groups for which E_Ψ,F_Ψ,G_Ψ form inverted right angle triangles whose area is equal to $\left(\frac{H}{2}\right)^2 - \left(\frac{\varphi(S)}{2}\right)^2$ where $S \neq m$.



If S=product of second and third elements of each group ($b'.c'$, $b''.c''$, ….. $b^{''\ldots e\ times}.c^{''\ldots e\ times}$) , then there exist at least Z-1 + $^{Z-1}C_2$ inverted right angle triangles whose area is $\left(\frac{H}{2}\right)^2-\left(\frac{\varphi(S)}{2}\right)^2$.

That is $w_5 \geq$ Z-1 + $^{Z-1}C_2$ or, $\sum_{i=1}^{n} i$ [{τ(m)-2-i}th triangular number] – $\sum_{j=1}^{4} wj \geq$ Z-1+$^{Z-1}C_2$, where n=τ(m) –3 .

Where total number of groups = $W_1+W_2+W_3+W_4+W_5$.

Proof : If m= $P_1.P_2\ldots.P_z$, Z ≥ 3.

Total pair of primes = Z-1+$^{Z-1}C_2$ and number of groups of the form (1,P',P'',P'.P'') is Z-1+$^{Z-1}C_2$

Where P' and P'' are one of $P_1,P_2,\ldots.P_Z$ (P'<P''). If groups are of the form (1,P',P'',P'.P'') then E,F,G form inverted right angle triangles, and every Z-1+$^{Z-1}C_2$ triangles follow the relation T= $\left(\frac{H}{2}\right)^2-\left(\frac{\varphi(S)}{2}\right)^2$. Where S= $p'.p''$. By considering the groups of four elements (elements belongs to the divisors of m) , there exist some other groups also which are not of the form (1,P',P'',P'.P'') but follow the relation T = $\left(\frac{H}{2}\right)^2-\left(\frac{\varphi(S)}{2}\right)^2$. If m=$P_1.P_2.P_3$, then total number of groups = $^{\tau(m)}C_4=^8C_4=70$.

Out of these 70 groups there exist definitely Z−1+$^{Z-1}C_2$, that is 3 groups for which the vertices form inverted right angle triangle whose area is $\left(\frac{H}{2}\right)^2-\left(\frac{\varphi(S)}{2}\right)^2$.

Above 3 groups are of the form (1,P',P'',P'.P'') where P' and P'' are one of P_1,P_2,P_3 (P'<P''). But for some m there exist some other groups which are not of the form (1,P',P'',P'.P'') but follow the relation T = $\left(\frac{H}{2}\right)^2-\left(\frac{\varphi(S)}{2}\right)^2$.

If m =$2.P_2.P_3$, then the vertices (E,F,G) for the group ($2,2.p_2,2.p_3,2.p_2.p_3$) form inverted right angle triangle whose area is $\left(\frac{H}{2}\right)^2-\left(\frac{\varphi(S)}{2}\right)^2$.

So, if m=$2.P_2.P_3$, then there exist 4 inverted right angle triangle whose area is $\left(\frac{H}{2}\right)^2-\left(\frac{\varphi(S)}{2}\right)^2$.

If m=$3.P_2.P_3$, then the vertices (E,F,G) for the group ($3,3.P_2,3.P_3,3.P_2.P_3$) form inverted right angle triangle but area is not equal to $\left(\frac{H}{2}\right)^2-\left(\frac{\varphi(S)}{2}\right)^2$.

So, If m=$3.P_2.P_3$ then there exist 3 inverted right angle triangle whose area is $\left(\frac{H}{2}\right)^2-\left(\frac{\varphi(S)}{2}\right)^2$.

Hence, there exist at least Z-1+$^{Z-1}C_2$ inverted right angle triangle whose area is $\left(\frac{H}{2}\right)^2-\left(\frac{\varphi(S)}{2}\right)^2$.

NOTE : (a) $a'<b'<c'<d'$, $a''<b''<c''<d''$, ….., $a^{''\ldots e\ times}<b^{''\ldots e\ times}<c^{''\ldots.e\ times}<d^{''\ldots.e\ times}$

(b) let m = 2.3.5 then , groups are (1,2,3,5) , (1,2,3,6) , (1,2,3,10) , (1,2,3,15) , (1,2,3,30) , (1,2,5,6) , (1,2,5,10) , (1,2,5,15) , (1,2,5,30) , (1,2,6,10) , (1,2,6,15) , (1,2,6,30) , (1,2,10,15) , (1,2,10,30) , (1,2,15,30) , (1,3,5,6) , (1,3,5,10) , (1,3,5,15) , …….. , (6,10,15,30) . If m=2.3.5 , then total number of groups = 70 .
(c) $d_1=a'=1$.

Theorem 2.1 in another form :
A lot consists of groups for which E_Ψ,F_Ψ,G_Ψ form inverted right angle triangle, then probability (q_1) of getting the inverted right angle triangle whose area is $\left(\frac{H}{2}\right)^2-\left(\frac{\varphi(S)}{2}\right)^2$

is greater than or equal to $\frac{z-1+^{z-1}C_2}{w_4+w_5}$

Or, $q_1 \geq \frac{z-1+^{z-1}C_2}{w_4+w_5}$ and probability (q_2) of getting the inverted right angle triangle whose area is not equal to

$\left(\frac{H}{2}\right)^2 - \left(\frac{\varphi(S)}{2}\right)^2$ is less than or equal to $\frac{w_4+w_5-^{z-1}C_2-z+1}{w_4+w_5}$

Or, $q_2 \leq \frac{w_4+w_5-^{z-1}C_2-z+1}{w_4+w_5}$.

q_1 *can also be written as* ,

$$q_1 \geq \frac{z-1+^{z-1}C_2}{\sum_{i=i}^{n} i\ [\{\tau(m)-2-i\}\text{th triangular number}] - \sum_{j=1}^{3} w_j}$$

Note : If $m = p_1.p_2.p_3$ then ,

$q_1 \geq 0.1364$ *and* $q_2 \leq 0.8637$.

III. RESULTS RELATED TO HARDY-RAMANUJAN NUMBER , AREA OF RECTANGLE AND TRAPEZIUM AND GOLDEN RATIO .

1) Result 3.1: If $m = 3.5.7$, out of the 70 groups if we take 27th and 41th groups that are $(1,7,15,35)$ and $(3,5,15,35)$ and

if $E_{27}[HCF(1,7), LCM\ (1,7)]$,$F_{27}[HCF(7,15), LCM(7,15)]$, $G_{27}[HCF(15,35), LCM(15,35)]$

and $E_{41}[HCF(3,5), LCM(3,5)]$,

$F_{41}[HCF(5,15), LCM(5,15)]$,

$G_{41}[HCF(15,35), LCM(15,35)]$ are the vertices of triangles E_{27},F_{27},G_{27} and $E_{41}F_{41}G_{41}$, then the average of $\left(\frac{H_{27}}{2}\right)^2 - \left(\frac{\varphi(S_{27})}{2}\right)^2$

And $\left(\frac{H_{41}}{2}\right)^2 - \left(\frac{\varphi(S_{41})}{2}\right)^2$ is Hardy – Ramanujan number

Where $E_{27}F_{27}G_{27}$ is inverted right angle triangle , $E_{41}F_{41}G_{41}$ is non-inverted right angle triangle. H_{27},H_{41} are the length of hypotenuse of triangles $E_{27}F_{27}G_{27}$ and $E_{41}F_{41}G_{41}$


S_{27}=7x15 , S_{41}=5x15 and area of both triangle is not equal to $\left(\frac{H}{2}\right)^2 - \left(\frac{\varphi(S)}{2}\right)^2$.

$\frac{\left(\frac{H_{27}}{2}\right)^2-\left(\frac{\varphi(S_{27})}{2}\right)^2+\left(\frac{H_{41}}{2}\right)^2-\left(\frac{\varphi(S_{41})}{2}\right)^2}{2} = 1729$.

2) Result 3.2: If m=$P_1.P_2$ (P_1<P_2), divisors of m are 1,P_1,P_2,$P_1.P_2$.
Let E [HCF(1,P_1), LCM(1,P_1)],

F [HCF(P_1,P_2), LCM(P_1,P_2)], and

G [HCF(P_2,$P_1.P_2$), LCM (P_2,$P_1.P_2$)] are the vertices of triangle EFG



Let Area of Rectangle S'U'EF = R and Area of trapezium V'W' GE = K then,

(a) $\varphi(m) = R - \sqrt{H^2 - R^2}$

(b) $\varphi(m) = 2R - \sqrt{H^2 + 4(K - R)}$

where H is the length of hypotenuse of right angle triangle EFG .

Proof : $EG = \sqrt{(p_2 - 1)^2 + (p_1 . p_2 - p_1)^2}$

$= (p_2 - 1).\sqrt{1 + {p_1}^2}$

$EG = H = (p_2 - 1).\sqrt{1 + {p_1}^2}$

Area of Rectangle = S'U'EF = $R = p_1(p_2 - 1)$

Area of Trapezium = V'W'GE =

$K = \frac{1}{2} p_1(p_2 - 1)\ (p_2 + 1)$

Put $R = p_1(p_2 - 1)$ and $H = (p_2 - 1).\sqrt{1 + {p_1}^2}$ in (a) that is $R - \sqrt{H^2 - R^2}$, We get

$$p_1(p_2 - 1) - \sqrt{(p_2 - 1)^2(1 + {p_1}^2) - {p_1}^2(p_2 - 1)^2}$$

$$= p_1(p_2 - 1) - (p_2 - 1)$$

$$= (p_2 - 1)(p_1 - 1)$$

$$= (p_1 - 1)(p_2 - 1)$$

$$= \varphi(m)$$

Hence (a) part is proved.

Now, put $R = p_1(p_2 - 1)$, $H = (p_2 - 1)\sqrt{1 + {p_1}^2}$

And $K = \frac{1}{2} p_1(p_2 - 1)\ (p_2 + 1)$ in (b) that is

$2R - \sqrt{H^2 + 4(K - R)}$, we get

$=2p_1(p_2\text{-}1)\text{-}[(p_2\text{-}1)^2(1+{p_1}^2)+2\{p_1(p_2\text{-}1)(p_2+1)\text{-}2p_1(p_2\text{-}1)\}]^{1/2}$

$=2p_1(p_2-1) - [(p_2-1)^2(1+p_1^2)+2\,p_1(p_2-1)(p_2+1-2)]^{1/2}$

$=2p_1(p_2-1) - [(p_2-1)^2(1+p_1^2)+2\,p_1(p_2-1)^2]^{1/2}$

$=2p_1(p_2-1) - [(p_2-1)^2(1+p_1^2+2p_1)]^{1/2}$

$=2p_1(p_2-1) - (p_2-1)(p_1+1)$

$=(p_2-1)[2p_1-p_1-1]$

$=(p_2-1)(p_1-1)$

$=(p_1-1)(p_2-1)$

$=\varphi(m)$

Hence (b) part is proved.

3) *Result 3.3:* If $m=p_1.p_2$ where p_1,p_2 are the primes ($p_1<p_2$), divisors of m are $1,p_1,p_2,p_1.p_2$. let E [HCF$(1,p_1)$, LCM$(1,p_1)$], F [HCF(p_1,p_2), LCM(p_1,p_2)] and G [HCF$(p_2,p_1.p_2)$,LCM$(p_2,p_1.p_2)$] are the vertices of triangle EFG, then roots of the quadratic equation

$2.\,(\varphi(p_1))^2x^2 - 2(\varphi(p_1))^2x^1 - p_1 = 0$ are $\frac{\sqrt{H^2-4T}\pm H}{2\sqrt{H^2-4T}}$

Proof :- $2.\,(\varphi(p_1))^2x^2 - 2(\varphi(p_1))^2x^1 - p_1 = 0$

$x=\frac{(p_1-1)\pm\sqrt{1+p_1^2}}{2(p_1-1)}$

Since , $H = (p_2-1)\sqrt{1+p_1^2}$ and $T = \frac{1}{2}p_1(p_2\text{-}1)^2$

Where T is the area of triangle EFG

$$\frac{\sqrt{H^2-4T}\pm H}{2\sqrt{H^2-4T}} = \frac{(p_1-1)\pm\sqrt{1+p_1^2}}{2(p_1-1)}$$

Hence, Proved.

4) *Result 3.4:*

$$If\ m = p_1.p_2\ where\ p_1\ is\ either\ 2\ or\ 3\ and$$

$p_1 < p_2, Divisors\ of\ m\ are\ 1, p_1, p_2, p_1.p_2.$

$$Let\ E[HCF(1,p_1), LCM(1,p_1)], F[HCF(p_1,p_2), LCM(p_1,p_2)$$


And $G[HCF(p_2,p_1.p_2), LCM(p_2,p_1.p_2)]$ are the vertices of triangle EFG , then

$2.\lim_{n\to\infty}\left(\frac{U_{n+1}}{U_n}\right) - 1 = \sqrt{\frac{4T+(\varphi(m))^2}{\varphi(p_2).\varphi(m)}} = \frac{H}{\varphi(p_2).\sqrt{\varphi(p_1)}}$

$$where\ T\ is\ the\ area\ of\ triangle\ EFG.$$

Proof : $If\ m = p_1.p_2, then\ \varphi(m) = (p_1-1).(p_2-1),$

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538
Volume 10 Issue I Jan 2022- Available at www.ijraset.com



Vertices of above triangle EFG are $E(1,p_1)$, $F(1,P_1.P_2)$ and $G(P_2,P_1.P_2)$.

$$T = \frac{1}{2}.EF.FG$$

$$Or, T = \frac{1}{2}.p_1.(p_2-1)^2$$

$$Put\ T\ and\ \varphi(m)\ in\ R.H.S,$$

$$\sqrt{\frac{2p_1(p_2-1)^2 + (p_1-1)^2(p_2-1)^2}{(p_1-1)(p_2-1)^2}} = \sqrt{\frac{p_1^2+1}{p_1-1}}$$

(i) $If\ p_1 = 2, then \sqrt{\frac{p_1^2+1}{p_1-1}} = \sqrt{5}$

(ii) $If\ p_1 = 3, then \sqrt{\frac{p_1^2+1}{p_1-1}} = \sqrt{5}$

$$Since, for\ n \geq 1 \lim_{n\to\infty}\left(\frac{U_{n+1}}{U_n}\right) = \frac{1+\sqrt{5}}{2}$$

$$Or, 2.\lim_{n\to\infty}\left(\frac{U_{n+1}}{U_n}\right) - 1 = \sqrt{5}$$

Hence , $2.\lim_{n\to\infty}\left(\frac{U_{n+1}}{U_n}\right) - 1 = \sqrt{\frac{4T+(\varphi(m))^2}{\varphi(p_2).\varphi(m)}} = \frac{H}{\varphi(p_2).\sqrt{\varphi(p_1)}}$.

IV. ANGLES OF TRIANGLE EFG IN TERMS OF EULERS PHI FUNCTION

1) *Theorem 4.1:* If $m=p_1.p_2$ where p_1,p_2 are the primes ($p_1<p_2$), divisors of m are $1,p_1,p_2,p_1.p_2$. let E [HCF($1,p_1$), LCM($1,p_1$)], F [HCF(p_1,p_2), LCM(p_1,p_2)] and G [HCF($p_2,p_1.p_2$), LCM($p_2,p_1.p_2$)] are the vertices of triangle EFG and angle E = α and angle G =β, then the angles of triangle EFG (in terms of euler's phi function) are $\alpha = \frac{1}{2}cot^{-1}\left[\frac{\varphi(m)\sqrt{2H^2-(\varphi(m))^2}}{H^2-(\varphi(m))^2}\right]$ and

$\beta = \frac{1}{2}cot^{-1}\left[\frac{\varphi(m)\sqrt{2H^2-(\varphi(m))^2}}{(\varphi(m))^2-H^2}\right]$

©IJRASET: All Rights are Reserved

Proof : Put $\varphi(m) = (p_1 - 1)(p_2 - 1)$ and

$H = (p_2 - 1)\sqrt{1 + {p_1}^2}$ in

$\alpha = \frac{1}{2}cot^{-1}\left[\frac{\varphi(m)\sqrt{2H^2-(\varphi(m))^2}}{H^2-(\varphi(m))^2}\right]$,

$= \frac{1}{2}cot^{-1}\left[\frac{(p_1-1)(p_2-1)\{2(p_2-1)^2(1+{p_1}^2)-(p_1-1)^2(p_2-1)^2\}^{1/2}}{(p_2-1)^2(1+{p_1}^2)-(p_1-1)^2(p_2-1)^2}\right]$

$= \frac{1}{2}cot^{-1}\left[\frac{(p_1-1)\{2(1+{p_1}^2)-(p_1-1)^2\}^{1/2}}{\{(1+{p_1}^2)-(p_1-1)^2\}}\right]$

$= \frac{1}{2}cot^{-1}\left[\frac{(p_1-1)\{2+2{p_1}^2-{p_1}^2-1+2p_1\}^{1/2}}{1+{p_1}^2-{p_1}^2-1+2p_1}\right]$

$= \frac{1}{2}cot^{-1}\left[\frac{(p_1-1)\{{p_1}^2+2p_1+1\}^{1/2}}{2p_1}\right]$

$= \frac{1}{2}cot^{-1}\left[\frac{(p_1-1)(p_1+1)}{2p_1}\right]$

$= \frac{1}{2}cot^{-1}\left[\frac{{p_1}^2-1}{2p_1}\right] = \frac{1}{2}tan^{-1}\left[\frac{2p_1}{{p_1}^2-1}\right]$ -------(∗)

Since, tanα = $\frac{GF}{EF}$

Or, tanα = $\frac{p_2-1}{p_1(p_2-1)} = \frac{1}{p_1}$

From (∗) equation,

$$\frac{1}{2}tan^{-1}\left[\frac{2p_1}{{p_1}^2-1}\right] = \alpha$$

Or, $tan(2\alpha) = \frac{2p_1}{{p_1}^2-1}$

Or, $\frac{2tan\alpha}{1-tan^2\alpha} = \frac{2p_1}{{p_1}^2-1}$

Put $\alpha = \frac{1}{p_1}$,

$$\frac{2x\frac{1}{p_1}}{1-\frac{1}{{p_1}^2}} = \frac{2p_1}{{p_1}^2-1}$$

L.H.S = R.H.S

Hence, Proved.

Since, $\alpha + \beta = \frac{\pi}{2}$

$$\frac{1}{2}tan^{-1}\left[\frac{2p_1}{{p_1}^2-1}\right] + \beta = \frac{\pi}{2}$$

$$\beta = \frac{\pi}{2} - \frac{1}{2}tan^{-1}\left[\frac{2p_1}{{p_1}^2-1}\right]$$

$$\beta = \frac{1}{2}\left[\pi - tan^{-1}\left\{\frac{2p_1}{{p_1}^2 - 1}\right\}\right]$$

Or, $\beta = \frac{1}{2}\left[\pi - cot^{-1}\left\{\frac{{p_1}^2-1}{2p_1}\right\}\right]$

$$\beta = \frac{1}{2}\left[cot^{-1}\left\{\frac{1-{p_1}^2}{2p_1}\right\}\right]$$

Hence, $\beta = \frac{1}{2}cot^{-1}\left[\frac{\varphi(m)\sqrt{2H^2-(\varphi(m))^2}}{(\varphi(m))^2-H^2}\right]$

V. OBSERVATIONS

(5.1) If $m = 2.3.5$, then $w_4 = 18$, $w_5 = 4$, and $w_4 + w_5 = 22$.
If m= 3.5.7 , then w_4=19, w_5=3, and w_4+w_5=22
So, $w_4 + w_5$ is constant but w_4, w_5 are not constant.
If m=$p_1.p_2.p_3$, for different m w_1, w_2, w_3 and w_4+w_5 are constants.
$w_1 = 10, w_2 = 36, w_3 = 2$ and $w_4 + w_5 = 22$.

$$w_2 > w_4 + w_5 > w_1 > w_3.$$

Total number of right angle triangles = $w_3 + w_4 + w_5$

$= \varphi(\sum_{k=1}^{5} w_k) = \varphi(8_{C_4}) =$

$\varphi[\sum_{i=i}^{5} i\ [\{\tau(m) - 2 - i\}th$ triangular number]] = 24 .

Or, $w_3 + w_4 + w_5 = \varphi(w_1 + w_2 + w_3 + w_4 + w_5)$

Or, $\sum_{k=1}^{5} w_k - \varphi(\sum_{k=1}^{5} w_k) = w_1 + w_2$

That is $70 - 24 = 10 + 36$

From Theorem 2.1 : $w_5 \geq z - 1 + {}^{z-1}C_2$

If $m = p_1.p_2.p_3$, then,

$\varphi(\sum_{k=1}^{5} w_k) \geq z - 1 + {}^{z-1}C_2 + w_3 + w_4$, Where $\sum_{k=1}^{5} w_k$ = Total No. of groups.

And $\varphi(\sum_{k=1}^{5} w_k) - z + 1 - {}^{z-1}C_2 - w_3 \geq w_4 \geq \frac{z-1+{}^{z-1}C_2}{q1} - w_5$. where q_1 is the probability of getting the inverted right angle triangle whose area is $\left(\frac{H}{2}\right)^2 - \left(\frac{\varphi(S)}{2}\right)^2$

VI. APPLICATION OF EULER'S TOTIENT FUNCTION AND NUMBER THEORETIC FUNCTIONS IN STUDENT'S PENCIL COMPASS

Mathematical instruments are used to understand mathematical constructions and concepts . If we talk about mathematical instruments then there are many mathematical instruments like protractor , ruler , set-square , divider , pencil compass and etc . There are two types of mathematical instruments , one is used by student and other are used by teachers . Mathematical instruments used by teachers are very large as compare to mathematical instruments used by student because they are used in black boards or white boards . The mathematical instruments used by student are smaller because they are usable in their books only . If we talk about the student's pencil compass , then one can form the circles with a radius of about 1 to 15 centimeters .

Here we have applied the concepts of number theory .

If it is assumed that a circle of radius up to 20 centimeter is being formed by student's pencil compass then also the following result will be correct . Normally circles of radius up to 14 or 15 centimeters are formed by the student's pencil compass .

VII. RESULT

If we draw all the possible circles by student's pencil compass having radius = r (r = 1,2,3,4,……,maximum possible radius to be made by student's pencil compass) , then 6 is the only number whose every positive proper divisors as a radius follow the relation

$\varphi(2r)$ + τ(2r) + σ(2r) = greatest integer function of P

and the number 6 (itself) as a radius follow the relation

$\varphi(2r)$ + τ(2r) + σ(2r) = lowest integer function of P .

$\varphi(2r)$ + τ(2r) + σ(2r) can also be written as $\varphi(d)$ + τ(d) + σ(d) ,

where P = π x d , π = 3.1415 (four digits after the decimal point) , P is circumference(perimeter) of circle and d is diameter of circle .

NOTE: (a) Draw all the possible circles by student's pencil compass means to make circles of each radius which is made by student's pencil compass .

Where $\varphi(d)$ is the is the euler's phi function which denote the number of positive integers not exceeding d that are relatively prime to d .

τ(d) denote the number of positive divisors of d and σ(d) denote the sum of these divisors .
In mathematical form : If r = 1 , $\varphi(2)$ + τ(2) + σ(2) = greatest integer function of 2 x 3.1415 x 1 = 6 ,
If r = 2 , $\varphi(4)$ + τ(4) + σ(4) = greatest integer function of 2 x 3.1415 x 2 = 12 ,
If r = 3 , $\varphi(6)$ + τ(6) + σ(6) = greatest integer function of 2 x 3.1415 x 3 = 18 ,
If r = 6 , $\varphi(12)$ + τ(12) + σ(12) = lowest integer function of 2 x 3.1415 x 6 = 38 ,
Where 1,2 and 3 are the positive proper divisors of 6 .

VIII. ACKNOWLEDGEMENT

I would like to thanks Dr. Pravin Hudge , Dr. V. Patil , Dr. Vikas Deshmane , and Mandar Khasnis sir for useful comments .


REFERENCES

[1] Gabor J. Szekely , contests in higher mathematics miklos Schweitzer competitions 1962-1991, page number 244 geometry (Balazs csikos)
[2] Elementary number theory by David M. Burton , TATA MC GRAW – HILL Edition .
[3] Pythagoreans introduction to number theory Ramin Takloo-Bighash , Bartolini springer international publishing , 2018 .
[4] Grade school triangles Jack Scalcut , The American Mathematical Monthly 117(8) , 673-685 , 2010 .
[5] Monthly research problems , 1969-73 Richard K Guy , The American Mathematical Monthly 80(10) , 1120-1128, 1973 .
[6] E.Grosswald , The average order of an arithmetic function , Duke mathematics journal 23(1956) , 41-44 .
[7] To count clean triangles we count on imph(n) , Mizan R Khan , Riaz R Khan arxiv preprint arxiv : 2012.11081 , 2020 .
[8] (2002) Number Theory . In : Mathematics problems and proofs , Springer , Boston , MA.https://doi.org/10.1007/0-306-46963-4_3 .
[9] An Introduction to the Theory of Numbers:G.H. Hardy,Edward M.Wright ,Andrew Wiles , Roger Heathbrown , Joseph Silverman.
[10] An Introduction to the Theory of Numbers Book : Niven , Zuckerman , Montgomery .
[11] A Classical Introduction to Modern Number Theory (Graduate Texts in Mathematics) .
[12] Handbook of mathematical , Scientific and engineering formulas , Tables functions , Graphs , Transforms max fogiel; Research and Education Association , Research and Education Assoc; 1984.
[13] A text book of algebra F.Y.B.Sc. / B.A. (semester – 1) , Sheth Publishers .
[14] An Introduction to the Theory of Numbers Book : Niven , Zuckerman , Montgomery .
[15] Wolfram Web Resources , Eric Weisstein .
[16] Mathematical instrument , WIKIPEDIA ,Gerard L'Estrange Turner Scientific Instruments , 1500-1900:An Introduction (university of California Press , 1998) .
[17] COBB,PAUL.1999. "Individual and Collective Mathematical Development : The case of statistical Data Analysis" . Mathematical Thinking and Learning 1 (1):5-43 .
[18] FENNEMA , ELIZABETH , and ROMBERG , THOMAS , eds.1999. Mathematics Classrooms That Promote Understanding . Mahwah , NJ:Erlbaum .

ABOUT AUTHOR

I am a student (undergraduate student) of third year (T.Y B.Sc) , C.H.M college ,Ulhasnagar . I completed my first year and second year (F.Y,S.Y B.Sc) from SICES degree college of arts , science , and commerce, Ambarnath(w) .

©IJRASET: All Rights are Reserved | SJ Impact Factor 7.538 | ISRA Journal Impact Factor 7.894 |



10.22214/IJRASET

45.98

IMPACT FACTOR: 7.129

IMPACT FACTOR: 7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)