



# **iJRASET**

International Journal For Research in  
Applied Science and Engineering Technology



---

# **INTERNATIONAL JOURNAL FOR RESEARCH**

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume: 13    Issue: V    Month of publication: May 2025**

**DOI: <https://doi.org/10.22214/ijraset.2025.70483>**

**[www.ijraset.com](http://www.ijraset.com)**

**Call:  08813907089**

**E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)**

# Formulation and Evaluation of Giloy Berberine-Loaded Hydrocolloidal Patches for Acne Treatment

Divya Borade<sup>1</sup>, Sanskruti Bambale<sup>2</sup>, Prajwali Bhalerao<sup>3</sup>

<sup>1,2</sup>Student, <sup>3</sup>Assistant Professor; (Pharmacognosy) Sinhgad Institute Of Pharmaceutical Sciences, Lonavala



**Abstract:** Giloy, scientifically known as *Tinospora Cordifolia*, is a widely recognized medicinal plant in Ayurveda. It is a deciduous climbing shrub with heart-shaped leaves belonging to Menispermaceae family, acting as one of the main sources of new pharmaceuticals and health care products. The plant is genetically diverse, containing different active components, including steroids, aliphatics, alkaloids, glycosides, and diterpenoid lactones. This herb possesses anti-inflammatory, antibacterial, immunomodulating agent, anti-viral, anti-diabetic properties in addition to being a strong antioxidant.<sup>1</sup> A primary method for determining the antimicrobial activity of a compound and its active substance was carried out on invitro model through agar well diffusion method on *Propionibacterium Acne*. Observations of antibacterial activity were carried out by reviewing clear zones indicating the inhibition zones.<sup>2</sup> Studies revealed that *T. cordifolia* is an excellent drug and does not have any adverse or toxic effects till now. There is a new invention for topical acne treatment called acne patch. The purpose of this study was to formulate an acne patch preparation from Giloy Stem extract with a combination of Xanthan gum and Propylene glycol and other excipients against *Propionibacterium acnes* bacteria. Acne patch preparation was evaluated for physical characteristics, and tested for its antibacterial effectiveness activity.


**Keywords:** Giloy, *P. acne*, Hydrocolloidal Patches, *T.Cordifolia*, Berberine, Xantham Gum.

## I. INTRODUCTION

Patches as a topical acne treatment:

Acne is commonly affecting many teenagers that makes so many scientists are exploring many formulas for acne treatment. There is a new invention for topical acne treatment called acne patch. In general, acne patches that are already on the market contain hydrocolloid synthesizing substances.<sup>3</sup> Acne patches on the market are usually made of hydrocolloids or hydrogels. Hydrocolloid dressings consist of two layers, namely, a colloid layer and a water impermeable layer. The so-called colloid layer is the inner layer and the water-impermeable layer is the outer layer. The function of the water-impermeable layer is that it can provide a protective layer and can help prevent the spread of pathogenic microorganisms, Based on the type of acne, acne patches are divided into several types:

Sr.no.	Acne Patches	Characteristics	
1	Microneedle patches	<ul style="list-style-type: none"> <li>For cystic or nodular acne</li> </ul> Containing microneedle that is dissolved and is very fine, has a small needle on one side  Can help deliver and penetrate the active ingredients into the deeper layers of the skin	 <p>Fig. 1 Microneedle patch</p>
2	Hydrogel patches	For active acne contains active ingredients to kill acne-causing bacteria and reduce inflammation. Can reduce bumps, pain, and redness in pimples, and can relieve inflamed pimples such as papules. Can also help reduce the size of lesions due to nodular or cystic acne.	 <p>Fig. 2 Hydrogel Patch</p>

3	Hydrocolloid patches	<p>For yellow or white pimples. Often used to speed up the healing process. Has the size of a pimple and is circular in shape. The shape is also very thin so that when used in public it is less visible. Can suck moisture from the pores, prevent other infections, and can prevent the formation of acne scars on the skin.</p>	 <p>Fig. 3 Hydrocolloid Patch</p>
---	----------------------	---	--

**A. Hydrocolloidal Patches:**

Hydrocolloid acne patches primarily target the Epidermis , specifically its outermost layers.

Targeted Skin Layer:

Stratum Corneum( *the outermost layer of the epidermis*)

- This is where the patch adheres and forms a moist environment.
- It absorbs fluid/ pus from whiteheads or popped pimples.
- Helps isolate the wound from bacteria, dirt, and friction.
- Hydrocolloid patches are is non medicated acne patch used to treat acne.
- The function of this patch is to give skin hydration and improve the skin barrier.

The Permeation mechanism of Hydrocolloid patches is passive and absorptive, targeting superficial acne lesions like whiteheads and pustules.

Particularly effective for surface-level acne (not deep cystic acne).

Can flatten pimples overnight by drawing out pus and reducing inflammation.

No chemical interaction with deeper layers of the skin.

**II. TINOSPORA CORDIFOLIA (GILOY)**

Classification of Plant:

Biological Source: Giloy is a deciduous climbing shrub that is tropical to Indian subcontinents.

Species: Cordifolia

Family: *Menispermaceae*

Kingdom: Plantae

Genus: Tinospora

Synonyms: Gulvel, Guduchi, Giloy



Fig. 4 T.Cordifolia

**III. CHEMICAL CONSTITUENTS**

Various chemical constituents have been found in different parts of the Giloy plant. They belongs to different classes such as alkaloids, terpenoids, steroids, glycosides, polysaccharides, flavonoids.

Sr. No.	Active compound type	Biological Source	Compounds
1.	Alkaloids (4-6%)	Stem, root	Berberine, palmatine, jatrorrhizine
2.	Glycosides (10- 15%)	Leaf, root	Cordifolioside, tinosporaside



3.	Terpenoids (5-7%)	Flowers	Beta sitosterol, caryophyllene
4.	Flavonoids (3-5%)	Leaf, flower, stem	Quercetin, luteolin, kaempferol
5.	Polysaccharides (20-30%)	Stem, leaves, root	Arabinose, galactose, glucose, xylose
6.	Steroids	Stem	B-Sitosterol, Giloinsterol, Makisterone A

#### IV. MATERIAL AND METHODS

##### I. Materials:

An aerial stem of *Tinospora cordifolia* was collected and authenticated.

*Propionobacterium Acnes* Bacterial Culture, Agar medium.

##### II. List of Chemicals:

Sr.No.	Chemicals
	Ethanol 70%
	Xanthan gum
	Propylene glycol
	Tea tree oil
	Aloevera gel
	Triethanolamine
	Distilled water

##### III. List of Instruments:

Sr. No.	Name of Equipment	Purpose
1	Weighing balance	Measurement of weight
2	Magnetic stirrer	Mixing and Agitation
3	Vernier Calliper	Measuring Thickness
4	Incubator	Maintain growth and development of microorganisms
5	Autoclave	Sterlization
6	Hot air oven	Drying
7	Dessicator	Determine Weight uniformity
8	Digital pH meter	Determine pH
9	UV Spectrophotometer	Quantification, Purity determination, and identification
10	Sieve	Size separation and ensure size uniformity

#### V. PROCEDURE FOR PREPARATION OF PATCH

*T.Cordifolia* stem was collected, cleaned and washed with water. Then, the stems were dried in a tray dry




Fig. 5 Dried stem of T.Cordifolia



Dried stems are coarsely ground; sifting was carried out using a 40-mesh sieve, weighed, and stored in airtight jars.



Fig. 6 Trituration of dried stem



The extract was prepared using the cold maceration method, and 100 ml ethanol was added to 100 g powdered *T. Cordifolia* at room temperature for 48 hrs.



Fig. 7 Extraction Process



Ethanol extract of *T. Cordifolia* was filtered, collected and stored.



Fig. 8 Filtration

The formulation of Patch begins with mixing of Xanthan gum and aloe vera gel until Homogenous mixture was formed. The ethanolic extract of plant and other excipients were added and mixed using magnetic stirrer.



Fig. 9 Dissolution of ingredients



Once the mixture was formed, it was spread uniformly on a Petri dish and without disturbance kept for evaporation at room temperature for 24 hrs.



Fig. 10 Patch formulation




Fig. 11 Final Formulated Patch

### VI. EVALUATION PARAMETERS

#### 1) Physical evaluation of extract:

Physical assessment (color, appearance) of *T. Cordifolia* (Giloy) ethanol extract was performed.

Parameter	Observation
Color	Light Brown
Appearance	Powdered Form

#### 2) Preliminary Phytochemical Screening:

Phytochemical screening of *T. Cordifolia* stem (ethanolic extract) was performed for the presence of alkaloids.

Sr. No.	Tests for Alkaloids	Methodology	Observations
1)	Dragendorff's Test	2-3 ml filtrate + few drops of Dragendorff's reagent	Orange brown ppt
2)	Mayer's Test	2-3 ml filtrate + few drops of Mayer's reagent	Gives ppt
3)	Hager's Test	2-3 ml filtrate + few drops of Hager's reagent	Yellow ppt
4)	Wagner's Test	2-3 ml filtrate + few drops of Wagner's reagent	Reddish brown ppt



Fig. 12 Reagents used for test



Fig. 13 Phytochemical test results

#### COLUMN CHROMATOGRAPHY

- Column Preparation
- Sample loading
- Elution with gradient solvent
- Fraction Collection
- Evaporate to get pure compound (Rotary Vacuum Evaporator)
- Monitoring with TLC



Fig. 14 Column Chromatography

**TLC (THIN LAYER CHROMATOGRAPHY):**

Thin layer chromatography (TLC) was performed on the 20 cm × 20 cm plates percolated with microcrystalline cellulose. Extracts are spotted on the plates, and TLC is allowed to run until the mobile phase reaches three-fourths of the TLC length.<sup>[22]</sup> TLC was performed using a hexane-to-ethyl acetate ratio of 4.3:1.8.




Fig. 15 TLC plate

$$R_f \text{ value} = \frac{\text{Distance travelled by solute}}{\text{Distance travelled by solvent}}$$

$$\text{Solute (Berberine)} = 2.8 \text{ cm}$$

$$\text{Solvent front} = 6.5 \text{ cm}$$

$$R_f \text{ value} = \frac{2.8}{6.5} = 0.43$$

**IN VITRO TESTING:**

Anti-acne activity assay of *T. Cordifolia* stem ethanol extract

The anti-acne activity of *T. Cordifolia* stem ethanol extract against *P. acnes* was tested at concentrations of 50, 100, 150, and 200 mg using the agar well diffusion method. A primary method for determining the antimicrobial activity of a compound and active substance. This method was carried out by making holes in the agar and inoculating test bacteria on Petri dishes containing agar. After that, samples were added to each hole in the cup at a predetermined concentration and incubated at 37°C for a day. Observations of antibacterial activity were carried out by reviewing clear zones indicating the inhibition zones. The inhibitory diameter was observed after incubation, and the results were recorded<sup>15</sup>

Sr. No.	Sample	Concentration (mg/ml)	Zone of inhibition (mm)
1)	<i>T. Cordifolia</i> stem extract	50	9
		100	12
		150	15
		200	18




Fig. 16 Anti-acne activity

**PREPARATION OF PATCH FORMULATION:**

*F(0)* = formula without extract, *F(1)* = formula with extract, and *F(2)* = formula with extract.

Xanthan gum and aloe vera gel were mixed until homogeneous mixture was formed then ethanolic extract (*T. Cordifolia*) were added, previously dissolved with 70% ethanol.



Tea tree oil and propylene glycol were then added. Once the mixture was uniform, the remaining distilled water was gradually added while stirring. Place in a Petri dish and let evaporate at room temperature for 24 hrs.

Ingredients	Functions	F(0)	F(1)	F(2)
<i>T. Cordifolia</i> Extract	Antibacterial	-	0.2 g	0.4 g
Xanthan Gum	Polymer	1 g	1 g	1.7 g
Aloe vera Gel	Humectant	0.5 g	1.7 g	2.3 g
Tea tree oil	Anti-microbial, Flavoring agent	0.12 g	0.2 g	0.4 g
Propylene Glycol	Humectant	0.5 g	1 g	1.5 g
Distilled Water	Solvent	10 ml	30 ml	40 ml
Triethanolamine	pH stabilizer	0.02 g	0.02 g	0.04 g

#### IV. EVALUATION PARAMETER OF PATCH

##### A. Folding resistance

The patch is repeatedly folded the same way for the folding resistance test until broken. Then, the number of folds is considered the value of the resistance to the folding of a patch.

##### B. Thickness of patch

The patch thickness tester is used to measure the thickness individually for each of the four patch formulations to test the thickness of the patch. Measurements are taken at three different points for each formulation.

##### C. Moisture absorption

After the patch was weighed, it was kept for 24 h at room temperature. After that, it was stored at 40°C for 24 h and weighed again.<sup>[11]</sup> The formula for % Humidity to determine the percentage of moisture absorption is as follow:

$$\% \text{ Humidity} = \frac{\text{initial weight} - \text{final weight}}{\text{initial weight}} \times 100\%$$

##### D. pH test

The pH of prepared patches was determined using a pH meter.

The pH of patch was found to be 6.

##### E. Drug content

The drug release profile of a herbal patch made with *Tinospora cordifolia*, xanthan gum, propylene glycol, tea tree oil, Aloe vera, and triethanolamine, needed in-vitro studies (e.g., Franz diffusion cell method) and data analysis.


Time (min)	Absorbance (AU)	Concentration (µg/ ml)	Cumulative release(%)
0	0.000	0.00	0%
1	0.025155	0.24	2.4%

2	0.045	0.43	4.3%
5	0.095	0.92	9.2%
10	0.155	1.50	15%
20	0.230	2.22	22.2%
30	0.315	3.05	30.5%
45	0.420	4.06	40.6%
60	0.510	4.95	49.5%
90	0.645	6.26	62.6%
120	0.755	7.34	73.4%

Sr. No.	Sample	Concentration (mg/ml)	Zone of inhibition(mm)
1)	<i>T. Cordifolia</i> stem extract	50	9
		100	12
		150	15
		200	18

**VII.RESULT**

Sr. No.	Evaluation Parameters	Observation
1)	Folding resistance	After 7 folds, patch adhesion started to peel off and after 15 folds the material showed tearing signs
2)	Thickness of patch	0.25 mm
3)	Moisture absorption	15.15%
4)	pH	6
5)	TLC	0.43



## VIII. DISCUSSION

Current study was undertaken to investigate the antimicrobial property of *T. cordifolia* that could serve as novel futuristic plant-based antibacterial agent for acne management. Berberine is the main alkaloid constituents of the plant used for its antimicrobial property. A primary method for determining the antimicrobial activity of a compound and its active substance was carried out by invitro model through agar well diffusion method on P.Acne. Hydrocolloidal patches are chosen because they often used to speed up the healing process, it has the size of a pimple and is circular in shape. The shape is also very thin so that when used in public it is less visible. And its main property is that it sucks moisture from the pores, prevent other infections, and can prevent the formation of acne scars on the skin. Study shown that herbal Hydrocolloidal patches has not yet been discovered.

## IX. CONCLUSION

*Tinospora cordifolia* holds significant potential as a natural ingredient in cosmetics due to its numerous therapeutic properties. The stem extract of *T. cordifolia* has shown antibacterial potential against major acne causing bacteria *P.acne*. The aim of the current research work is to prepare herbal acne patches by using herbal and natural ingredients. Formulation were prepared and evaluated for organoleptic properties and other evaluation parameters including pH, thickness, moisture absorption, folding resistance. Organoleptic tests show that F1 and F2 are transparent, while F3 is little bit light brown. All patch formulations have a thickness of 0.22-0.25mm, folding endurance 7-15 times, moisture absorption of 15.5%. Studies revealed that *T. cordifolia* is an excellent drug and does not have any adverse or toxic effects till now.

## REFERENCES

- [1] Promila, S. S., & Devi, P. (2017). Pharmacological potential of *Tinospora cordifolia* (Willd.) Miers ex hook. & Thoms.(Giloy): A review. *Journal of pharmacognosy and phytochemistry*, 6(6), 1644-1647.
- [2] Somalwar, S., & Somalwar, A. R. (2022). *Tinospora Cordifolia*: A Novel Antimicrobial Agent. *IJRB AT*, 46-51.
- [3] Kapao, N., & Wattanutchariya, W. (2018). Development of natural acne patch from local materials using quality function deployment technique. In *MATEC Web of Conferences* (Vol. 192, p. 01050). EDP Sciences.
- [4] Kuo, C. W., Chiu, Y. F., Wu, M. H., Li, M. H., Wu, C. N., Chen, W. S., & Huang, C. H. (2021). Gelatin/chitosan bilayer patches loaded with cortex *Phellodendronamurense*/Centella asiatica extracts for anti-acne application. *Polymers*, 13(4), 579.
- [5] Zhang, T., Sun, B., Guo, J., Wang, M., Cui, H., Mao, H., ... & Yan, F. (2020). Active pharmaceutical ingredient poly (ionic liquid)-based microneedles for the treatment of skin acne infection. *Acta Biomaterialia*, 115, 136-147.
- [6] Mandal, A., Clegg, J. R., Anselmo, A. C., & Mitragotri, S. (2020). Hydrogels in the clinic. *Bioengineering & translational medicine*, 5(2), e10158.
- [7] Wu, M., Peng, Q. Y., Han, L. B., & Zeng, H. B. (2021). Self-healing hydrogels and underlying reversible intermolecular interactions. *Chinese Journal of Polymer Science*, 39(10), 1246-1261.
- [8] De Paepe, K., Wibaux, A., Ward, C., & Rogiers, V. (2009). Skin efficacy and biophysical assessment of glycerol-containing hydrocolloid patches. *Skin Pharmacology and Physiology*, 22(5), 258-265.
- [9] Palange, A., Barge, M., & Naikwadi, N. (2024). Review in Acne Vulgaris along with its treatment. *Asian Journal of Pharmaceutical Research and Development*, 12(3), 192-196.
- [10] "Histological and Anatomical Studies of *Tinospora cordifolia* (Willd.) Miers: A Medicinal Plant", Authors: M. B. Sharma, P. S. Chavan, and A. K. Singh, Publication: *Indian Journal of Plant Sciences*, Volume: 7, Issue: 3, Pages: 223-230, Year: 2021
- [11] Singh, K., Panghal, M., Kadyan, S., Chaudhary, U., & Yadav, J. P. (2014). Antibacterial activity of synthesized silver nanoparticles from *Tinospora cordifolia* against multi drug resistant strains of *Pseudomonas aeruginosa* isolated from burn patients. *Journal of Nanomedicine & Nanotechnology*, 5(2), 1.
- [12] Nipanikar, S. U., Chitlange, S. S., & Nagore, D. (2017). Pharmacological evaluation of hepatoprotective activity of AHPL/AYTAB/0613 tablet in carbon tetrachloride-, ethanol-, and paracetamol-induced hepatotoxicity models in Wistar albino rats. *Pharmacognosy research*, 9(Suppl 1), S41.
- [13] Zinjarde, S. S., Bhargava, S. Y., & Kumar, A. R. (2011). Potent  $\alpha$ -amylase inhibitory activity of Indian Ayurvedic medicinal plants. *BMC complementary and alternative medicine*, 11(1), 1-10.
- [14] Gupta, R., & Sharma, V. (2011). Ameliorative effects of *Tinospora cordifolia* root extract on histopathological and biochemical changes induced by aflatoxin-B1 in mice kidney. *Toxicology international*, 18(2), 94.
- [15] Jantarat, C., Sirathanarun, P., Chuchue, T., Konpian, A., Sukkua, G., & Wongprasert, P. (2018). In vitro antimicrobial activity of gel containing the herbal ball extract against *Propionibacterium acnes*. *Scientia pharmaceutica*, 86(1), 8.
- [16] Development of natural acne patch from local materials using quality function deployment technique Nattawadee Kapao and Wassanai Wattanutchariya
- [17] Formulation Study of Tea Tree Oil Patches Paola Minghetti , Antonella Casiraghi1 , Francesco Cilurzo , Veniero Gambaro and Luisa Montanari
- [18] Formulation and evaluation of polyherbal anti-acne combination by using in-vitro model Pranav Kumar Prabhakar 1 , Deepak Nath 2, Saurabh Singh 2 , Amit Mittal 2 , Dileep Singh Baghel 2
- [19] Patches For Acne Treatment: An update on the Formulation And Stability Test Taqiyyah Qothrunnadaa' , Aliya Nur Hasanah.
- [20] Antibacterial and qualitative phytochemical analysis of Giloy extract for application of herbal finish on cotton fabric K Medhaa , N Aryaa ,\*, K Malikb , S Ahlawatc & N Chauhan
- [21] Studies on Antioxidant Activity and Total Phenolic Content of *Tinospora cordifolia* (Miers.) Stem Using in Vitro Models Neha Upadhyay, Showkat Ahmad Ganie\*, Rajneesh Agnihotri, Rajendra Sharma



- [22] Formulation of antiacne serum based on lime peel essential oil and in vitro antibacterial activity test against Propionibacterium acnes Noor Fitri; Ifat Fatimah; Lutfi Chabib; Febi Indah Fajarwati
- [23] Tinospora cordifolia (Giloy): An insight on the multifarious pharmacological paradigms of a most promising medicinal ayurvedic herb Abhishek Gupta a,b,\* , Priyanka Gupta b , Gunjan Bajpai
- [24] Tinospora cordifolia (Giloy Or Amrita): An Updated Insight On The Multifarious Pharmacological And Pharmacotherapeutic Paradigms Of A Most Promising And Miraculous Medicinal Ayurvedic Herb Saurabh Saklani Jatin Mehra<sup>1</sup> , Garima Bisht<sup>1</sup> , Sanjay Rishishwar<sup>2</sup> , Poonam Rishishwar<sup>2</sup>





10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)