iJRASET
International Journal For Research in Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 11 **Issue:** III **Month of publication:** March 2023

DOI: https://doi.org/10.22214/ijraset.2023.49282

www.ijraset.com

Call: 08813907089 | E-mail ID: ijraset@gmail.com

International Journal for Research in Applied Science & Engineering Technology (IJRASET)
ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538
Volume 11 Issue III Mar 2023- Available at www.ijraset.com

Generalized Investigated in Topological Space

Kharatti Lal
Dept. of Applied Science – Mathematics, Govt. Millennium polytechnic College Chamba, CHAMBA - Himachal Pradesh – 176310 (INDIA)

Abstract: ***In this paper, some characterizations and proportion of notion a investigated. Throughout this paper (X, τ) and (Y, σ) (simply, X and Y) represent topological spaces on which separation axioms are assumed unless otherwise mentioned. We introduce a new class of sets called regular generalized open sets which is properly placed in between the class of open sets and the class of - open sets. Throughout this paper (X, τ) represents a topological space on which no separation axiom is assumed unless otherwise mentioned. For a subset A of a topological space X, cl (A) and int (A) denote the closure of A and the interior of A respectively. X/A or Ac denotes the complement of A in X. introduced and investigated semi open sets, generalized closed sets, regular semi open sets, weakly closed sets, semi generalized closed sets , weakly generalized closed sets, strongly generalized closed sets, generalized pre - regular closed sets, regular generalized closed sets, and generalized α-generalized closed sets respectively.***

Keywords: ***Topological space, Cluster Point, Open and Closed set, β* -*** **Continuous, Subset, .Regular open closed set, Separation axioms**

I. INTRODUCTION

A function $f : (X, \tau) \to (Y, \sigma)$ is called super-continuous (resp. *a*-continuous α-continuous pre-continuous δ - smi - continuous Z - continuous γ - continuous continuous Z^*- continuous , β - continuous e^*- continuous) if $f^{-1}(V)$ is δ - open (resp. *a*-open, α-open, per open, δ-semiopen, Z-open, γ-open, *e*- open, Z^* - open, β-open, e^*-open) in X, for each $V \in \sigma$. the notion of β-open sets and β-continuity in topological space. The concepts of Z^*- open set and Z^*- continuity introduced by Mubarki. The purpose of this paper introduce and study the notions of β^*- open sets, β^* - continuous functions and (β^*- open sets. For a subset A of a (X, τ), $cl(A)$, $int(A)$ and $X \setminus A$ denote the closure of A, the interior of A and the complement of A, respectively. A subset A of a topological space (X, τ) is called regular open (resp. regular closed) if $A = int(cl(A))$ (resp. $A = cl(int(A))$). A point x of X is called δ - cluster point of A if $int(cl(U)) \cap A = \phi$, for every open set U of X containing x.
The set of all δ-cluster points of A is called δ-closure of A and is denoted $cl\ \delta(A)$. A set A is δ-closed if and if $A = cl\ \delta(A)$. The complement of a *a*-open (resp. α-open, δ-semiopen, δ-preopen, Z-open, γ-open, *e*-open, Z^*-open, β-open, e^*-open) sets is called *a*-close (resp. α-closed δ-semi-closed , δ-pre-closed Z-closed γ-closed , *e* - closed Z^*-closed β-closed , e^*- closed The intersection of all δ - preclosed (resp. β-closed) set containing A is called the δ - preclosure (resp. β-closure) of A and is denoted by δ - *pcl* (*A*) (resp. β-*cl*(*A*)). The union of all δ-preopen (resp. β-open) sets contained in A is called the δ- pre - interior (resp. β - interior) of A and is denoted by δ-*pint*(*A*) (resp. β-*int*(*A*)). The family of all δ-open (resp. δ-semiopen, δ- preopen, Z^*- open, β-open, e^*- open) sets is denoted by $\delta O(X)$ (resp. $\delta SO(X)$, $\delta PO(X)$, $Z^* O(X)$, $\beta O(X)$, $e^* O(X)$).

Lemma 1.1
Let A be a subset of a space (X, τ). Then:
(1) δ -*pint*(*A*) = $A \cap int(cl\delta(A))$ *and* δ-*pcl*(*A*) = $A \cup cl\ (int\ \delta(A))$,
(2) $\beta - Int(A) = A \cap cl\ (int(cl(A)))$ *and* $\beta - cl\ (A) = A \cup int\ (cl(int\ (A)))$.

β* - ***Open sets***

Definition 2.1
A subset A of a topological space (X, τ) is said to be:
(1) a β^* - open set if $A \subseteq cl(int(cl(A))) \cup int(cl\delta(A))$,
(2) a β^* - closed set if $in\ t\ (cl(int(A))) \cap cl\ (int\ \delta\ c\ (A)) \subseteq A$.
(3) The family of all β^* - open (resp. β^*- closed) subsets of a space (X, τ) will be as always denoted by $\beta^* O(X)$ (resp. $\beta^* C(X)$).

©IJRASET: All Rights are Reserved

Remark 3.1

The following diagram holds for each a subset A of X.

None of these implications are reversible as shown in the following examples

Example 4.1

Let $X = \{a, b, c, d\}$, with topology $\tau = \{\phi, \{a\}, \{c\}, \{a, b\}, \{a, c\}, \{a, b, c\}, \{a, c, d\}, X\}$. Then:

(1) A subset $\{b, c\}$ of X is β^* - open but it is not β-open,

(2) A subset $\{b, d\}$ of X is e^* - open but it is not β^* - open,

Example 4.2

Let $X = \{a, b, c, d, e\}$ and $\tau = \{\phi, \{a, b\}, \{c, d\}, \{a, b, c, d\}, X\}$. Then $\{a, e\}$ is β^*- open but it is not Z^* - open.

Remark 3.3

By the following example we show that the intersection of any two β^*-open sets is not β^* - open.

Example 4.4

Let $X = \{a, b, c\}$ with topology $\tau = \{\phi, \{a\}, \{b\}, \{a, b\}, X\}$. Then $A = \{a, c\}$ and $B = \{b, c\}$ are β^*-open sets. But, $A \cap B = \{c\}$ is not β^*- open

.

Definition 2.2

Let (X, τ) be a topological space. Then:

(1) The union of all β^* - open sets of contained in A is called the β^*-

(2) interior of A and is denoted by β^*-*int* (A),

(3) The intersection of all β^* - closed sets of X containing A is called

(4) The β^* - closure of A and is denoted by β^* - *cl* (A).

Theorem 5.1

Let A, B be two subsets of a topological space (X, τ). Then the following are hold:

(1) β^*- $int(X) = X$ *and* β^* - $int(\phi) = \phi$,

(2) B^* - $int(A) \subseteq A$,

(3) *If* $A \subseteq B$, *then* β^*- $int(A) \subseteq \beta^*$ - $int(B)$,

(4) $x \in \beta^*$- $int(A)$ *if and only if there exist* β^* - *open* W *such that* $x \in W \subseteq A$,

(5) A *is* β^* - *open set if and only if* $A = \beta^*$ - $int(A)$,

(6) B^* - $int(\beta^*$ - $int(A)) = \beta^*$ - $int(A)$,

(7) B^* - $int(A \cap B) \subseteq \beta^*$ - $int(A) \cap \beta^*$ - $int(B)$,

(8) β^*- $int(A) \cup \beta^*$-$int(B) \subseteq \beta^*$-$int(A \cup B)$.

Example 4.5

Let $X = \{a, b, c, d\}$ with topology $\tau = \{\phi, \{a\}, \{b\}, \{a, b\}, X\}$.

(1) If $A = \{a, c\}$, $B = \{b, c\}$, then β^*- $cl(A) = A$, β^*- $cl(B) = B$ and β^*-)

(2) $lc\ (A \cup B) = X$. Thus β^*- $cl\ (A \cup B) \not\subset \beta^*$ - $cl(A) \cup \beta^*$ - $cl(B)$,

(3) If $A = \{a, c\}$, $C = \{a, b\}$, then β^* - $cl(C) = X$, β^* - $cl\ (A) = A$ and β^*- (A∩C) = $\{a\}$. Thus β^* - $cl(A) \cap \beta^*$- $cl(C) \not\subset \beta^*$- $cl(A \cap C)$,

(4) If $E = \{c, d\}$, $F = \{b, d\}$, then $E \cup F = \{b, c, d\}$ and hence β^*- $Int\ (E) = \phi$, β^* - $int\ (F) = F$ and β^* - $int\ (E \cup F) = \{b, c, d\}$. Thus β^* - $Int\ (E \cup F) \not\subset \beta^*$ - $int(E) \cup \beta^*$ - $int(F)$.

Theorem 5.2

For a subset A in a topological space (X, τ), the following statements are true:

(1) $B^{*}\text{-}cl(X/A) = X \setminus \beta^{*}\text{-}int(A)$,

(2) $\beta^{*}\text{-}int(X \setminus A) = X/\beta^{*}\text{-}cl(A)$.

Proof.

It follows from the fact the complement of β^{*}- open set is a β^{*} - closed

And $\cap_i(X/A_i) = X/\cup_i A_i$.

Theorem 5.3

Let A be a subset of a topological space (X, τ). Then the following are

Equivalent to :

(1) *A is a* β^{*} *- open set,*

(2) $A = \beta$ *- int(A)* $\cup$ *pint*δ *(A)*

.**Proof**

(1) ⇒ (2). Let *A* be a β^{*}- open set. Then $A \subseteq cl\ (int(cl\ (A))) \cup int\ (cl\delta(A))$ and hence by (Lemma 1.1) $A \subseteq (A \cap cl\ (int(cl(A)))) \cup (A \cap int\ (cl\delta(A))) = \beta\text{-}int\ (A) \cup pint\delta\ (A) \subseteq A$, (2) ⇒ (1).

Theorem 2.4

For a subset A of space (X, τ).Then the following are equivalent:

(1) *A is a* β ** - closed set,*

(2) $A = \beta$ *- cl(A)* $\cap$ *pcl* $\delta(A)$,

Proof

Theorem 5.5

β^{*} * - Continuous function

Example 4.6

Let X $\{a, b, c, d\}$ with topology $\tau = \{\phi, \{a\}, \{c\}, \{a, b\}, \{a, c\}, \{a, b, c\}, \{a, c, d\}, X\}$. Then, the function f: $(X, \tau) \to (X, \tau)$ defined by $f(a) = a$, $f(b) = f(c) = c$ and $f(d) = d$ is β^{*} - continuous but it is not β^{*} - continuous. The function f: $(X, \tau) \to (X, \tau)$ defined by $f(a) = d$, $f(b) = a$, $f(c) = c$ and $f(d) = b$ is e^{*}- continuous but it is not β^{*}- continuous .

Example 4.7

Let $X = \{a, b, c, d, e\}$ with topology $\tau = \{\phi, \{a, b\}, \{c, d\}, \{a, b, c, d\}, X\}$. Then function $f : (X, \tau) \to (X, \tau)$ which defined by $f(a) = a$, $f(b) = e$, $f(c) = c$, $f(d) = d$ and $f(e) = b$ is β^{*}- continuous but it is not Z^{*}- continuous.

Theorem 5.5

Let $f : (X, \tau) \to (Y, \sigma)$ *be a function. Then the following statements are*

Equivalent :

(1) *f is* β^{*} *- continuous,*

(2) For each $x \in X$ *and* $V \in \sigma$ *containing f(x), there exists* $U \in \beta^{*} O(X)$

(3) *containing x such that* $f(U) \subseteq V$,

The inverse image of each closed set in Y is β^{*} *- closed in X,*

(4) $int(cl(int(f^{-1}(B)))) \cap cl(int\ \delta(f^{-1}(B))) \subseteq f^{-1}(cl(B))$, *for each* $B \subseteq Y$,

(5) $f^{-1}(int(B)) \subseteq cl(int(cl(f^{-1}(B)))) \cup int(cl\delta(f^{-1}(B)))$, *for each* $B \subseteq Y$,

(6) β^*- $cl(f^{-1}(B)) \subseteq f^{-1}(cl(B))$, for each $B \subseteq Y$,
(7) $f(\beta^*$ - $cl(A)) \subseteq cl(f(A))$, for each $A \subseteq X$,
(8) $f^{-1}(int(B)) \subseteq \beta^*$ - $int(f^{-1}(B))$, for each $B \subseteq Y$.

Proof

(1) ⇔ (2) and (1) ⇔ (3) are obvious,
(3) ⇒ (4). Let $B \subseteq Y$. Then by (3) $f^{-1}(cl(B))$ is β^*-closed. This
means $f^{-1}(cl(B)) \supseteq int(cl(int(f^{-1}(cl(B))))) \cap cl(int\ \delta(f^{-1}(cl(B)))) \supseteq int(cl(int(f^{-1}(B)))) \cap cl(int\ \delta(f^{-1}(B)))$,
(4) ⇒ (5). By replacing Y/B instead of B in (4), we have
$Int(cl(int(f^{-1}(Y/B)))) \cap cl(int\ \delta(f^{-1}(Y\backslash B))) \subseteq f^{-1}(cl(Y/B))$, and
therefore $f^{-1}(int(B)) \subseteq cl(int(cl(f^{-1}(B)))) \cup int(cl\delta(f^{-1}(B)))$, for each $B \subseteq Y$,
(5) ⇒ (1). Obvious,
(3) ⇒ (6). Let $B \subseteq Y$ and $f^{-1}(cl(B))$ be β^* - closed in X. Then
B^*- $cl(f^{-1}(B)) \subseteq \beta^*$- $cl(f^{-1}(cl(B))) = f^{-1}(cl(B))$,
(7). Let $A \subseteq X$. Then $f(A) \subseteq Y$. By (6), we have $f^{-1}(cl(f(A))) \supseteq \beta^*$-
$Cl(f^{-1}(f(A))) \supseteq \beta^*$- $cl(A)$. Therefore, $cl(f(A)) \supseteq ff^{-1}(cl(f(A))) \supseteq f(\beta^*$ - $cl(A))$,
(7) ⇒ (3). Let $F \subseteq Y$ be a closed set. Then, $f^{-1}(F) = f^{-1}(cl(F))$. Hence by
(7), $f(\beta^*$- $cl(f^{-1}(F))) \subseteq cl(f(f^{-1}(F))) \subseteq (F) = F$, thus, β^*- $cl(f^{-1}(F)) \subseteq f^{-1}(F)$,
so, $f^{-1}(F) = \beta^*$-$cl(f^{-1}(F))$. Therefore, $f^{-1}(F) \in \beta^*C(X)$,
$Int(f^{-1}(int(B))) \subseteq \beta^*$- $int(f^{-1}(B))$. Therefore, $f^{-1}(int(B)) \subseteq \beta^*$- $int(f^{-1}(B))$,
(8) ⇒ (1). Let $U \subseteq Y$ be an open set. Then $f^{-1}(U) = f^{-1}(int(U)) \subseteq \beta^*$-$int(f^{-1}(U))$.
Hence, $f^{-1}(U)$ is β^* - open in X. Therefore, f is β^* - continuous.
Remarks 3.3
The composition of two β^* - continuous functions need not be β^* - continuous as show by the following example.

Example 4.7

Let $X = Y = Z = \{a, b, c, d\}$ with topologies $\tau = \{\phi, \{a\}, \{b\}, \{a, b\}, X\}$
. Let $f: X \to Y$ and $g: Y \to Z$ be functions defined
by $f(a) = b$, $f(b) = b$, $f(c) = c$, $f(d) = d$ and $g(a) = a$, $g(b) = c$,$g(c) = a$, $g(d) = d$,
respectively. Then f and g are β^*- continuous but $g \circ f$ is not β^*- continuous.

Theorem 5.6

If $f: (X, \tau) \to (Y, \sigma)$ is a β^* - continuous function and A is δ - open in X, then the restriction $f\backslash A: (A, \tau) \to (Y, \sigma)$ is β^*- continuous.

Proof.

Let V be an open set of Y. Then by hypothesis $f^{-1}(V)$ is β^* - open in X. we have $(f\backslash A)^{-1}(V) = f^{-1}(V) \cap A$
$\beta^* \in O(A)$. Thus, it follows that f/A is β^*- continuous.

Lemma 1.2

Let A and B be two subsets of a space (X, τ). If $A \in \delta O(X)$ and $B \in \beta^* O(A)$, the $A \cap B \in \beta^* O(X)$.

Theorem 5.7

Let $(X, \tau) \to (Y, \sigma)$ be a function and $\{Gi: i \in I\ s\}$ be a cover of X by δ-open sets of (X, τ). If $F/Gi : (Gi, \tau\ Gi) \to (Y, \sigma)$ is β^*- continuous for each $i \in I$, then f is β^*- continuous.

Proof.

Let V be an open set of (Y, σ). Then by $(V) = X \cap (V) = \cup \{Gi \cap f^{-1}(V): i \in I\} = \cup \{(f\backslash Gi)^{-1}(V): i \in I\}$. Since $f \setminus Gi$ is β^*-
Continuous for each $i \in I$, then $(f / Gi)^{-1}(V) \in YO(Gi)$ for each $i \in I$. we s $(f\backslash Gi)^{-1}(V)$ is β^* - continuous in X. Therefore, f is β^* - continuous in (X, τ).

Definition 2.3 The β^*- s frontier of a subset A of X, denoted by β^*- Fr (A), is defined by β^*- Fr (A) = β^*- cl $A) \cap \beta^*$- Cl (X/A) equivalently β^* - Fr (A) = β^*- cl (A) / β^*- int (A)

.

Theorem 5.8

The set of all points x of X at which a function f: $(X, \tau) \rightarrow (Y, \sigma)$ is not β^* - continuous is identical with the union of the β^* - frontiers of the inverse images of open sets containing f (x).

Proof. Necessity. Let x be a point of X at which f is not β^*- continuous. Then, there is an open set V of Y containing $f(x)$ such that $U \cap (X \backslash f^{-1}(V)) \neq \phi$, for every $U \in \beta^* O(X)$ containing x. Thus, we have $x \in \beta^*$- cl $(X/f^{-1}(V)) = X$ / β^*- int $(f^{-1}(V))$ and $x \in f^{-1}(V)$. Therefore, we have $x \in \beta^*$- Fr $(f^{-1}(V))$.Sufficiency. Suppose that $x \in \beta^*$- Fr $(f^{-1}(V))$, for some V is open set containing $f(x)$. Now, we assume that f is β^*- continuous at $x \in X$. Then there exists $U \in \beta^* O(X)$ containing x such that $f(U) \subseteq V$. Therefore, we have $x \in U \subseteq f^{-1}(V)$ and hence $x \in \beta^*$- int $(f^{-1}(V)) \subseteq X/ \beta^*$ - $Fr(f^{-1}(V))$. This is a contradiction. This means that f is not β^* - continuous at $x \varepsilon V$.

Theorem 5.9

If f: (X, τ) s$\rightarrow$ (Y, σ) is a β^*- continuous injection and (Y, σ) is Ti, then (X, τ) is β^*- Ti, where i = 0, 1, 2.

Proof.

We prove that the theorem for $i = 1$. Let Y be T_1 and x, y be distinct points in X. There exist open subsets U, V in Y such that $f(x) \in U$, $f(y) \notin U$, $f(x) \notin V$ and $f(y) \in V$. Since f is β^* - continuous, $f^{-1}(U)$ and $f^{-1}(V)$ are β^* - open subsets of X such that $x \in f^{-1}(U)$, $y \notin f^{-1}(U)$, $x \notin f^{-1}(V)$ and $y \in f^{-1}(V)$. Hence, X is β^* - T_1.

Theorem 5.10

If f: $(X, \tau) \rightarrow (Y, \sigma)$ is β^* - continuous, g:$(X, \tau) \rightarrow (Y, \sigma)$ is super-continuous and Y is Hausdorff, then the set $\{x \in X: f(x) = G(x)\}$ is β^* - closed in X. f is β^* - continuous in (X, τ).

Proof :

Let $A = \{x \in X : f(x) = g(x)\}$ and $x \notin A$. Then $f(x) \neq g(x)$. Since Y is Hausdorff, there exist open sets U and V of Y such That $f(x) \in U$, $g(x) \in V$ and $U \cap V = \phi$. Since f is β^* - continuous, there exists a β^* - open set G containing x such that $f(G) \subseteq U$. Since g is super - continuous, there exist an δ-open set H of X containing x such that $g(H) \subseteq V$. Now, put $W = G \cap H$, we have W is a β^* - open set containing x and $f(W) \cap g(W) \subseteq U \cap V = \phi$. Therefore, we obtain $W \cap A = \phi$ and hence x ε s β^* - cl (A). This shows that A is β^* - closed in X. Let $f : X \rightarrow Y$ be a function. The subse $\{(x, f(x)) : x \in X\}$ of the product space $X \times Y$ is called the graph of f and is denoted by $G(f)$.

Explain : Definition 2.4 A function $f : X \rightarrow Y$ has a (β^*, τ) - graph if for each $(x, y) \in (X \times Y) / G(f)$, there exist a β^*- open U of X containing x and an open set V of Y containing y such that $(U \times V) \cap G(f) = \phi$.

Proof: It follows readily from the above definition.

Theorem 5.11

If $f : X \rightarrow Y$ is a β^* - continuous function and Y is Hausdorff, then f has a (β^*, τ) s - graph.

Proof.

Let $(x, y) \in X \times Y$ such that $y \neq f(x)$. Then there exist open see that $y \in U$, $f(x) \in V$ and $V \cap U = \phi$. Since f is β^* - continuous, there exists β^* - open W containing x such that $f(W) \subseteq V$. This I $(W) \cap U \subseteq V \cap U = \phi$.

Definition 2.5

A space X is said to be β^*- compact if every β^* - open cover of X has a finite subcover.


Theorem 5.12
If f: (X, τ) → (Y, σ) has a (β, τ) - graph, then f (K) is closed in (Y, σ) for each subset K which is β * - compact relative to (X, τ).*

Proof.
Suppose that $y \in f(K)$.Then $(x, y) \in G(f)$ for each $x \in K$. Since $G(f)$ is (β^*, τ) - graph, there exist a β^*- open set U_x containing x and an open set V_x of Y containing y such that $f(U_x) \cap V_x = \phi$. The family $\{U_x : x \in K\}$ is a cover of K by β^*-open sets. Since K is β^*-compact relative to (X, τ), there exists a finite subset that $K \subseteq \cup \{U_x : x \in K_0\}$. Let $V = \cap \{V_x : x \in K_0\}$. have $f(K) \cap V \subseteq (\cup_{x \in K_0} f(U_x)) \cap V \subseteq \cup_{x \in K_0} (f(u_x)) \cap V = \phi$. It follows that, $y \in cl(f(K))$. Therefore, $f(K)$ is closed in (Y, σ).

Corollary 6.1
If f: (X, τ) → (Y, σ) is β-continuous function and Y is Hausdorff, then f(K) is closed in (Y, σ) for each subset K which is β*-compact relative to (X, τ).*

Theorem 5.13
*If f: X → Y is a β * - continuous function and Y is a Hausdorff space, then f has a (β *, τ) - graph.*

Proof
Let $(x, y) \in X \times Y$ such that $y \neq f(x)$ and Y be a Hausdorff space. Then there sexist two open sets U and V such that $y \in U$, $f(x) \in V$ and $V \cap U = \phi$. Since f is β^* - continuous, there exists a β^*-open set W containing x such that $f(W) \subseteq V$. This implies that $f(W) \cap U \subseteq V \cap U = \phi$. Therefore f has a (β^*, τ) - graph.



Corollary 6.2
If f: X → Y is β - continuous and Y is Hausdorff, then G(f) is β*- closed in X × Y.*

Theorem 5.14
If f: X → Y has a (β, τ)-graph and g: Y → Z is a β*- continuous function, then the set {(x, y): f(x) = g(y)} is β*- closed in X × Y.*

Proof:
Let $A = \{(x, y): f(x) = g(y)\}$ and (x, y) . We have $f(x) \neq g(y)$ and then $(x, g(y)) \in (X \times Z) \setminus G(f)$. Since f has a (β^*, τ) - graph, then there exist a β^* - open set U and an set V containing x and $g(y)$, respectively such that $f(U) \cap V = \phi$. Since g is a β^*- continuous function, then there exist an β^*- open set G containing y such that $g(G) \subseteq V$. We have $f(U) \cap g(G) = \phi$. This implies that $(U \times G) \cap A = \phi$. Since $U \times G$ is β^*- open, then (x, y) β^*- $cl(A)$. Therefore, A is β^*- closed in $X \times Y$.

Corollary 6.3
If f: X → Y, g: Y → Z are β-continuous functions and Z is Hausdorff, then the Set {(x, y): f(x) = g(y)} is β* - closed in X × Y.*

Theorem 5.15
If f: X → Y is a β- continuous function and Y is Hausdorff, then the set {(x, y) ∈ X × X: f(x) = f(y)} is β*- closed in X × X.*

Proof:

Let $A = \{(x, y): f(x) = f(y)\}$ and let $(x, y) \in (X \times X)\backslash A$. Then $f(x) \neq f(y)$. Since Y is Hausdorff, then there exist open sets U and V containing $f(x)$ and $f(y)$, respectively, such that $U \cap V = \phi$. But, f is β^*-continuous, then there exist β^* - open sets H and G in X containing x and y, respectively, such that $f(H) \subseteq U$ and $f(G) \subseteq V$. This implies $(H \times G) \cap A = \phi$ we have $H \times G$ is a β^* - open set in $X \times X$ containing (x, y). Hence, A is β^* - closed in $X \times X$.

Theorem 5.16

If $f: (X, \tau) \rightarrow (Y, \sigma)$ is β^*- continuous and S is closed in $X \times Y$, then $\upsilon x (S \cap G(f))$ is β^*- closed in X, where υxrepresents the projection of $X \times Y$ onto X.

Proof.


Let S be a closed subset of $X \times Y$ and $x \in \beta^*$-cl Let $U \in \tau$ containing x and $V \in \sigma$ containing $f(x)$. Since f is β^*-continuous, $x \in f^{-1}(V) \subseteq \beta^*$- $int(f^{-1}(V))$. Then $U \cap \beta^*$- $\text{int}(f^{-1}(V)) \cap \upsilon x (S \cap G(f))$ contains some point z of X. This implies that $(z, f(z)) S$ and $f(z) \in V$. Thus we have $(U \times V) \cap S \neq \phi$ and hence $(x, f(x)) cl(S)$. Since A is closed, then $(x, f(x)) S \cap G(f)$ and $x \in \upsilon x (S \cap G(f))$. Therefore $\upsilon x (S \cap G(f))$ is β^*-closed in (X, τ).

II. CONCLUSION

A topological space (X, τ) is said to be β^*-connected if it is not the union of two nonempty disjoint β^*- open sets. *If (X, τ) is a β^* - connected space and $f:(X, \tau) \rightarrow (Y, \sigma)$ has a (β^*, τ)-graph and β^*- continuous function, the constant.* Suppose that f is not constant. There exist disjoint points $x, y \in X$ such that $f(x) = f(y)$. Since $(x, f(x)) G(f)$, then $y \neq f(x)$, hence by, there exist open sets U and V containing x and $f(x)$ respectively such that $f(U) \cap V = \phi$. Since f is β^* - continuous, there exist a β^* - open sets G containing y such that $f(G) \subseteq V$. Since U and V are disjoint β^*- open sets of (X, τ), it follows that (X, τ) is not β^*- connected Therefore, f is constant. *Let $(X1, \tau1)$, $(X2, \tau2)$ and (X, τ) be topological spaces. Define a function $f: (X, \tau) \rightarrow (X1 \times X2, \tau1 \times \tau2)$ by $f(x) = (f(x1), f(x2))$. Then $fi: X \rightarrow (Xi, \tau i)$, where $(i = 1, 2)$ is β^*-continuous if f is β^*- continuous.*

REFERENCES

[1] Dontchev. J. and Noiri.T. Quasi-normal spaces and closed sets, Acta Math. Hungar. 89(3) (2000), 211– 219.
[2] O. Njastad On some classes of nearly open sets pacific .J. Math. 15 (1965), pp 47 – 53.
[3] A.S. Mashhour, M.E. Abd. EI – Monsef, S.N. EI – Deeb on pre – continuous and weak pre – Continuous mappings Proc. Math. Phys. Soc. Egypt, 53 (1982), pp 47 – 53.
[4] A.I.EL – Magharabi, A.M. Mubarki Z – open sets z – Continuitty in topological space. Int.Journal Math Arch. 2(10) (2011), PP. 1819 – 1827.
[5] E.Ekici On a open sets and decompositions of continuity and super continuity Annales Univ. Sci. Budapest, 51 (2008), p p 39 - 51.



10.22214/IJRASET

45.98

IMPACT FACTOR: 7.129

IMPACT FACTOR: 7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)