



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 13 Issue: V Month of publication: May 2025

DOI: https://doi.org/10.22214/ijraset.2025.71225

www.ijraset.com

Call: © 08813907089 E-mail ID: ijraset@gmail.com



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538

Volume 13 Issue V May 2025- Available at www.ijraset.com

### Genomic Innovation in Agriculture: The Rise of CRISPR-Cas9 in Plant Improvement

Tahir Metkari<sup>1</sup>, Sonakshi Bankar<sup>2</sup> Progressive Convent School And Junior College Kolki, Phaltan.

Abstract: CRISPR-Cas9 technology has transformed plant biotechnology by allowing precise genome editing, leading to remarkable enhancements in crop characteristics like yield, disease resistance, stress tolerance, and nutritional value. This article examines the molecular workings of CRISPR-Cas9, highlights its wide-ranging applications in various plant species, reviews significant case studies, and addresses current challenges and prospects for the future. Real-world examples are backed by recent peer-reviewed research.

Keywords: CRISPR-Cas9, Plant biotechnology, Genetics, Breeding, Agriculture.

### I. INTRODUCTION

Global food demand is rapidly increasing due to population growth, shifting dietary habits, and environmental stresses caused by climate change. Traditional plant breeding techniques are often slow and imprecise, typically requiring several generations to stabilize desired traits (Tester & Langridge, 2010). CRISPR-Cas9, originally an adaptive immune system in bacteria, has been adapted for precise genome editing, offering unmatched accuracy and efficiency (Jinek et al., 2012). This technology creates targeted DNA double-strand breaks (DSBs), which are then repaired by the cell's natural mechanisms, enabling precise knockout, insertion, or modification of genes linked to important agricultural traits (Doudna & Charpentier, 2014; Bortesi & Fischer, 2015).

### II. MECHANISM OF CRISPR-CAS9

CRISPR-Cas9 utilizes a single-guide RNA (sgRNA) to guide the Cas9 endonuclease to a specific complementary DNA sequence located next to a protospacer adjacent motif (PAM), usually the sequence 5'-NGG-3' (Jinek et al., 2012). Upon binding, Cas9 creates a double-strand break (DSB) at the target site, which the cell then repairs through one of two main pathways:

- Non-Homologous End Joining (NHEJ): An error-prone repair mechanism that often results in insertions or deletions (indels), potentially disrupting gene function (Puchta, 2005).
- Homology-Directed Repair (HDR): A high-fidelity repair process that uses a homologous DNA template to introduce precise nucleotide changes or insertions (Puchta, 2005).

Improvements in delivery techniques, including Agrobacterium-mediated transformation, biolistic methods, and the use of ribonucleoprotein complexes, have significantly increased CRISPR efficiency across various crop species (Bortesi & Fischer, 2015; Chen et al., 2019).

### III. APPLICATIONS IN PLANT BIOTECHNOLOGY

### A. Enhancing Crop Yield

CRISPR-Cas9 targeting of yield-associated genes has led to significant productivity gains. For instance, knocking out the GN1a gene in rice, which encodes cytokinin oxidase/dehydrogenase that reduces cytokinin levels, increases grain number and overall yield (Li et al., 2016). Likewise, deletion of CKX genes in wheat elevates cytokinin concentrations, resulting in enhanced grain yield and biomass accumulation (Rong et al., 2022; Zhang et al., 2021; Gasparis et al., 2019).

### B. Improving Nutritional Quality

Gene editing improves nutritional traits by manipulating metabolic pathways. In rice, editing SWEET transporter genes not only boosts disease resistance but also modifies sugar transport, increasing carbohydrate content (Oliva et al., 2019). In tomatoes, disruption of sugar metabolism regulators has raised sugar levels by up to 30% without compromising fruit size or yield (Wang, 2021). Additionally, genes involved in carotenoid biosynthesis have been edited in rice to enhance provitamin A levels, addressing vitamin A deficiency (Mishra et al., 2024).



### International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538

Volume 13 Issue V May 2025- Available at www.ijraset.com

### C. Developing Disease Resistance

Editing susceptibility (S) genes provides durable resistance against pathogens. Mutations in the MLO gene in wheat confer resistance to powdery mildew, a well-established strategy in cereals (Bui et al., 2023; Li et al., 2022). In rice, alterations in SWEET gene promoters prevent bacterial blight by blocking pathogen effector binding (Oliva et al., 2019). Furthermore, CRISPR-Cas9 has been applied to develop resistance to viruses and fungi by modifying host factors essential for infection (Andolfo et al., 2016).

### D. Enhancing Stress Tolerance

Abiotic stresses like drought and salinity limit crop yields worldwide. CRISPR-mediated edits of drought-responsive DREB genes in chickpea enhanced tolerance by activating stress response pathways (Singh et al., 2024). Similarly, editing the HyPRP1 gene in tomato improved tolerance to heat, salt, and drought stresses (Tran et al., 2023; Saikia et al., 2020). These precise genetic modifications enable crops to better endure harsh environmental conditions without sacrificing productivity.

### IV. CASE STUDIES

- 1) Soybean: Targeted editing of the GmEOD1 gene using CRISPR resulted in increased seed size and overall yield, achieving trait enhancement without adversely affecting plant growth (Tang et al., 2023).
- 2) Rice: Modification of the GN1a gene led to a higher number of grains per panicle, boosting yield in field conditions (Li et al., 2016).
- 3) Tomato: Alterations in sugar metabolism genes produced sweeter tomatoes while maintaining fruit size and yield (Wang, 2021).
- 4) Wheat: Knockout of MLO genes conferred resistance to powdery mildew, reducing the need for fungicide applications (Bui et al., 2023).
- 5) Chickpea: Editing stress-responsive genes enhanced drought tolerance, a critical trait for crops grown in semi-arid environments (Singh et al., 2024).

### V. CHALLENGES AND FUTURE PROSPECTS

### A. Off-target Effects and Specificity

Although CRISPR-Cas9 offers high precision, off-target mutations remain a significant concern (Zhang et al., 2015). To address this, approaches such as engineered high-fidelity Cas9 variants, paired nickases, and truncated sgRNAs have been developed to enhance specificity (Slaymaker et al., 2016; Kleinstiver et al., 2016). Cutting-edge research by Voytas and collaborators has introduced base editors and prime editors in plants, enabling precise nucleotide alterations without generating double-strand breaks, thereby minimizing off-target effects (Lin et al., 2020).

### B. Regulatory and Public Acceptance

Regulatory frameworks vary worldwide; some countries exempt CRISPR-edited crops that do not contain foreign DNA from GMO regulations (Wolt et al., 2016). However, public acceptance remains mixed, highlighting the need for transparent communication about the safety and benefits of CRISPR technology (Uddin et al., 2022).

### C. Technical Challenges

Delivering CRISPR components efficiently into difficult-to-transform plant species and regenerating edited plants pose ongoing challenges (Bortesi & Fischer, 2015). Innovations such as tissue culture-free gene editing, viral vector-mediated delivery, and nanoparticle-based transformation hold promise for overcoming these limitations (Duan et al., 2021).

### D. Future Directions

Future advancements include multiplexed genome editing to simultaneously modify multiple traits, epigenome editing to regulate gene expression without altering DNA sequences, and integrating CRISPR with synthetic biology for tailored trait development (Chen et al., 2019; Wolter & Puchta, 2018). Combining CRISPR with conventional breeding approaches will accelerate the development of crops suited for sustainable agriculture.



### International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 13 Issue V May 2025- Available at www.ijraset.com

### REFERENCES

- [1] Andolfo, G., Iovieno, P., Frusciante, L., & Ercolano, M. R. (2016). Genome-editing technologies for enhancing plant disease resistance. Frontiers in Plant Science, 7, 1813
- [2] Bortesi, L., & Fischer, R. (2015). The CRISPR/Cas9 system for plant genome editing and beyond. Biotechnology advances, 33(1), 41-52.
- [3] Bui, T. P., Le, H., Ta, D. T., Nguyen, C. X., Le, N. T., Tran, T. T., & Do, P. T. (2023). Enhancing powdery mildew resistance in soybean by targeted mutation of MLO genes using the CRISPR/Cas9 system. BMC Plant Biology, 23(1), 533.
- [4] Chen, K., Wang, Y., Zhang, R., Zhang, H., & Gao, C. (2019). CRISPR/Cas genome editing and precision plant breeding in agriculture. Annual review of plant biology, 70(1), 667-697.
- [5] Doudna, J. A., & Charpentier, E. (2014). The new frontier of genome engineering with CRISPR-Cas9. Science, 346(6213), 1258096.
- [6] Duan, L., Ouyang, K., Xu, X., Xu, L., Wen, C., Zhou, X., & Liang, Y. (2021). Nanoparticle delivery of CRISPR/Cas9 for genome editing. Frontiers in Genetics, 12, 673286.
- [7] Endo, M., et al. (2019). Carotenoid engineering in rice for biofortification. Plant Physiology, 179(1), 33-46.
- [8] Gasparis, S., et al. (2019). CRISPR/Cas9-targeted mutagenesis of wheat CKX genes increases grain yield. Plant Biotechnology Journal, 17(3), 557-566.
- [9] Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J. A., & Charpentier, E. (2012). A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. science, 337(6096), 816-821.
- [10] Kleinstiver, B. P., Pattanayak, V., Prew, M. S., Tsai, S. Q., Nguyen, N. T., Zheng, Z., & Joung, J. K. (2016). High-fidelity CRISPR-Cas9 nucleases with no detectable genome-wide off-target effects. Nature, 529(7587), 490-495.
- [11] Li, M., Li, X., Zhou, Z., Wu, P., Fang, M., Pan, X., & Li, H. (2016). Reassessment of the four yield-related genes Gn1a, DEP1, GS3, and IPA1 in rice using a CRISPR/Cas9 system. Frontiers in plant science, 7, 377.
- [12] Li, S., Lin, D., Zhang, Y., Deng, M., Chen, Y., Lv, B., & Gao, C. (2022). Genome-edited powdery mildew resistance in wheat without growth penalties. Nature, 602(7897), 455-460.
- [13] Lin, Q., et al. (2020). Prime genome editing in rice and wheat. Nature Biotechnology, 38(5), 582-585.
- [14] Mishra, M., Afzal, S., & Singh, N. K. (2024). CRISPR/Cas9-Mediated Genome Engineering for Production of Biofortified Crops. In Phytoremediation and Biofortification (pp. 303-333). Apple Academic Press.
- [15] Oliva, R., Ji, C., Atienza-Grande, G., Huguet-Tapia, J. C., Perez-Quintero, A., Li, T., & Yang, B. (2019). Broad-spectrum resistance to bacterial blight in rice using genome editing. Nature biotechnology, 37(11), 1344-1350.
- [16] Puchta, H. (2005). The repair of double-strand breaks in plants: mechanisms and consequences for genome evolution. Journal of experimental botany, 56(409),
- [17] Rong, C., Liu, Y., Chang, Z., Liu, Z., Ding, Y., & Ding, C. (2022). Cytokinin oxidase/dehydrogenase family genes exhibit functional divergence and overlap in rice growth and development, especially in control of tillering. Journal of experimental botany, 73(11), 3552-3568.
- [18] Saikia, B., Singh, S., Debbarma, J., Velmurugan, N., Dekaboruah, H., Arunkumar, K. P., & Chikkaputtaiah, C. (2020). Multigene CRISPR/Cas9 genome editing of hybrid proline rich proteins (HyPRPs) for sustainable multi-stress tolerance in crops: the review of a promising approach. Physiology and Molecular Biology of Plants, 26(5), 857-869.
- [19] Singh, S., Mishra, V., Riyazuddin, R., Chugh, V., & Upadhyay, S. K. (2024). CRISPR/Cas9-Mediated Genome Editing for Trait Improvement and Stress Tolerance in Leguminosae (Legume Family). Plant Breeding.
- [20] Slaymaker, I. M., Gao, L., Zetsche, B., Scott, D. A., Yan, W. X., & Zhang, F. (2016). Rationally engineered Cas9 nucleases with improved specificity. Science, 351(6268), 84-88.
- [21] Tang, Q., Wang, X., Jin, X., Peng, J., Zhang, H., & Wang, Y. (2023). CRISPR/Cas technology revolutionizes crop breeding. Plants, 12(17), 3119.
- [22] Tester, M., & Langridge, P. (2010). Breeding technologies to increase crop production in a changing world. Science, 327(5967), 818-822.
- [23] Tran, M. T., Son, G. H., Song, Y. J., Nguyen, N. T., Park, S., Thach, T. V., & Kim, J. Y. (2023). CRISPR-Cas9-based precise engineering of SIHyPRP1 protein towards multi-stress tolerance in tomato. Frontiers in Plant Science, 14, 1186932.
- [24] Uddin, A., Gallardo, R. K., Rickard, B., Alston, J., & Sambucci, O. (2022). Consumer acceptance of new plant-breeding technologies: An application to the use of gene editing in fresh table grapes. Plos one, 17(12), e0270792.
- [25] Wang, B., Li, N., Huang, S., Hu, J., Wang, Q., Tang, Y., & Yu, Q. (2021). Enhanced soluble sugar content in tomato fruit using CRISPR/Cas9-mediated SIINVINH1 and SIVPE5 gene editing. PeerJ, 9, e12478.
- [26] Wolt, J. D., Wang, K., & Yang, B. (2016). The regulatory status of genome-edited crops. Plant biotechnology journal, 14(2), 510-518.
- [27] Wolter, F., & Puchta, H. (2018). Application of CRISPR/Cas to understand cis-and trans-regulatory elements in plants. Plant transcription factors: methods and protocols. 23-40.
- [28] Xu, Z., et al. (2019). Editing SWEET gene promoters for resistance to bacterial blight in rice. Nature Communications, 10, 4925.
- [29] Zhang, X. H., Tee, L. Y., Wang, X. G., Huang, Q. S., & Yang, S. H. (2015). Off-target effects in CRISPR/Cas9-mediated genome engineering. Molecular therapy Nucleic acids, 4.
- [30] Zhang, S., Zhang, R., Gao, J., Song, G., Li, J., Li, W., & Li, G. (2021). CRISPR/Cas9-mediated genome editing for wheat grain quality improvement. Plant Biotechnology Journal, 19(9), 1684.









45.98



IMPACT FACTOR: 7.129



IMPACT FACTOR: 7.429



## INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call: 08813907089 🕓 (24\*7 Support on Whatsapp)