



# IJRASET

International Journal For Research in  
Applied Science and Engineering Technology



---

# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume:** 14    **Issue:** III    **Month of publication:** March 2026

**DOI:** <https://doi.org/10.22214/ijraset.2026.78985>

[www.ijraset.com](http://www.ijraset.com)

Call:  08813907089

E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)

# Interplanetary Magnetic Field-Magnetosphere Coupling: A Review

Shriram Lahauriya<sup>1</sup>, A. C. Pandey<sup>2</sup>, Kalpana Singh<sup>3</sup>

<sup>1</sup>Department of Physics, Government P.G. Autonomous College, Datia (M.P.) India- 475661

<sup>2</sup>Department of Physics, Govt. New Science College, Rewa (MP)-486001

<sup>3</sup>Institute of Technology and Management, NH 24, Bakshi Ka Talab, Lucknow, U.P. -226008, India

**Abstract:** *The interaction between the solar wind and Earth's magnetosphere is primarily governed by the Interplanetary Magnetic Field (IMF). The IMF, carried by the solar wind plasma from the Sun, plays a critical role in transferring energy, momentum, and plasma into the magnetosphere. The orientation and magnitude of IMF components, particularly the  $B_z$  component, control the efficiency of solar wind-magnetosphere coupling through magnetic reconnection processes. This review summarizes the physical mechanisms governing IMF-magnetosphere coupling, including reconnection, solar wind pressure effects, magnetospheric convection, and geomagnetic storms. It also reviews observational studies, numerical modeling approaches, and recent advances in understanding energy transfer processes. The paper concludes with future directions for heliophysics research and space weather prediction.*

## I. INTRODUCTION

The space environment surrounding Earth is strongly influenced by the continuous flow of plasma from the Sun known as the solar wind. Embedded within this plasma is the Interplanetary Magnetic Field (IMF), which originates from the solar magnetic field and is transported outward through the heliosphere[1-3].

The interaction between the IMF and Earth's magnetosphere forms a fundamental component of space physics and plays a major role in space weather phenomena.

Solar wind-magnetosphere coupling determines the dynamics of the magnetosphere, the occurrence of geomagnetic storms, and the generation of auroral phenomena[4-7].

Early theoretical work demonstrated that the orientation of the IMF controls the transfer of energy into the magnetosphere. When the IMF has a southward component relative to Earth's magnetic field, magnetic reconnection occurs at the magnetopause, allowing solar wind energy and plasma to enter the magnetosphere[8-9]. Understanding this coupling is crucial for predicting space weather effects that impact satellites, communication systems, and power grids.

## II. INTERPLANETARY MAGNETIC FIELD

The IMF originates from the solar magnetic field and is carried outward by the solar wind plasma due to the high electrical conductivity of the plasma, which effectively "freezes" magnetic field lines into the solar wind flow[10].

### A. Parker Spiral Structure

Because the Sun rotates while emitting solar wind, the IMF forms a spiral structure known as the Parker spiral. This spiral geometry determines the orientation of the magnetic field at different distances from the Sun[11]. At the distance of Earth (1 AU), the IMF strength typically ranges from **1–10 nT**, although stronger values may occur during solar eruptions.

### B. IMF Components

The IMF vector is commonly expressed in three components:

- $B_x$  – radial component
- $B_y$  – azimuthal component
- $B_z$  – north-south component

Among these, the  $B_z$  component is the most important for magnetospheric coupling.

### III. STRUCTURE OF EARTH'S MAGNETOSPHERE

The magnetosphere is the region surrounding Earth where the planet's magnetic field dominates the motion of charged particles[12].

Major regions include:

- 1) Bow shock
- 2) Magnetosheath
- 3) Magnetopause
- 4) Magnetotail
- 5) Radiation belts

The solar wind compresses the magnetosphere on the dayside and stretches it into a long magnetotail on the nightside.

Solar wind dynamic pressure determines the size and shape of the magnetosphere, while the interplanetary electric field controls energy transfer processes[13-14].

### IV. MECHANISMS OF IMF-MAGNETOSPHERE COUPLING

#### A. Magnetic Reconnection

Magnetic reconnection is the primary mechanism for transferring energy from the solar wind into the magnetosphere.

When the IMF is oriented opposite to Earth's magnetic field at the magnetopause, magnetic field lines merge and reconnect. Newly connected field lines allow solar wind plasma to enter the magnetosphere[15].

Reconnection leads to:

- plasma transport
- particle acceleration
- magnetospheric convection

This process is most efficient when the IMF  $B_z$  component is negative (southward).

#### B. Magnetospheric Convection

Magnetic reconnection drives large-scale plasma circulation known as magnetospheric convection.

The convection cycle consists of:

- Dayside reconnection
- Transport of open magnetic flux to the magnetotail
- Nightside reconnection
- Return flow of plasma toward the dayside

This cycle transfers energy from the solar wind into the magnetosphere and ionosphere.

#### C. Energy Transfer and Coupling Functions

Several coupling functions have been proposed to quantify solar wind-magnetosphere interaction.

One widely used parameter is Akasofu's  $\epsilon$  parameter, which estimates the rate of solar wind energy input into the magnetosphere.

Studies indicate that the  $B_z$  component has the strongest correlation with magnetospheric activity and ring current injection[16-17].

### V. IMF ORIENTATION EFFECTS

#### A. Southward IMF

Southward IMF enhances magnetic reconnection and leads to strong coupling between the solar wind and magnetosphere.

Consequences include:

- geomagnetic storms
- enhanced auroral activity
- increased radiation belt fluxes

Sustained southward IMF can drive strong geomagnetic storms and continuous energy transfer into the magnetosphere.

#### B. Northward IMF

When the IMF is northward, reconnection occurs in the magnetotail lobes rather than on the dayside magnetopause.

This produces:

- weaker geomagnetic activity
- complex ionospheric convection patterns
- reduced energy transfer

Under these conditions, the ionospheric convection pattern often shows four-cell structures instead of the typical two-cell configuration.

## VI. ROLE OF IMF BY COMPONENT

Although  $B_z$  dominates energy transfer, the  $B_y$  component significantly influences magnetospheric asymmetries. Research shows that IMF  $B_y$  induces asymmetric currents and plasma convection patterns in the magnetosphere and ionosphere. These asymmetries affect auroral activity in both hemispheres[18].

## VII. MAGNETOSPHERIC RESPONSE TO IMF VARIATIONS

The magnetosphere responds rapidly to changes in IMF orientation. Studies show that geomagnetic activity often follows IMF changes with a time delay of about **20–30 minutes**, corresponding to the time required for solar wind disturbances to propagate through the magnetosphere. Rapid transitions from northward to southward IMF can trigger magnetospheric substorms and enhanced plasma convection[19].

## VIII. GEOMAGNETIC STORMS

Geomagnetic storms occur when large amounts of energy from the solar wind are transferred into the magnetosphere[20].

Major drivers include:

- Coronal Mass Ejections (CMEs)
- Interplanetary shocks
- Magnetic clouds

These events enhance the ring current and produce large disturbances in Earth's magnetic field.

## IX. MAGNETOSPHERE–IONOSPHERE COUPLING

The magnetosphere and ionosphere are strongly coupled through field-aligned currents and plasma convection. Energy transferred from the solar wind flows through the magnetosphere and is deposited in the ionosphere, producing:

- 1) auroras
- 2) ionospheric heating
- 3) enhanced currents

The ionospheric response depends strongly on IMF orientation and solar wind parameters.

## X. NUMERICAL MODELING OF IMF–MAGNETOSPHERE INTERACTION

Modern space physics relies heavily on numerical simulations.

Common models include:

- 1) Magnetohydrodynamic (MHD) simulations
- 2) Global magnetosphere models
- 3) Space Weather Modeling Framework (SWMF)

Simulations have shown that sustained southward IMF produces strong convection and directly driven magnetospheric states.

## XI. SPACE WEATHER IMPLICATIONS

IMF–magnetosphere coupling is central to space weather forecasting.

Space weather events can affect:

- 1) satellites
- 2) GPS signals
- 3) communication systems
- 4) power grids
- 5) astronaut safety

Real-time monitoring of IMF conditions using spacecraft such as ACE and DSCOVR provides early warning of geomagnetic storms. Space Weather Implications refer to the effects of solar activity on Earth's environment, technology, and human systems. Space weather originates mainly from the Sun and includes phenomena like solar flares, coronal mass ejections (CMEs), and high-speed solar wind streams [21-22].

#### A. Key Space Weather Phenomena

- 1) Solar Flares – Sudden bursts of radiation from the Sun
- 2) Coronal Mass Ejections (CMEs) – Massive clouds of charged particles ejected into space
- 3) Solar Wind – Continuous flow of charged particles
- 4) Geomagnetic Storms – Disturbances in Earth's magnetic field [23]

#### B. Major Implications

##### 1. Satellite and Space Systems

- Damage to satellites used by NASA and communication companies
- Disruption of GPS, navigation, and weather forecasting systems
- Increased atmospheric drag causing satellites to lose orbit

##### 2. Communication Systems

- High-frequency (HF) radio communication blackout
- Disruption in aviation communication, especially on polar routes
- Interference in mobile and internet signals

##### 3. Power Grids

- Geomagnetically induced currents (GICs) can overload transformers
- Large-scale blackouts (e.g., similar to effects seen during the Quebec Blackout of 1989)

##### 4. Human Health

- Increased radiation exposure for astronauts aboard the International Space Station
- Potential risks to airline crew and passengers at high altitudes

##### 5. Navigation Systems

- Errors in GPS positioning affecting aviation, military, and civilian navigation
- Impact on precision farming and autonomous systems

##### 6. Auroras

- Beautiful natural displays like Northern Lights (Aurora Borealis)
- Caused by interaction of solar particles with Earth's magnetosphere

#### C. Scientific & Technological Importance

- 1) Helps in understanding the Sun–Earth connection
- 2) Important for designing radiation-hardened electronics (relevant to your interest in advanced electronic devices)
- 3) Critical for space missions planning and safety

## XII. FUTURE RESEARCH DIRECTIONS

Future research will focus on:

- 1) high-resolution measurements of the solar wind
- 2) improved coupling models
- 3) multi-satellite observations
- 4) artificial intelligence in space weather prediction

Upcoming missions will provide improved understanding of solar wind dynamics and magnetospheric processes.

## XIII. CONCLUSION

The interaction between the Interplanetary Magnetic Field and Earth's magnetosphere governs the transfer of energy from the Sun into the near-Earth environment. Magnetic reconnection, driven primarily by the orientation of the IMF Bz component, controls

magnetospheric dynamics, geomagnetic storms, and auroral activity [24-25]. Continued observations and modeling efforts are essential for improving our understanding of solar wind–magnetosphere coupling and its effects on space weather [26].

## REFERENCES

- [1] R. Miteva and S. W. Samwel, "Catalog of Geomagnetic Storms with Dst Index  $\leq -50$  nT and Their Solar and Interplanetary Origin (1996–2019)," *Atmosphere*, 14, 12, 1744, 2023, doi: 10.3390/atmos14121744.
- [2] Singh Sham. Space weather disturbances and their geoeffective-ness during solar cycle 23 and 24. *Iranian Journal of Physics Research*, 24, 3, 117-133 (2024). <https://doi.org/10.47176/ijpr.24.3.31863>.
- [3] Sham Singh, Ajay Vasishth, Bikramjit Singh, A. C. Panday, A. P. Mishra. The Plasma Approximation. *IJSRST*. 3 (1), 108-117 (2017). <https://doi.org/10.32628/IJSRST173125>.
- [4] Kumar, P., Pal, M., & Singh, S. Analysis of intense geomagnetic storm on 24 April 2023 with interplanetary parameters. *J. Atmos. Terr. Phys.* 269,106481 (2025). <https://doi.org/10.1016/j.jastp.2025.106481>
- [5] Kumar, P., Pal, M., & Singh, S. Analysis 2023 Storms based on different time scales (Dst, Kp & Sym/H). *The Journal of Space Safety Engineering*. (2024). <https://doi.org/10.1016/j.jsse.2024.12.002>
- [6] Kumar, P., Pal, M., & Singh, S. Analysis of solar plasma parameters during intense geomagnetic storm (SYM/H < -100). *Radiation Effects and Defects in Solids*. 179(7-8), 985–993 (2024). DOI: 10.1080/10420150.2024.2378423
- [7] W. Rukundo, "The Ionospheric Dynamics of the African Sector Responding to a Severe Geomagnetic Storm; the Storm of 3–5 November 2021," *Space Weather*. 21, 3 (2023). doi: 10.1029/2022sw003219.
- [8] Singh, Sham, and A. P. Mishra "Cosmic ray intensity increases during high solar activity period for the solar cycles 22 and 23." *Indian J Phys* 93, 139–145 (2019). <https://doi.org/10.1007/s12648-018-1284-3>
- [9] L. Lefèvre et al., "Detailed Analysis of Solar Data Related to Historical Extreme Geomagnetic Storms: 1868 – 2010," *Solar Physics*. 291, 5, 1483 (2016). doi: 10.1007/s11207-016-0892-3.
- [10] X. An, X. Meng, H. Chen, W. Jiang, R. Xi, and Q. Chen, "Modeling Global Ionosphere Based on Multi-Frequency, Multi-Constellation GNSS Observations and IRI Model," *Remote Sensing*. 12, 3, 439 (2020) doi: 10.3390/rs12030439.
- [11] I. G. Richardson and H. V. Cane, "Solar wind drivers of geomagnetic storms over more than four solar cycles," *AIP conference proceedings*. (2013). doi: 10.1063/1.4811075.
- [12] A. A. Abunin, M. A. Abunina, A. V. Belov, and I. M. Chertok, "Peculiar Solar Sources and Geospace Disturbances on 20–26 August 2018," *Solar Physics*. 295, 1, (2020) doi: 10.1007/s11207-019-1574-8.
- [13] Sham Singh., & A. P. Mishra. "Interaction of solar plasma near-Earth with reference to Geomagnetic storms during maxima of solar cycle 24. *Indian J Phys* 89, 1227–1234 (2015). <https://doi.org/10.1007/s12648-015-0703-y>
- [14] X. Luo, S. Gu, Y. Lou, C. Xiong, B. Chen, and X. Jin, "Assessing the Performance of GPS Precise Point Positioning Under Different Geomagnetic Storm Conditions during Solar Cycle 24," *Sensors*. 18, 6, 1784 (2018). doi: 10.3390/s18061784.
- [15] D. Telloni et al., "Study of the Influence of the Solar Wind Energy on the Geomagnetic Activity for Space Weather Science." (2020).
- [16] Sham Singh, D. Shrivastava, and A.P. Mishra, "effect of solar and inter-planetary disturbances on space weather." *Indian journal of scientific research*. 3, 121-125, (2012).
- [17] P. Kumar, M. Pal, and S. Singh, "Analysis of Intense Geomagnetic Storm on 24 April 2023 with Interplanetary Parameters," *Journal of Atmospheric and Solar-Terrestrial Physics*, p. 106481, Feb. 2025, doi: 10.1016/j.jastp.2025.106481.
- [18] Sham Singh, M. Pal, P. Kumar, A. Rani, N. Thakur, K. Singh and A. Mishra. (2025). The Relationship between Cosmic Ray Intensity, Sunspot Cycle with Geomagnetic Activity. *Proceedings of Science*. 1312. <https://doi.org/10.22323/1.444.1312>.
- [19] Y. Nakagawa, S. Nozawa, and A. Shinbori, "Relationship between the low-latitude coronal hole area, solar wind velocity, and geomagnetic activity during solar cycles 23 and 24," *Earth, planets and space*, vol. 71, no. 1, Feb. 2019, doi: 10.1186/s40623-019-1005-y.
- [20] I. G. Richardson, "Geomagnetic activity during the rising phase of solar cycle 24," *Journal of space weather and space climate*. 3, (2013) doi: 10.1051/swsc/2013031.
- [21] S. Watari, A. Nakamizo, and Y. Ebihara, "Solar events and solar wind conditions associated with intense geomagnetic storms," *Earth Planets and Space*. 75 1 (2023) doi: 10.1186/s40623-023-01843-2.
- [22] Sham Singh (2024). Cosmic-Ray Modulation in Relation to Solar and Heliospheric Parameters. *Proceedings of Science*. 1310. <https://doi.org/10.22323/1.444.1310>.
- [23] Sham Singh, A. C. Panday, Kalpana Singh & A. P. Mishra. (2017). Effect of geomagnetic storms and their association with solar wind velocity and IMF during solar cycle 23 and 24. *International Journal of Pure and Applied Physics*. 13(1) 3 (2017).
- [24] N. Gopalswamy, P. Mäkelä, S. Yashiro, S. Akiyama, and H. Xie, "Solar activity and space weather." Feb. 01, 2022. Accessed: Jul. 21, 2024. <https://arxiv.org/pdf/2201.02724v1.pdf>
- [25] P. Kumar, M. Pal, and S. Singh, "Correlations the Interplanetary Characteristics and the Occurrence of Geomagnetic Large Storms." *Scope*. 13, 1 (2024). doi: 10.54882/14202414202317855
- [26] Sham Singh, Kalpana Singh, Ajay Vasishth, A. C. Panday, Shabir Ahmad Shabir & A. P. Mishra. *IJSRSET*, Effect of Geomagnetic Storms and Their Association with Solar Wind Velocity during 1996-2016. 3(5), 456 (2017). <https://doi.org/10.32628/IJSRSET173498>.



10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)