



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 13 **Issue:** III **Month of publication:** March 2025

DOI: <https://doi.org/10.22214/ijraset.2025.67958>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

Non-Linear Programming Problem using Pentagonal Fuzzy Numbers

K . Vaishali

PG and Research Department of Mathematics, Seethalakshmi Ramaswami College (Autonomous), Tiruchirappalli, (Affiliated to Bharathidasan University, Tiruchirappalli)

Abstract: In this study, pentagonal fuzzy numbers are used as the decision parameters. To solve these fuzzy nonlinear programming problems, we first convert all of the pentagonal fuzzy numbers into crisp values using robust ranking method. Next, we obtain a crisp nonlinear programming problem. Finally, we apply the Kuhn-Tucker conditions to obtain the best solution.

Keywords: Fuzzy Non-Linear programming problem, Robust Ranking Method, Pentagonal Fuzzy Number.

I. INTRODUCTION

Traditional optimization techniques have been used successfully for many years. The decision maker's decision is based on the information available from the mathematical modelling environment. However, we have a lot of trouble formulating and addressing real-world problems because of the ambiguities and inexactness of the decision parameters. These inaccuracies are caused by subjectivity and human opinion, lack of information, estimation error, and imperfect representation of information, among other things. As a result, conventional mathematical techniques are unable to formulate and successfully solve real-world problems. Thus, fuzzy optimization techniques offer a useful and powerful tool for optimization under fuzzy environment.

In 1965, Zadeh introduced Fuzzy set [19]. It plays a crucial role in solving the real-world problems. Thereafter, Bellman and Zadeh have applied the decision-making concept in fuzzy nature [3]. In 1991, Fuzzy set theory and its applications was discussed by Zimmermann [20]. Arun Pratap Singh [2] presented Allocation of subjects in an educational institution by Robust ranking method [8]. Monalisha Pattnaik [11] applied Robust ranking for two phase fuzzy Optimization. S. Yahya Mohamed [18] Solving Fuzzy Travelling Salesman Problem Using Octagon Fuzzy Numbers with alpha-Cut. P. Umamaheswari et al [15] presented fuzzy problem converted into parametric form. Then applying by Kuhn Tucker conditions, the optimal solution of the problem is obtained.

Darunee Hunwisai et al [7], M. S. Annie Christi [1] applied Robust ranking for fuzzy transportation problems [14]. Sudha et al [13] suggested a new approach for fuzzy neutrosophic quadratic problem. Kokila et al [9] used Robust ranking method for solving Fuzzy Octagonal number. S. U. Malini et al [10] applied Modi method Fuzzy Octagonal number. Thangaraj Beaula and Vijaya [4-6] have adopted different fuzzy numbers and ranking methods to solve critical path problems. Vijaya et al [16,17] have used Pythagorean Fuzzy numbers and Neutrosophic Fuzzy numbers to find the solution of Decision making problem and critical path problems. Vijaya et al have used complex Fermatean Neutrosophic fuzzy sets to solve decision making problem.

Vijaya, V., and D. Rajalaxmi. "Decision making in fuzzy environment using Pythagorean fuzzy numbers." Mathematical Statistician and Engineering Applications 71.4 (2022): 846-854. Vijaya, V., D. Rajalaxmi, and H. Manikandan. "Finding critical path in a fuzzy project network using neutrosophic fuzzy number." Advances and Applications in Mathematical Sciences 21.10 (2022): 5743-5753. Vijaya V, Said Broumi, & Manikandan H. (2025). Complex Fermatean Neutrosophic Sets and their Applications in Decision Making. Neutrosophic Sets and Systems, 81, 827-839.

In this paper, we solve fuzzy nonlinear programming problems in which all of the problem's decision parameters are pentagonal fuzzy numbers. First, we use the robust ranking approach to turn all of the pentagonal fuzzy numbers into crisp values, and then we get a crisp nonlinear programming problem, which we solve using the Kuhn Tucker conditions.

II. PRELIMINARIES

Definition: 2.1

If X is a universal set and $x \in X$, then a fuzzy set \tilde{A} defined as a collection of ordered pairs, $\tilde{A} = \{(x, \mu_{\tilde{A}}(x)), x \in X\}$ where $\mu_{\tilde{A}}(x)$ is called the membership function that maps X to the membership space M .

Definition:2.2

A fuzzy number $A = (a_1, a_2, a_3, a_4, a_5)$ is called a pentagonal fuzzy number. If its membership function is given by

$$\mu_{\tilde{A}}(x) = \begin{cases} 0 & \text{for } x < a_1, a_2 \leq x \\ \frac{x-a_1}{a_2-a_1} & \text{for } a_1 \leq x \leq a_2 \\ \frac{x-a_2}{a_3-a_2} & \text{for } a_2 \leq x \leq a_3 \\ 1 & \text{for } x = a_3 \\ \frac{a_4-x}{a_4-a_3} & \text{for } a_3 \leq x \leq a_4 \\ \frac{a_5-x}{a_5-a_4} & \text{for } a_4 \leq x \leq a_5 \\ 0 & \text{for } x > a_5 \end{cases}$$

Here for PFN $A = (a_1, a_2, a_3, a_4, a_5)$, a_3 is the middle point and (a_1, a_2) and (a_4, a_5) are the left and right side points of a_3 respectively. The middle point a_3 has the grade of membership 1 and w_1, w_2 are the grades of points a_2, a_4 .

Note that, every PFN is associated with two weights w_1 and w_2 .

Definition:2.3

A fuzzy number $A = (a_1, a_2, a_3, a_4, a_5)$ is a pentagonal fuzzy number (PFN) having membership function

$$\mu_{(x, w_1, w_2)} = \begin{cases} w_1 \left(\frac{x-a_1}{a_2-a_1} \right) & \text{for } a_1 \leq x \leq a_2 \\ 1 - (1 - w_1) \left(\frac{x-a_2}{a_3-a_2} \right) & \text{for } a_2 \leq x \leq a_3 \\ 1 & \text{for } x = a_3 \\ 1 - (1 - w_2) \left(\frac{x-a_3}{a_4-a_3} \right) & \text{for } a_3 \leq x \leq a_4 \\ w_2 \left(\frac{x-a_5}{a_4-a_5} \right) & \text{for } a_4 < x < a_5 \\ 0 & \text{for } x > a_5 \end{cases}$$

Case(i):

When $w_1 = w_2 = 0$, then the pentagonal fuzzy number is reduced to a triangular fuzzy number. That is, $A = (a_1, a_2, a_3, a_4, a_5) \approx (a_2, a_3, a_4)$ where membership function is given by

$$\mu_{\tilde{A}}(x) = \begin{cases} 0 & \text{for } x \leq a_2 \\ 1 - \left(\frac{x-a_2}{a_3-a_2} \right) & \text{for } a_2 \leq x \leq a_3 \\ 1 & \text{for } x = a_3 \\ 1 - \left(\frac{x-a_3}{a_4-a_3} \right) & \text{for } a_3 \leq x \leq a_4 \\ 0 & \text{for } x \geq a_4 \end{cases}$$

Case (ii):

When $w_1 = w_2 = 1$, then the pentagonal fuzzy number becomes a trapezoidal fuzzy number. That is, the membership function is given by

$$\mu_{\tilde{A}}(x) = \begin{cases} 0 & \text{for } x \leq a_1 \\ \left(\frac{x-a_1}{a_2-a_1} \right) & \text{for } a_1 \leq x \leq a_2 \\ 1 & \text{for } a_2 \leq x \leq a_5 \\ \left(\frac{a_4-x}{a_5-a_4} \right) & \text{for } a_4 \leq x \leq a_5 \\ 0 & \text{for } x > a_5 \end{cases}$$

Definition:2.4

A fuzzy set \tilde{A} defined on X is called a normal fuzzy set if there exist at least one $x \in X$ such that $\mu_{\tilde{A}}(x) = 1$.

Definition:2.5

If \tilde{a} is a fuzzy number, the Robust ranking index is defined by

$$R(\tilde{a}) = 0.5 \int_0^1 (a_{\alpha}^L, a_{\alpha}^U) d\alpha$$

Where $(a_{\alpha}^L, a_{\alpha}^U) = \{2\alpha(a_2 - a_1) + a_1, -2\alpha(a_5 - a_4) + a_5, 2\alpha(a_3 - a_2) + 2a_2 - a_3, 2\alpha(a_4 - a_3) - a_4 + 2a_3\}$ is the α -cut of the fuzzy number \tilde{a} .

Definition:2.6

Given a fuzzy set ' \tilde{A} ' defined on ' X ' and any $\alpha \in [0,1]$, the α -cut is denoted by $\tilde{A}(\alpha)$ and is defined as, $\tilde{A}(\alpha) = \{x: \mu(x) \geq \alpha, \alpha \in [0,1]\}$.

III. FUZZY NON-LINEAR PROGRAMMING PROBLEMS

The fuzzy nonlinear programming problem (FNPP) defined as

Maximize (or) Minimize $\tilde{f}(x) = \tilde{f}(x_1, x_2, \dots, x_n)$

Subject to

$\tilde{g}_j(\tilde{x}_j) \leq \text{or } \approx \text{or } \geq b_i; i=1, 2, \dots, m$

$\tilde{x}_j \geq 0$ for all $j=1, 2, \dots, n$

The fuzzy version Kuhn-Tucker conditions given below,

Kuhn-Tucker condition 1: $\nabla f(x) - \sum_{j=1}^n \mu_j \nabla g_j(x) = 0$ (1)

Kuhn-Tucker condition 2: $\mu_j \nabla g_j(x) = 0, \forall j=1, 2, 3, \dots, n$ (2)

Kuhn-Tucker condition 3: $g_j(x) - b_i \leq 0, j=1, 2, 3, \dots, n$ (3)

IV. NUMERICAL EXAMPLE

Consider a fuzzy nonlinear programming problem

Max $\tilde{w} = (1,1,2,3,4)\tilde{c}_1^2 + (1,1,2,2,3)\tilde{c}_2^2$

Subject to

$(0,1,2,2,3)\tilde{c}_1 + (2,3,3,4,5)\tilde{c}_2 \leq (3,4,5,5,6)$

$(1,2,3,3,4)\tilde{c}_1 - (0,1,1,2,2)\tilde{c}_2 \leq (1,2,3,4,5)$ and

$\tilde{c}_1, \tilde{c}_2 \geq 0$

Now convert the given fuzzy problem into crisp form using the Robust

Ranking method

Max $\tilde{w} = R(1,1,2,3,4)\tilde{c}_1^2 + R(1,1,2,2,3)\tilde{c}_2^2$

Subject to

$R(0,1,2,2,3)\tilde{c}_1 + R(2,3,3,4,5)\tilde{c}_2 \leq R(3,4,5,5,6)$

$R(1,2,3,3,4)\tilde{c}_1 - R(0,1,1,2,2)\tilde{c}_2 \leq R(1,2,3,4,5)$ and

$R(\tilde{a}) = 0.5 \int_0^1 (a_\alpha^L, a_\alpha^U) d\alpha$

Where $(a_\alpha^L, a_\alpha^U) = \{2\alpha(a_2 - a_1) + a_1, -2\alpha(a_5 - a_4) + a_5, 2\alpha(a_3 - a_2) + 2a_2 - a_3, 2\alpha(a_4 - a_3) - a_4 + 2a_3\}$

Therefore

$R(1,1,2,3,4) = 0.5 \int_0^1 \{2\alpha(0) + 1 - 2\alpha(1) + 4 + 2\alpha(1) + 2 - 2 + 2\alpha(1) - 3 + 2(2)\} d\alpha$

$= 0.5 \int_0^1 \{1 - 2\alpha + 4 + 2\alpha + 2 - 2 + 2\alpha - 3 + 4\} d\alpha$

$= 0.5 \int_0^1 \{6 + 2\alpha\} d\alpha$

$= 3.0$

Similarly, we get

$R(1,1,2,2,3) = 3.0, R(0,1,2,2,3) = 2.5, R(2,3,3,4,5) = 6.0, R(3,4,5,5,6) = 8.5$

$R(1,2,3,3,4) = 4.5, R(0,1,1,2,2) = 3.0, R(1,2,3,4,5) = 9.0$

We obtain crisp nonlinear programming problem is

Max $\tilde{w} = 3.0\tilde{c}_1^2 + 3.0\tilde{c}_2^2$

Subject to

$2.5\tilde{c}_1 + 6.0\tilde{c}_2 \leq 8.5$

$4.5\tilde{c}_1 - 3.0\tilde{c}_2 \leq 9.0$

$\tilde{c}_1, \tilde{c}_2 \geq 0$

By applying KT condition (1) we get

$6.0\tilde{c}_1 - 2.5\phi_1 - 4.5\phi_2 = 0$

$6.0\tilde{c}_1 - 6.0\phi_1 - 3.0\phi_2 = 0$

By applying KT condition (2) we get

$$\phi_1 [2.5\tilde{c}_1 + 6.0\tilde{c}_2 - 8.5] = 0$$

$$\phi_2 [4.5\tilde{c}_1 - 3.0\tilde{c}_2 - 9.0] = 0$$

By applying KT condition (3) we get

$$2.5\tilde{c}_1 + 6.0\tilde{c}_2 - 8.5 \leq 0$$

$$4.5\tilde{c}_1 - 3.0\tilde{c}_2 - 9.0 \leq 0$$

Solving the above equations, we get the optimal solution is

$$\tilde{c}_1 = 2.304 \text{ and } \tilde{c}_2 = 0.456$$

$$\text{Max } \tilde{w} = 3.0\tilde{c}_1^2 + 3.0\tilde{c}_2^2$$

$$\begin{aligned} \text{Max } \tilde{w} &= 3.0(2.304)^2 + 3.0(0.456)^2 \\ &= 15.925 + 0.623 \end{aligned}$$

$$\text{Max } \tilde{w} = 16.548$$

V. CONCLUSION

We used the proposed method in this research to achieve optimal solution to non-linear programming problem with linear constraints, and we confirmed the accuracy of the proposed method using numerical examples.

REFERENCES

- [1] Annie Christi M. S. (2017), Solutions of Fuzzy Transportation Problem Using Best Candidates Method and Different Ranking Techniques, World Academy of Science, Engineering and Technology International Journal of Mathematical and Computational Sciences,11(4).
- [2] Arun Pratap Singh (2021), A Comparative Study of Centroid Ranking Method and Robust Ranking Technique in Fuzzy Assignment Problem, Global Journal of Technology and Optimization, 12(3),
- [3] Bellman, R. E., Zadeh, L. A. (1970), Decision-making in a fuzzy environment, Management Science, 17(4), pp.141-164.
- [4] Beaula Thangaraj, and V. Vijaya (2012). "Critical Path in a project network using a new representation for Trapezoidal Fuzzy Numbers." International Journal of Mathematics Research 4.5:549-557.
- [5] Beaula Thangaraj, and V. Vijaya (2015). "A new method to find critical path from multiple paths in project networks." Int J Fuzzy Math Arch 9.2: 235-243.
- [6] Beaula, T., and V. Vijaya (2013), "A Study on Exponential Fuzzy Numbers Using α -Cuts." International Journal of Applied 3.2: 1-13.
- [7] DaruneeHunwisai& Poom Kumam (2017), A method for solving a fuzzy transportation problem via Robust ranking technique and ATM, Cogent Mathematics.
- [8] Jayamani,K.Ashwini, N.Srinivasan (2017), Method for Solving Fuzzy Assignment Problem using ones Assignment Method and Robusts Ranking Technique, Journal of Applied Science and Engineering Methodologies,3(2), pp.488-501.
- [9] Kokila G. & R. Anitha (2017), A New Method for solving fuzzy Octogonal number using Robust ranking method, International Journal of Current Research and Modern Education.
- [10] Lalitha M., Dr. C. Loganathan September (2016),An Objective Fuzzy Nonlinear Programming Problem with Symmetric Trapezoidal Fuzzy Numbers,International Journal of Mathematics Trends and Technology, 37 (1),pp.29-35.
- [11] Malini S. U. &Felbin C. Kennedy (2013), An Approach for Solving Fuzzy Transportation Problem Using Octagonal Fuzzy Numbers, Applied Mathematical Sciences, Vol 7, pp.2661 - 2673.
- [12] Nemat Allah Taghi-Nezhad (2018), Fatemeh Taleshian, A Solution Approach for Solving Fully Fuzzy Quadratic Programming Problems, Journal of Applied Research on Industrial Engineering,5(1), pp.50-61.
- [13] Sanjaya Kumar Behera and Jyoti Ranjan nayak (2012), Optimal solution of fuzzy nonlinear programming problems with linear constraints, International Journal of advances in Science and Technology, 4(2).
- [14] Ms. Swati Subhash Desai, Dr. N.N.Pandey February (2011), Applications of Linear Programming Problems and Non Linear Programming Problems in Industry, Variorum – Multi- Disciplinary e-Research Journal, 1(3).
- [15] Umamaheswari P. andK. Ganesan (2017), A Solution Approach to Fuzzy Nonlinear Programming Problems, International Journal of Pure and Applied Mathematics, 113(13), pp.291-300.
- [16] Vijaya, V., and D. Rajalaxmi (2022). "Decision making in fuzzy environment using Pythagorean fuzzy numbers." Mathematical Statistician and Engineering Applications 71.4: 846-854.
- [17] Vijaya, V., D. Rajalaxmi (2022), and H. Manikandan. "Finding critical path in a fuzzy project network using neutrosophic fuzzy number." Advances and Applications in Mathematical Sciences 21.10: 5743-5753.
- [18] S. Yahya Mohamed and M. Divya (Nov. - Dec.2016), Solving Fuzzy Travelling Salesman Problem Using Octagon Fuzzy Numbers with alpha-Cut and Ranking Technique, IOSR Journal ofMathematics, 12(6), PP 52-56.
- [19] Zadeh, L.A. (1965), Fuzzy Sets, Information and Control, 8 (3), pp. 338-353.
- [20] Zimmermann, H.J, Fuzzy Set Theory and Its Applications, (4th Ed.).



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)