



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 10 **Issue:** VIII **Month of publication:** August 2022

DOI: <https://doi.org/10.22214/ijraset.2022.46253>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

A Note on the Wedderburn Decomposition and Unit Group of the Semisimple Group Algebra $\mathbb{F}_p S_n$

Shubham Mittal

Department of Mathematics, IISER Mohali

Abstract: In this short note, we give an algorithm to compute the unit group of a semisimple group algebra $\mathbb{F}_p S_n$ for any $n \geq 5$. To show the practicality of the algorithm, we explicitly find the unit groups of the semisimple group algebras $\mathbb{F}_p S_5$, $\mathbb{F}_p S_6$, $\mathbb{F}_p S_7$ and $\mathbb{F}_p S_8$.

Keywords: Group algebra, Wedderburn decomposition, Unit group, Symmetric group, Young-Tableaux

I. UNIT GROUP OF $\mathbb{F}_p S_n$

Let $\mathbb{F}_p$ be a field, where p is a prime. The study of unit groups of the semisimple group algebras is a classical research problem (cf. [1, 7, 15, 8, 9, 10, 11, 13] and the references therein). The unit groups of semisimple group algebras of symmetric groups S_3 and S_4 have been studied in [5, 14]. In this paper, we continue in this direction and deduce the Wedderburn decomposition of the semisimple group algebra $\mathbb{F}_p S_n$ for any $n \geq 5$ and as a by-product obtain the unit group. It is well known that for a finite group G , the group algebra $\mathbb{F}_p G$ is semisimple if and only if p does not divide $|G|$ [6]. Consequently, $\mathbb{F}_p G$ can be written as a direct sum of matrix rings over finite fields of characteristic p . To be more precise, we have $\mathbb{F}_p G \cong \bigoplus_{t=1}^j M_{n_t}(\mathbb{F}_{p^{f_t}})$ for any semisimple group algebra $\mathbb{F}_p G$. Let ℓ be the exponent of G and ζ be the primitive ℓ^{th} root of unity. In accordance to [2], we define $I_{\mathbb{F}} = \{r \mid \mathbb{F} \mapsto \mathbb{F}^r \text{ is an automorphism of } \mathbb{F}(\zeta) \text{ over } \mathbb{F}\}$. An element $h \in G$ is a p -regular element if its order is not divisible by p . For a p -regular element $h \in G$, let the sum of all the conjugates of h be denoted by γ_h , and the cyclotomic- $\mathbb{F}$ class of γ_h be denoted by $S(\gamma_h) = \{\gamma_{hr} \mid r \in I_{\mathbb{F}}\}$. On utilizing [2, Proposition 1.2] and the result that symmetric group S_n splits over every field, one can see that $\mathbb{F}_p S_n \cong \bigoplus_{t=1}^j M_{n_t}(\mathbb{F}_p)$ (i.e. $p_t = p$ for all $t = 1$ to j), where j is the number of conjugacy classes of S_n that are further equal to partitions $p(n)$ of n . Consequently, the n_t 's appearing in the Wedderburn decomposition of $\mathbb{F}_p S_n$ are nothing but the dimensions of irreducible representations of $\mathbb{F}_p S_n$. To this end, we now show the following.

Theorem 1. The n_t 's appearing in the Wedderburn decomposition of $\mathbb{F}_p S_n$ are equal to the number of standard Young-Tableaux of all the partitions of n .

Proof. It is known that the p -restricted partitions of n indexes the irreducible representations of $\mathbb{F}_p S_n$ (cf. [4]). The p -restricted partitions of n are all those integer partitions in which the difference between successive parts can be at most $p - 1$. It turns out that in the semisimple case (since $p \geq n + 1$), the irreducible representations are given by the Specht modules (cf. [4] for their definition) - in the modular case, the irreducible representations are quotients of Specht modules (obtained by modular reduction of a lattice in $\mathbb{Q}$ -irreducible). Moreover, the quotients of Specht modules are indexed by partitions of n and the respective dimension is equal to the number of standard Young-Tableaux of the respective partition type. Further, the number of standard Young-Tableaux of a partition can be obtained by Hook length formula [12] associated with a Young diagram [3] obtained from the partition. All in all, n_t 's are equal to the number of standard Young-Tableaux of all the partitions of n . $\square$


To this end, due to Theorem 1, we are now ready to give an algorithm to compute the unit group of semisimple group algebra $\mathbb{F}_p S_n$ for any n . Using this algorithm, we explicitly calculate the unit groups of the semisimple group algebras $\mathbb{F}_p S_5$, $\mathbb{F}_p S_6$, $\mathbb{F}_p S_7$ and $\mathbb{F}_p S_8$. As usual, we denote the group of general linear matrices of order r over field $\mathbb{F}_p$ by $GL(r, \mathbb{F}_p)$.

Algorithm: **Input:** positive integer n and a prime p such that $\mathbb{F}_p S_n$ is semisimple.

- write all the partitions of n (total partitions are $p(n)$)
- draw "Young's diagram" for each partition
- calculate "number of standard Young-Tableaux" for each partition using Hook length formula. Let these be $(r_1, \dots, r_{p(n)})$.

Output: The Wedderburn decomposition of $\mathbb{F}_p S_n$ is isomorphic to $\bigoplus_{i=1}^{p(n)} M_{r_i}(\mathbb{F}_p)$ and the unit group of $\mathbb{F}_p S_n$ is isomorphic to $\bigoplus_{i=1}^{p(n)} GL(r_i, \mathbb{F}_p)$.

In order to show the worthiness of above algorithm, first, we deduce the unit group of $\mathbb{F}_p S_6$ for any $p > 5$. Clearly, the partitions $p(6)$ of 6 are 11 and listed as follows: 6, (5,1), (4,2), (4,1,1), (3,3), (3,2,1), (3,1,1,1), (2,2,2), (2,2,1,1), (2,1,1,1,1), (1,1,1,1,1,1). The respective Young's diagram corresponding to these partitions are:



Using Hook length formula, the respective number of standard Young-Tableaux of a partition corresponding to above diagrams are 1,5,9,10,5,16,10,5,9,5 and 1. Therefore, the Wedderburn decomposition of the semisimple group algebra $\mathbb{F}_p S_6$ is isomorphic to $\mathbb{F}_p^2 \oplus M_5(\mathbb{F}_p)^4 \oplus M_9(\mathbb{F}_p)^2 \oplus M_{10}(\mathbb{F}_p)^2 \oplus M_{16}(\mathbb{F}_p)$. Consequently, the unit group

$$U(\mathbb{F}_p S_6) \cong (\mathbb{F}_p^*)^2 \oplus GL(5, \mathbb{F}_p)^4 \oplus GL(9, \mathbb{F}_p)^2 \oplus GL(10, \mathbb{F}_p)^2 \oplus GL(16, \mathbb{F}_p).$$

Next, we deduce the unit group of $\mathbb{F}_p S_5$ for any $p > 5$. Clearly, the partitions $p(5)$ of 5 are 7 and listed as follows: 5, (4,1), (3,2), (3,1,1), (2,2,1), (2,1,1,1), and (1,1,1,1,1). Further, one can verify that the respective number of standard Young-Tableaux of a partition corresponding to Young diagrams in this case are 1,4,5,6,5,4 and 1. Therefore, the Wedderburn decomposition of the semisimple group algebra $\mathbb{F}_p S_5$ is isomorphic to $\mathbb{F}_p^2 \oplus M_4(\mathbb{F}_p)^2 \oplus M_5(\mathbb{F}_p)^2 \oplus M_6(\mathbb{F}_p)$. Consequently, the unit group

$$U(\mathbb{F}_p S_5) \cong (\mathbb{F}_p^*)^2 \oplus GL(4, \mathbb{F}_p)^2 \oplus GL(5, \mathbb{F}_p)^2 \oplus GL(6, \mathbb{F}_p).$$

Finally, using our algorithm, one can verify that for the groups S_7 and S_8 , the unit groups of semisimple group algebras $\mathbb{F}_p S_7$ and $\mathbb{F}_p S_8$ are as follows:

- for $p > 7$, $U(\mathbb{F}_p S_7)$ is isomorphic to

$$(\mathbb{F}_p)^2 \oplus GL(6, \mathbb{F}_p)^2 \oplus GL(14, \mathbb{F}_p)^4 \oplus GL(15, \mathbb{F}_p)^2 \oplus GL(20, \mathbb{F}_p) \oplus GL(21, \mathbb{F}_p)^2 \oplus GL(35, \mathbb{F}_p)^2.$$

• for $p > 7$, $U(\mathbb{F}_p S_8)$ is isomorphic to

$$(\mathbb{F}_p^*)^2 \oplus GL(7, \mathbb{F}_p)^2 \oplus GL(14, \mathbb{F}_p)^2 \oplus GL(20, \mathbb{F}_p)^2 \oplus GL(21, \mathbb{F}_p)^2 \oplus GL(28, \mathbb{F}_p)^2 \oplus GL(35, \mathbb{F}_p)^2 \oplus GL(42, \mathbb{F}_p) \oplus GL(56, \mathbb{F}_p)^2 \oplus GL(64, \mathbb{F}_p)^2 \oplus GL(70, \mathbb{F}_p)^2 \oplus GL(90, \mathbb{F}_p).$$

II. DISCUSSION


We have given a simple algorithm with which one can easily characterize the unit group of the semisimple group algebra $\mathbb{F}_p S_n$ for any n . Therefore, this paper settles one of the stated problems of [8] related to the unit group of the group algebras of symmetric groups. The results proved in this paper are for finite fields of the form $\mathbb{F}_p$, however, one can see that the Wedderburn decomposition of the semisimple group algebra $\mathbb{F}_p S_n$ is independent of the order of finite field. To be more precise, the field under consideration can have p elements or p^k elements for some k .

III. ACKNOWLEDGMENT

I would like to thank Dr. Gaurav Mittal for his valuable suggestions and unsolicited support to organize this paper.

REFERENCES

- [1] G. K. Bakshi, S. Gupta, I. B. S. Passi, "The algebraic structure of finite metabelian group algebras", Comm. Algebra, 43(1), (2015), 2240-2257.
- [2] R. A. Ferraz, "Simple components of the center of $\mathbb{F}G/J(\mathbb{F}G)$ ", Comm. Algebra, 36(9), 3191-3199, 2008.
- [3] W. Fulton, Young tableaux, with applications to representation theory and geometry, Cambridge University Press, 1997.
- [4] G. D. James, The representation theory of the symmetric groups, Springer, 1978.
- [5] M. Khan, R. K. Sharma, J. B. Srivastava, "The unit group of $\mathbb{F}S_4$ ", Acta Math. Hungar., 118 (1-2), 105-113, 2008.
- [6] C. P. Milies, S. K. Sehgal, An introduction to group rings, Kluwer Acad. Pub., 2002.
- [7] G. Mittal, R. K. Sharma, "On unit group of finite group algebras of non-metabelian groups upto order 72", Math. Bohemica, 146(4), 429-455, 2021.
- [8] G. Mittal, R. K. Sharma, "Unit group of semisimple group algebras of some non-metabelian groups of order 120", Asian-European J. Math., 15(3), 2250059, 2022.
- [9] G. Mittal, R. K. Sharma, "Computation of Wedderburn decomposition of groups algebras from their subalgebra", Bull. Korean Math. Soc., 59(3), 781-787 2022.
- [10] G. Mittal, R. K. Sharma, "Wedderburn decomposition of a semisimple group algebra $\mathbb{F}_q G$ from a subalgebra of factor group of G ", Int. Elect. J. Algebra, 32, 91-100, 2022.
- [11] G. Mittal, R. K. Sharma, "On unit group of finite group algebras of non-metabelian groups of order 108", J. Algebra Comb. Discrete Appl., 8(2), 59-71, 2021.
- [12] B. Sagan, The Symmetric Group. Representations, Combinatorial Algorithms, and Symmetric Functions, Springer-Verlag, 2001.
- [13] M Sahai, SF Ansari, "Unit groups of the finite group algebras of generalized quaternion groups", J. Algebra Its Appl., 19(6), 2020.
- [14] R. K. Sharma, J. B. Srivastava, M. Khan, "The unit group of $\mathbb{F}S_8$ ", Acta Mathematica Academiae Paedagogicae Nyiregyhaziensis, 23(2), 129-142, 2007.
- [15] R. K. Sharma, G. Mittal, "Unit group of semisimple group algebra $\mathbb{F}_q SL(2, 5)$ ", Math. Bohemica, 147(1), 1-10, 2022.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)