



# **iJRASET**

International Journal For Research in  
Applied Science and Engineering Technology



---

# **INTERNATIONAL JOURNAL FOR RESEARCH**

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume:** 12    **Issue:** III    **Month of publication:** March 2024

**DOI:** <https://doi.org/10.22214/ijraset.2024.58887>

**[www.ijraset.com](http://www.ijraset.com)**

**Call:** ☎ 08813907089

**E-mail ID:** [ijraset@gmail.com](mailto:ijraset@gmail.com)

# Observations on Ternary Quadratic Diophantine

## Equation $8(x^2 + y^2) - 15xy + 7x + 7y + 49 = 40z^2$

C.Saranya<sup>1</sup>, R.Salini<sup>2</sup>

<sup>1</sup>Assistant Professor, <sup>2</sup>PG student, PG and Research Department of Mathematics, Cauvery College for Women (Autonomous)  
(Affiliated to Bharathidasan University), Tiruchirappalli, Tamil Nadu, India

**Abstract:** The Ternary Quadratic Diophantine Equation is given by  $8(x^2 + y^2) - 15xy + 7x + 7y + 49 = 40z^2$ .

Number of non-zero distinct integer solutions. Some interesting relations among the solutions and the way of factorization, along with the illusion of non-zero distinct integer solutions to the above equations are obtained.

**Keywords:** Diophantine equation, Integral solutions, Quadratic equation with three unknowns, Ternary equation.

### I. INTRODUCTION

Diophantine equation have provoked the interest of different mathematicians. It is simple to solve Diophantine equations with a degree greater than three by reducing them to equations degree 2 or 3. In [1-3], hypothesis of numbers is talked about. Diophantine quadratic equations are discussed in [5,6, and 12]. In [4,7-11] cubic, biquadratic and higher order equations are considered for its integer solutions

This correspondence deals with one more attracting ternary quadratic equation addressing a non-homogeneous cone for its infinitely non-zero integer points. In like, manner, a couple of fascinating relations among the arrangements are correlated.

### II. NOTATIONS

$T_{8,n}$  = Octagonal number of rank 'n'.

$T_{10,n}$  = Decagonal number of rank 'n'.

$T_{18,n}$  = Octadecagonal number of rank 'n'.

$T_{22,n}$  = Icosidigonal number of rank 'n'.

$T_{26,n}$  = Icosihexagonal number of rank 'n'.

$T_{28,n}$  = Icosioctagonal number of rank 'n'.

$Gno_n$  = Gnomonic number of rank 'n'.

### III. METHOD OF ANALYSIS

The ternary quadratic Diophantine equation for its non-zero integral solution is given by

$$8(x^2 + y^2) - 15xy + 7x + 7y + 49 = 40z^2 \quad (1)$$

The replacement of linear transformations  $x = u + v$  and  $y = u - v$  (2)

in (1) we get,

$$(u + 7)^2 + 31v^2 = 40z^2 \quad (3)$$

Below are four examples of distinct non-zero integer solutions to (1).

PATTERN: 1

Assume  $z = z(a, b) = a^2 + 31b^2$  (4)

wherein (a, b) is a pair of non-zero integers.

$$40 = (3 + i\sqrt{31})(3 - i\sqrt{31}) \quad (5)$$

Using (4) and (5) in (3), by the method of factorization,

$$\left((u+7) + i\sqrt{31}v\right)\left((u+7) - i\sqrt{31}v\right) = (3 + i\sqrt{31})(3 - i\sqrt{31}) \left[ \left(a + i\sqrt{31}b\right)^2 \left(a - i\sqrt{31}b\right)^2 \right] \quad (6)$$

When we compare real and imaginary components and put like terms together, we get

$$u = u(a, b) = 3a^2 - 93b^2 - 62ab - 7$$

$$v = v(a, b) = a^2 - 31b^2 + 6ab$$

The corresponding integer solutions of equation (1) are given by using the values of  $u$  &  $v$  in equation (2).

$$x = x(a, b) = 4a^2 - 124b^2 - 56ab - 7$$

$$y = y(a, b) = 2a^2 - 62b^2 - 68ab - 7$$

$$z = z(a, b) = a^2 + 31b^2$$

#### OBSERVATIONS:

1.  $x(a, a) - y(a, a) + z(a, a) + 4T_{10,a} \equiv 0 \pmod{12}$
2.  $y(1, 1) - x(1, 1) - z(1, 1)$  is a perfect square.
3.  $x(a, 1) - y(a, 1) + z(a, 1) - T_{8,a} - 7Gno_a \equiv 0 \pmod{24}$
4.  $x(a, a) - y(a, a) + 4T_{26,a} \equiv 0 \pmod{44}$
5.  $x(a, a) - y(a, a) - z(a, a) + 10T_{18,a} \equiv 0 \pmod{70}$

#### PATTERN:2

Consider (5), we write 40 as

$$40 = \frac{(19 + i3\sqrt{31})(19 - i3\sqrt{31})}{16} \quad (7)$$

The corresponding integer solutions of equation (1) are given by using the values of  $u$  &  $v$  in equation (2).

$$x = x(A, B) = 88A^2 - 2728B^2 - 592AB - 7$$

$$y = y(A, B) = 64A^2 - 1984B^2 - 896AB - 7$$

$$z = z(A, B) = 16A^2 + 496B^2$$

#### OBSERVATIONS:

1.  $x(1, 1) - y(1, 1) + z(1, 1)$  is a nasty number
2.  $y(A, A) - x(A, A) - 32T_{28,A} \equiv 0 \pmod{384}$

3.  $x(A, A) - y(A, A) + z(A, A) - 8T_{26,A} \equiv 0 \pmod{88}$
4.  $x(A, A) - y(A, A) - z(A, A) + 116T_{18,A} \equiv 0 \pmod{812}$
5.  $x(A, 1) - y(A, 1) - 2T_{26,A} - 163Gno_A \equiv 0 \pmod{581}$

PATTERN: 3

Consider(5), we write 40 as,

$$40 = \frac{(7 + i9\sqrt{31})(7 - i9\sqrt{31})}{64} \quad (8)$$

The corresponding integer solutions of equation (1) are given by using the values of  $u$  &  $v$  in equation (2)

$$x = x(A, B) = 128A^2 - 3968B^2 - 4352AB - 7$$

$$y = y(A, B) = -16A^2 + 496B^2 - 4576AB - 7$$

$$z = z(A, B) = 64A^2 + 1984B^2$$

OBSERVATIONS:

1.  $y(1, 1) - x(1, 1)$  is a perfect square.
2.  $x(A, A) - y(A, A) + z(A, A) + 256T_{18,A} \equiv 0 \pmod{1792}$
3.  $x(A, 1) - y(A, 1) - 12T_{26,A} - 178Gno_A \equiv 0 \pmod{4286}$
4.  $x(A, A) - y(A, A) + 1024T_{10,A} \equiv 0 \pmod{3072}$
5.  $x(A, A) - y(A, A) - z(A, A) + 512T_{26,A} \equiv 0 \pmod{5632}$

PATTERN: 4

Consider(5), we write 40 as

$$40 = \frac{(63 + i\sqrt{31})(63 - i\sqrt{31})}{100} \quad (9)$$

The corresponding integer solutions of equation (1) are given by using the values of  $u$  &  $v$  in equation (2).

$$x = x(A, B) = 640A^2 - 19840B^2 + 640AB - 7$$

$$y = y(A, B) = 620A^2 - 19220B^2 - 1880AB - 7$$

$$z = z(A, B) = 100A^2 + 3100B^2$$

OBSERVATIONS:

1.  $x(A, A) - y(A, A) + z(A, A) - 640T_{18,A} \equiv 0 \pmod{4480}$
2.  $x(A, A) - y(A, A) - 160T_{26,A} \equiv 0 \pmod{1760}$
3.  $x(A, 1) - y(A, 1) - z(A, 1) + 10T_{18,A} - 1225Gno_A \equiv 0 \pmod{2495}$
4.  $x(A, A) - y(A, A) - z(A, A) + 128T_{22,A} \equiv 0 \pmod{1152}$

$$5. \quad x(A,1) - z(A,1) - 54T_{22,A} - 563Gno_A \equiv 0 \pmod{22384}$$

#### IV. CONCLUSION

Four distinct patterns of non-zero distinct integer solutions to the non-homogeneous cone given by are presented in this paper. To finish up, one might look for different examples of non-zero number unmistakable arrangements and their relating properties for different decisions of ternary quadratic Diophantine conditions.

#### REFERENCES

- [1] Carmichael, R.D., The theory of numbers and Diophantine Analysis, Dover Publications, New York, 1959.
- [2] Dickson L.E, History of Theory of Numbers, Vol.11, Chelsea Publishing company, New York, 1952.
- [3] Mordell. L.J, Diophantine equations, Academic Press, London, 1969 Telang, S.G., Number theory, Tata McGraw Hill publishing company, New Delhi, 1996.
- [4] Gopalan.M.A., Vidhyalakshmi.S and Umarani.J., "On ternary Quadratic Diophantine equation  $6(x^2 + y^2) - 8xy = 21z^2$ ", Sch.J. Eng. Tech. 2(2A); 108- 112, 2014.
- [5] Janaki.G and Saranya.C., Observations on the Ternary Quadratic Diophantine Equation  $6(x^2 + y^2) - 11xy + 3x + 3y + 9 = 72z^2$ , International Journal of Innovative Research in Science, Engineering and Technology, Vol-5, Issue-2, Pg.no: 2060-2065, Feb 2016.
- [6] Janaki.G and Vidhya.S., On the integer solutions of the homogeneous biquadratic diophantine equation  $x^4 - y^4 = 82(z^2 - w^2)p^2$ , International Journal of Engineering Science and Computing, Vol. 6, Issue 6, pp.7275-7278, June, 2016.
- [7] Gopalan.M.A and Janaki.G, Integral solutions of  $(x^2 - y^2)(3x^2 + 3y^2 - 2xy) = 2(z^2 - w^2)p^3$ , Impact J.Sci.,Tech., 4(1), 97-102, 2010.
- [8] Janaki.G and Saranya.P., On the ternary Cubic diophantine equation  $5(x^2 + y^2) - 6xy + 4(x + y) + 4 = 40z^3$ , International Journal of Science and Research- online, Vol 5, Issue 3, Pg.No:227-229, March 2016.
- [9] Janaki.G and Saranya.C., Integral Solutions of the non-homogeneous heptic equation with five unknowns  $5(x^3 - y^3) - 7(x^2 + y^2) + 4(z^3 - w^3 + 3wz - xy + 1) = 972p^7$ , International Journal of Engineering Science and Computing, Vol. 6, Issue 5, pp.5347-5349, May, 2016.
- [10] Janaki.G and Saranya.C., Integral Solutions of the ternary cubic equation  $3(x^2 + y^2) - 4xy + 2(x + y + 1) = 972z^3$ , International Research Journal of Engineering and Technology, Vol. 4, Issue 3, pp.665-669, March, 2017.
- [11] Janaki.G and Saranya.C., Integral Solutions of the homogeneous biquadratic diophantine equation  $3(x^4 - y^4) - 2xy(x^2 - y^2) = 972(z + w)p^3$ , International Journal for Research in Applied Science and Engineering Technology, Vol. 5, Issue 8, pp.1123-1127, Aug 2017.





10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)