iJRASET
International Journal For Research in Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 12 **Issue:** III **Month of publication:** March 2024

DOI: https://doi.org/10.22214/ijraset.2024.58886

www.ijraset.com

Call: 08813907089 | E-mail ID: ijraset@gmail.com

International Journal for Research in Applied Science & Engineering Technology (IJRASET)
ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538
Volume 12 Issue III Mar 2024- Available at www.ijraset.com

Observations on Ternary Quadratic Diophantine Equation $12(x^2+y^2)-23xy+2x+2y+4=56z^2$

C. Saranya[1], M. Janani[2]
[1]*Assistant Professor* , [2]*PG Student, PG and Research Department of Mathematics, Cauvery College for Women (Autonomous) (Affiliated to Bharathidasan University), Tiruchirappalli, Tamil Nadu, India*

Abstract: ***The Ternary Quadratic Diophantine Equation*** $12(x^2+y^2)-23xy+2x+2y+4=56z^2$ ***is analyzed for its infinite number of non-zero integral solutions. Four interesting patterns satisfying the cone are identified. There are a few interesting connections between the solutions and some unique number patterns.***
Keywords: Diophantine equation, Integral solutions, Quadratic equation with three unknowns, Ternary equation.
.

I. INTRODUCTION

Diophantine equations have stimulated the interest of various mathematicians. Diophantine equations with higher degree greater than three can be reduced in to equations of degree 2 or 3 and it can be easily solved. In [1-3], theory of numbers is discussed. In [5,6 & 12], quadratic Diophantine equations are discussed. In [4, 7-11], cubic, biquadratic and higher order equations are considered for its integral solutions.

This communication concerns with yet another interesting ternary quadratic equation $12(x^2+y^2)-23xy+2x+2y+4=56z^2$ representing a non-homogeneous cone is determined for its infinitely many non-zero integral points. Likewise, a few interesting relations among the solutions are analyzed.

II. NOTATIONS

$T_{6,n}$ = Hexgonal number of rank ‘n’.

$T_{8,n}$ = Octogonal number of rank ‘n’.

$T_{10,n}$ = Decagonal number of rank ‘n’.

$T_{14,n}$ = Tetradecogonal number of rank ‘n’.

$T_{16,n}$ = Hexadecagonal number of rank ‘n’.

$T_{18,n}$ = Octadecagonal number of rank ‘n’.

$T_{22,n}$ = Icosidigonal number of rank ‘n’.

$T_{24,n}$ = Icositetragonal number of rank ‘n’.

$T_{26,n}$ = Icosihexagonal number of rank ‘n’.

Gno_n = Gnomonic number rank ‘n’.

III. METHOD OF ANALYSIS

The ternary quadratic diophantine equation to be tackled for its non-zero integral solution is

$$12(x^2+y^2)-23xy+2x+2y+4=56z^2 \quad (1)$$

using the linear transformations

$$x=u+v \quad \text{and} \quad y=u-v \quad (2)$$

in (1) leads to, $$(u+2)^2+47v^2=56z^2 \quad (3)$$

We illustrate below four different patterns of non-zero distinct integer solutions to (1)

©IJRASET: All Rights are Reserved

A. Pattern : 1

Assume $z = z(a,b) = a^2 + 47b^2$ (4)

where a and b are non-zero integers.

and write $56 = (3+i\sqrt{47})(3-i\sqrt{47})$ (5)

Using (4) and (5) in (3), and using factorization method,

$$\left((u+2)+i\sqrt{47}v\right)\left((u+2)-i\sqrt{47}v\right) = (3+i\sqrt{47})(3-i\sqrt{47})\left[\left(a+i\sqrt{47}b\right)^2\left(a-i\sqrt{47}b\right)^2\right] \quad (6)$$

Equating the like terms and comparing real and imaginary parts, we get

$$u = u(a,b) = 3a^2 - 141b^2 - 94ab - 2$$

$$v = v(a,b) = a^2 - 47b^2 + 60ab$$

Substituting the above values of u & v in equation (2), the corresponding integer solutions of (1) are given by

$$x = x(a,b) = 4a^2 - 188b^2 - 88ab - 2$$

$$y = y(a,b) = 2a^2 - 94b^2 - 100ab - 2$$

$$z = z(a,b) = a^2 + 47b^2$$

Observations

1. $x(a,a) - y(a,a) + z(a,a) + 4T_{18,a} \equiv 0 \pmod{28}$
2. $2y(1,1) - 2x(1,1) - 2z(1,1)$ is a perfect square.
3. $y(a,a) + z(a,a) - x(a,a) - 32T_{10,a} \equiv 0 \pmod{96}$
4. $y(a,a) - x(a,a) - 8T_{22,a} \equiv 0 \pmod{72}$
5. $x(a,1) - y(a,1) + z(a,1) - T_{8,a} - 7Gno_a \equiv 0 \pmod{40}$
6. y(1,1)-x(1,1) is a duck number

B. Pattern : 2

We substitute (5) with 56 as

$$56 = \frac{(29+i5\sqrt{47})(29-i5\sqrt{47})}{36} \quad (7)$$

Following the methodology introduced over, the relating non-zero particular whole number answers for (1) are Gotten as

$$x = x(A,B) = 204A^2 - 9588B^2 - 2472AB - 2$$

$$y = y(A,B) = 144A^2 - 6768B^2 - 3168AB - 2$$

$$z = z(A,B) = 36A^2 + 1692B^2$$

Observations

1. $x(A,A) - y(A,A) + z(A,A) + 48T_{16,A} \equiv 0 \pmod{288}$
2. $x(A,1) - y(A,1) - 6T_{22,A} - 375Gno_A \equiv 0 \pmod{2445}$
3. $x(A,A) - y(A,A) + 172T_{26,A} \equiv 0 \pmod{1892}$
4. $y(A,1) - x(A,1) - z(A,1) + 16T_{14,A} + 388Gno_A \equiv 0 \pmod{740}$
5. Each of the following expressions represents a Duck number

i. $x(1,1)-2y(1,1)$

ii. $x(1,1)-2y(1,1)-z(1,1)$

iii. $-y(1,1)-z(1,1)$

C. Pattern : 3

We substitute (5) with 56 as

$$56=\frac{(67+i\sqrt{47})(67-i\sqrt{47})}{81} \tag{8}$$

Following the methodology introduced over, the relating non-zero particular whole number answers for (1) are gotten as

$$x=x(A,B)=612A^2-28764B^2-360AB-2$$

$$y=y(A,B)=594A^2-27918B^2-2052AB-2$$

$$z=z(A,B)=81A^2+38078B^2$$

Observations

1. $x(A,A)-y(A,A)+z(A,A)-132T_{26,A}-726Gno_A\equiv 0 \pmod{38885}$
2. $y(A,A)-x(A,A)+132T_{26,A}\equiv 0 \pmod{1452}$
3. $x(A,1)-z(A,1)-177T_{8,A}-357Gno_A\equiv 0 \pmod{66487}$
4. $4x(1,1)-4y(1,1)$ is a palindromic number
5. $z(1,1)-x(1,1)-y(1,1)$ is a duck number

D. Pattern : 4

We substitute (5) with 56 as

$$56=\frac{(83+i5\sqrt{47})(83-i5\sqrt{47})}{144} \tag{9}$$

Following the methodology introduced over, the relating non-zero particular whole number answers for (1) are gotten as

$$x=x(A,B)=1056A^2-49632B^2-3648AB-2$$

$$y=y(A,B)=936A^2-43992B^2-7632AB-2$$

$$z=z(A,B)=144A^2+6768B^2$$

Observations


1. $x(A,1)-y(A,1)-z(A,1)+2T_{26,A}-1981Gno_A\equiv 0 \pmod{10427}$
2. $x(A,1)-y(A,1)-10T_{24,A}-2047Gno_A\equiv 0 \pmod{3593}$
3. $y(1,1)-x(1,1)+z(1,1)$ is a palindromic number
4. $y(1,1)-x(1,1)$ is a nasty number.
5. $y(1,1)-z(1,1)-2x(1,1)$ is a duck number.

IV. CONCLUSION

This paper discusses four distinct patterns of non-zero distinct integer solutions to the non-homogeneous cone given by $12(x^2+y^2)-23xy+2x+2y-4=56z^2$. To conclude, one may search for other patterns of solutions and their corresponding properties.

REFERENCES

[1] Carmichael, R.D., The theory of numbers and Diophantine Analysis, Dover Publications,New York, 1959.
[2] Dickson L.E, History of Theory of Numbers, Vol.11, Chelsea Publishing company, New York,1952.
[3] Mordell. L.J, Diophantine equations, Academic Press,London,1969 Telang, S.G., Number theory, Tata McGraw Hill publishing company, New Delhi, 1996.
[4] Gopalan.M.A and Janaki.G, Integral solutions of $(x^2-y^2)(3x^2+3y^2-2xy=2(z^2-w^2)p^3$, Impact J.Sci.,Tech., 4(1), 97-102, 2010.
[5] Gopalan. M.A., Vidhyalakshmi.S and Umarani.J., On ternary Quadratic Diophantine equation $6(x^2+y^2)-8xy=21z^2$, Sch.J. Eng. Tech. 2(2A); 108-112,2014
[6] Janaki.G and Saranya.C., Observations on the Ternary Quadratic Diophantine Equation $6(x^2+y^2)-11xy+3x+3y+9=72z^2$, International Journal of Innovative Research in Science, Engineering and Technology, Vol-5, Issue-2, Pg.no: 2060-2065, Feb 2016.
[7] Janaki.G and Saranya.P., On the ternary Cubic Diophantine equation $5(x^2+y^2)-6xy+4(x+y)+4=40z^3$, International Journal of Science and Research- online, Vol 5, Issue 3, Pg.No:227-229, March 2016.
[8] Janaki.G and Vidhya.S., On the integer solutions of the homogeneous biquadratic diophantine equation $x^4-y^4=82(z^2-w^2)p^2$, International Journal of Engineering Science and Computing, Vol. 6, Issue 6, pp.7275-7278, June, 2016.
[9] Janaki.G and Saranya.C., Integral Solutions of the non-homogeneous heptic equation with five unknowns $5(x^3-y^3)-7(x^2+y^2)+4(z^3-w^3+3wz-xy+1)=972p^7$, International Journal of Engineering Science and Computing, Vol. 6, Issue 5, pp.5347-5349, May, 2016.
[10] Janaki.G and Saranya.C., Integral Solutions of the ternary cubic equation $3(x^2+y^2)-4xy+2(x+y+1)=972\ z^3$, International Research Journal of Engineering and Technology, Vol. 4, Issue 3, pp.665-669, March, 2017.
[11] Janaki.G and Saranya.C., Integral Solutions of the homogeneous biquadratic diophantine equation $3(x^4-y^4)-2xy(x^2-y^2)=972(z+w)p^3$, International Journal for Research in Applied Science and Engineering Technology, Vol. 5, Issue 8, pp.1123-1127, Aug 2017.
[12] Saranya. C and Kayathri. P., Observations On Ternary Quadratic Equation $3x^2+2y^2=275z^2$, Advances and Applications in Mathematical Sciences, Vol 21, Issue 3, pp. 1549-1556 January 2022.



10.22214/IJRASET

45.98

IMPACT FACTOR: 7.129

IMPACT FACTOR: 7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)