



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 13 **Issue:** VI **Month of publication:** June 2025

DOI: <https://doi.org/10.22214/ijraset.2025.72431>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

Optimization of Three Component, One Pot Synthesis of aminobenzyl naphthol Exploiting Electrophilicity of azomethines Under Varying Conditions

Sushma Rani¹, M. K. Singh²

Department of Chemistry, Patna University

Abstract: *The synthesis of aminobenzyl naphthols by organocatalysis has generated interest due to the mild reaction conditions and environmental benefits. It has been shown that (1,4-diazacyclo[2.2.2]octane) has demonstrated high efficiency as an organocatalyst for the three component condensation reaction. These reactions lead to the synthesis of a novel class of aminobenzyl naphthols under various solvent conditions offering remarkable advantages like: mild reaction conditions, high yields, selectivity and simplicity.*

This protocol is particularly appealing for the synthesis of complex organic molecules with potential applications in pharmaceuticals and material science. Overall, the use of organocatalysts in aminobenzyl naphthol synthesis presents a versatile approach with potential for the further development maintaining the guidelines of greener chemistry.

Keywords: 2-Naphthol, aminobenzyl naphthol, azomethines, organocatalyst, three component, (1,4-diazabicyclo[2.2.2]octane)

I. INTRODUCTION

Compounds containing 1,3-amino oxygenated functional groups are indeed significant in medicinal chemistry and natural products [1]. These structures often contribute to biological activity due to their ability to participate in hydrogen bonding and interaction with biological targets. The design of synthetic drugs [1-3] often incorporates these functional groups to enhance efficacy and optimize pharmacokinetic properties. Aminonaphthols, commonly referred to as Betti bases [4], are an important class of compounds characterized by a naphthol moiety with an amino group. Aminonaphthols or Betti bases, exhibit a range of beneficial biological properties, including analgesic, antibacterial, hypotensive and bardycardiac activities [5-8] (Fig. 1). The diverse biological activities of aminonaphthols highlight their importance in medicinal chemistry and the potential for further exploration in drug development [4,9].

Since Betti's initial report in 1900, numerous methods have been developed for the synthesis of aminonaphthols. The preparation of aminonaphthols typically involves (Scheme 1) the reaction of 2-naphthol with an aldehyde and aliphatic amine [4] through a process known as an imine. In this reaction, the aldehyde reacts with the amine to form an intermediate imine, which subsequently undergoes nucleophilic attack by the 2-naphthol. Hydrolysis of performed amidoalkyl naphthols is another effective approach to synthesize aminonaphthols [10-19]. The synthesis of amidoalkyl naphthols has generated significant interest due to their versatile applications in various fields, including pharmaceuticals, agrochemicals and material science. These compounds often exhibit biological activities, making them valuable in drug development. Many traditional methods for synthesizing amidoalkyl naphthols often come with challenges including: low product yields, expensive catalysts, long reaction times and laborious workup procedures [4-9]. These disadvantages have prompted researchers to seek more efficient, cost effective methods, such as greener solvents, alternative catalysts, or one pot reactions. The synthesis of aryl aminonaphthols after presents [23-24] additional challenges [20-22] compared to aliphatic counter parts. The synthesis of hetero aryl aminonaphthols using electron rich heteroaromatic amines as amine sources [25-27]. Organocatalysts have indeed demonstrated significant potential in multi-component reaction (MCRs) [28] due to high efficiency and selectivity.

The advantages of using organic catalysts can be: easy preparation or availabilities, milder reaction conditions, selectivity and compatibility with numerous functional groups [29]. Recent studies have highlighted the effectiveness of organocatalysts, particularly derivatives of N,N-dialkylethanolamine, in catalyzing Michael addition [30] and Friedel-Crafts alkylation reactions [31,32].

To the best of our knowledge, there have been reported studies on the synthesis of Betti bases using organocatalysts. Betti bases, which are significant in various applications including medicinal chemistry, typically involve multi-component reaction. Exploring organocatalysts for this synthesis could offer new pathways for efficient and selective production, potentially leveraging their advantages in milder conditions and broad functional group compatibility.

The use of bifunctional organocatalysts for synthesizing alkyl and aryl naphthol derivatives through a one-pot three component [33-37] condensation is intriguing. This method effectively combines β -naphthol, aldehydes, and arylamines, potentially leading to efficient synthesis pathways with minimal byproducts. The bifunctional nature of the organocatalysts likely enhances reactivity and selectivity, streamlining the process. Additionally, characterizing the final products using techniques like NMR or mass spectroscopy would be beneficial for confirming the success of the synthesis.







Fig 1: An aminonaphthols with antipain and antibacterial activities



Scheme 1: Methods for synthesis of aminonaphthols

II. EXPERIMENTAL

All chemicals were purchased from Merck or Aldrich chemical companies and were used without further purification. This ensured convenience and efficiency in the experimental procedures, maintaining the integrity of the synthesized products while minimizing the handling and processing time.

Melting points were determined using a MEL-TEMP model 1202D apparatus and are uncorrected. FTIR spectra were recorded using a Bruker Tensor 27 spectrometer in the form of KBr discs.

The ^1H NMR spectra were recorded on a Bruker Spectrospin Avance 400 spectrometer.

^{13}C NMR spectra were recorded on the instrument at a frequency of 100 MHz. Chemical shifts were reported with respect to solvent signals as internal standards and coupling constant (J) are given in Hz. Multiplicities are indicated as the following: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; dd, doublet of doublets. Preparative layer chromatography was prepared using silica gel (Merk Kieselgel 60 Hf254, no. 7739).

III. RESULT AND DISCUSSION

Our studies started with the reaction of 2-naphthol (2a) with benzaldehyde (13a) and aniline (14a) as our model reaction to investigate. A range of organocatalysts was evaluated under different conditions, using different organic solvents. The results are summarized in table 1.

As given in table 1, different polar organic solvents such as ethanol (EtOH), chloroform (CHCl_3) and dichloromethane (CH_2Cl_2) resulted in low to moderate yields at room temperature (15-45%, entries 1-12). The best results were achieved under non-polar benzene solvent, demonstrating significantly higher yields. Organocatalyst II gave a promising outcomes (60%, entry 14), other organocatalysts I & III produced lower yields (30-35%, entries 13-15) and organocatalysts (IV-VI) provided poor results (45-50%, entries 16-18).





Fig 2: Organocatalysts I-VI

Since the reactions as well as products are air stable, there was no need to carry out the reactions in an inert atmosphere, which simplified the experimental set up and reduced the overall complexity of the process. This characteristic of the solvent benzene makes it suitable for broader applications in this type of organic synthesis. To optimize the reaction conditions, the model reaction was exposed to different amounts of organocatalyst II (5, 7, 8, 10 & 15 mol%) at different temperature using benzene solvent. (table 2) Perusal of suggests in table 2, the best results were obtained when the reaction was performed using benzene as a solvent and 8 mol% of DABCO as the organocatalyst at 55°C . To investigate the scope and generality of this procedure, the condensation reactions of 2-naphthols with a range of aromatic aldehydes and aromatic amines was conducted in the presence of benzene solvent and DABCO as a catalyst (fig2) at 55°C (scheme 3, Table 3). As Shown in table 2, the reactions were carried out efficiently within 40 to 55 minutes, resulting in the production of the desired products in good to excellent yields. This demonstrates the effectiveness of the DABCO catalysed condensation process under solvent. Highlighting its potential for rapid and efficient organic synthesis. (fig3)

Different substituted benzaldehydes successfully reacted with aniline and 2-naphthol, yielding the corresponding products (15a-p) in 80-88% yields.

The reactions demonstrated good functional group tolerance, indicating the versatility of the DABCO catalysed process. This further confirms the methods applicability for a wide range of substrates in organic synthesis.



Scheme 2: synthesis of Betti bases

Table 1: Optimization of organocatalytic synthesis of arylaminonaphthol under various conditions.


Entry	Catalyst (11 mol%)	Solvent	Time(h)	Yield(%)
1	I	CHCl ₃	26	20
2	I	CH ₂ Cl ₂	24	15
3	I	EtOH	22	25
4	II	CHCl ₃	22	45
5	II	CH ₂ Cl ₂	26	45
6	II	EtOH	24	45
7	III	CHCl ₃	26	20
8	IV	CHCl ₃	22	25
9	V	CHCl ₃	24	35
10	VI	CHCl ₃	22	40
11	VI	CH ₂ Cl ₂	26	35
12	VI	EtOH	24	40
13	I	C ₆ H ₆	2	30
14	II	C ₆ H ₆	2	60
15	III	C ₆ H ₆	2	35
16	IV	C ₆ H ₆	2	45
17	V	C ₆ H ₆	2	49
18	VI	C ₆ H ₆	2	50

*reaction was carried out at rt using various solvents


Table 2: Optimization of temperature and amount of DABCO in the model reaction under solvent benzene

Entry	Catalyst amount (mol%)	Temp(°C)	Time(min)	Yield(%)
1	10	45	55	50
2	10	40	45	58
3	15	65	40	68
4	15	60	45	71
5	5.7	50	45	61
6	8	55	40	76


Naphthols



Aldehydes




Amines



Diversity of reactants used

Fig 2:



Scheme 3: Synthesis of Betti bases

Table 3 : Synthesis of arylaminonaphthols using DABCO under solvent benzene

Entry	Reactants	Products	Time(min)	Yield(%)
1	2a/13a/14a	15a	40	80
2	2a/13b/14a	15b	45	88
3	2a/13c/14a	15c	45	85
4	2a/13d/14a	15d	40	82
5	2a/13e/14a	15e	45	85
6	2a/13f/14a	15f	35	80
7	2a/13g/14a	15g	40	81
8	2a/13h/14a	15h	45	87
9	2a/13a/14a	15i	40	85
10	2a/13a/14c	15j	45	75
11	2a/13a/14d	15k	38	88
12	2a/13a/14e	15l	40	85
13	2a/13a/14f	15m	35	87
14	2a/13d/14b	15n	45	86
15	2a/13e/14c	15o	40	85
16	2a/13d/14e	15p	45	88

Reaction conditions 2-naphthol/aldehyde/amines=1:1:1.15 and DABCO (8mol%) at 55°C under solvent benzene.

A. General Procedure for the synThesis of Aminonaphthols

A mixture of 2- naphthol(1mmol), aldehyde(1mmol) and an amine(1.15 mmol) was treated with DABCO Catalytic additive. The mixture was stirred at 55° C under solvent benzene in a magnetic stirrer for an appropriate duration. The progress of the reaction was monitored by TLC(ethyl acetate/hexane).Upon completion of the reaction, the mixture was allowed to cool at room temperature. The solids were collected through the suction filtration process. Crude products washed with ether, dried in vacuum, recrystallized with MeOH/EtOH/Isopropanol/Acetone whichever found suitable. All compounds were characterized using the following analytical techniques like melting point(mp),IR, ¹H NMR, ¹³C NMR and elemental (C,H,N)analysis.

1-[Phenyl(phenylamino)methyl]naphthalen-2-ol (15a)

White solid ; Yield 80% ; mp : 130-135° C ; FT-IR(KBr) ν 3517,3403,3156, 3006,1705,1591,1328,857 cm⁻¹; ¹H NMR (400MHz, CDCl₃) : δ 4.21(1H, bs, N-H, disappeared in the presence of D₂O), 6.23 (1H, s, methine-H), 6.86 (2H, d, J=7.8 Hz, Ar-H) ν 7.22-7.53 (10H, m, Ar-H), 7.86-7.92 (3H, m, Ar-H), 11.64 (1H, bs, OH, disappeared in the presence of D₂O)ppm; ¹³C NMR (100 MHz, CDCl₃): δ 62.4, 115.4, 117.5, 118.4, 121.8,121.9, 122.9, 125.8, 125.9, 127.2, 127.7, 127.9, 128.3, 128.5, 128.7, 128.9, 139.9, 145.7, 157.8ppm. Anal. Calcd. For C₂₃H₁₉NO: C, 84.87 ; H, 5.88; N,4.30 ;Found :C, 84.80; H,5.80; N,4.27%

1-[(2-Chlorophenyl)(phenylamino)methyl]naphthalen-2-ol (15b)

Colourless crystals; Yield 88% ; mp: 155-158 °C; FT-IR (KBr) ν 3518, 3401, 3159, 3006, 1704, 1590, 1329, 854 cm⁻¹; ¹H NMR (400 MHz, CDCl₃): δ 4.11 (1H, bs, N-H), 6.67 (1H, s, methine-H), 6.90 (2H, d, J=8Hz, Ar-H), 7.03 (1H, t, J=7.6Hz, Ar-H), 7.15-7.28 (5H, m, Ar-H), 7.33-7.48 (3H, m, Ar-H), 7.87-7.99 (2H, m, Ar-H), 11.74 (1H, bs, OH)ppm; ¹³C NMR (100MHz, CDCl₃): δ 58.6, 111.9, 115.7, 118.9, 120.5, 121.4, 121.9, 126.5, 126.8, 127.9, 128.6, 128.9, 129.5, 129.7, 130.7, 132.9, 136.5, 145.11, 156.2 ppm. Anal. Calcd. For C₂₃H₁₈ClNO: C, 76.77 ; H, 5.04; N,3.89 ;Found :C, 76.58; H,4.98; N,3.86%

1-[(3-Chlorophenyl)(phenylamino)methyl]naphthalene-2-ol (15c)

Colourless Crystals; Yield 85% ; mp: 148-150° C ; FT-IR (KBr) ν 3388, 3347, 3064, 2905, 1604, 1498, 1241, 764 cm⁻¹; ¹H NMR (400MHz, CDCl₃): δ 4.14 (1H, bs, N-H), 6.14 (1H, s, methane-H),6.84 (2H, d, J=8.2Hz, Ar-H), 6.95 (1H, t, J=7.5Hz, Ar-H), 7.15-7.22 (4H, m, Ar-H), 7.41 (1H, t, J=7.5Hz, Ar-H), 7.47-7.53 (3H, m, Ar-H), 7.74 (1H, s, Ar-H), 7.83-7.89 (3H, m, Ar-H), 11.46 (1H, bs, OH) ppm; ¹³C NMR (100MHz, CDCl₃): δ 60.9, 112.3, 115.5, 119.2,120.2, 120.9, 121.9, 122.4, 125.6, 125.9, 128.1, 128.6, 129.3, 129.9, 130.5, 130.8, 142.2, 145.5, 155.5 ppm. Anal. Calcd. For C₂₃H₁₈ClNO: C,76.77; H,5.04; N,3.89; Found: C,76.58; H,4.98; N,3.86%

1-[(4-Chlorophenyl)(phenylamino)methyl]naphthalen-2-ol (15d)

Colourless Crystals; Yield 82%; mp: 123-125° C; FT-IR (KBr) ν 3422, 3358, 3064, 2965, 1624, 1599, 1491, 1240, 764 cm^{-1} ; ^1H NMR (400MHz, CDCl_3): δ 4.14 (1H, bs, N-H), 6.21 (1H, s, methane-H), 6.84 (1H, d, $J=8.2\text{Hz}$, Ar-H), 6.90-6.98 (1H, m, Ar-H), 7.10 (4H, t, $J=8\text{Hz}$, Ar-H), 7.21-7.29 (3H, m, Ar-H), 7.37-7.54 (7H, m, Ar-H), 7.83-7.89 (2H, m, Ar-H), 11.46 (1H, bs, OH)ppm; ^{13}C NMR (100MHz, CDCl_3): δ 60.11, 112.3, 115.5, 119.2, 121.8, 122.9, 123.9, 124.4, 125.6, 127.5, 127.9, 128.6, 128.8, 129.9, 132.7, 140.8, 145.5, 151.5, 155.5 ppm. Anal. Calcd. For $\text{C}_{23}\text{H}_{18}\text{ClNO}$: C,76.77; H,5.04; N,3.89; Found: C,76.48; H,4.98; N,3.81%




Fig 4 : Products 15a-15p

1-[(phenylamino)(p-tolyl)methyl]naphthalen-2-ol (15e)

Colourless Crystals; Yield 85% ; mp: 133-135° C; FT-IR (KBr) ν 3422, 3352, 3064, 2975, 1604, 1599, 1491, 1240 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.39 (3H, s, $-\text{CH}_3$), 4.21 (1H, bs, N-H), 6.14 (1H, s, methane-H), 6.78 (2H, d, $J=8.2\text{Hz}$, Ar-H), 6.99 (1H, t, $J=7.8\text{Hz}$, Ar-H), 7.13-7.19 (5H, m, Ar-H), 7.27-7.38 (4H, m, Ar-H), 7.83-7.89 (3H, m, Ar-H), 11.66 (1H, bs, OH) ppm; ^{13}C NMR (100MHz, CDCl_3): δ 20.12, 61.5, 112.9, 115.4, 118.9, 120.7, 120.9, 121.9, 125.8, 126.8, 127.9, 128.0, 128.4, 128.6, 128.9, 130.7, 137.8, 145.8, 155.5 ppm. Anal. Calcd. For $\text{C}_{24}\text{H}_{21}\text{NO}$: C,84.92; H,6.24; N,4.13; Found: C,86.68; H,6.21; N,4.09%

1-[(3,4-Dimethoxyphenyl)(phenylamino)methyl]naphthalen-2-ol (15f)

Light brown solid; Yield 80% ; mp: 140-142° C ; FT-IR (KBr) ν 3399, 3318, 3019, 2965, 1599, 1511, 1240, 1028 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 3.89 (3H, s, $-\text{OCH}_3$), 3.85 (3H, s, $-\text{OCH}_3$), 4.21 (1H, bs, N-H), 6.14 (1H, s, methine-H), 6.85-6.89 (3H, m, Ar-H), 6.92 (1H, t, $J=7.5\text{Hz}$, Ar-H), 6.99-7.04 (2H, m, Ar-H), 7.13-7.18 (3H, m, Ar-H), 7.31 (1H, t, $J=7.5\text{Hz}$, Ar-H) 7.41 (1H, m, Ar-H), 7.75-7.85 (3H, m, Ar-H), 11.58 (1H, bs, OH) ppm; ^{13}C NMR (100MHz, CDCl_3): δ 54.11, 61.8, 109.8, 110.6, 113.4, 115.5, 118.9, 119.5, 120.4, 120.9, 121.8, 125.7, 127.9, 128.2, 128.6, 128.8, 130.8, 132.7, 145.8, 148.5, 148.8, 155.5 ppm. Anal. Calcd. For $\text{C}_{25}\text{H}_{23}\text{NO}_3$: C,77.90; H,6.01; N,3.63; Found: C,77.68; H,5.98; N,3.58%

1-[(2-Bromophenyl)(phenylamino)methyl]naphthalen-2-ol (15g)

Light brown solid; Yield 81% ; mp: 152-154° C ; FT-IR (KBr) ν 3429, 3348, 3059, 3019, 2970, 1624, 1604, 1499, 1240, 698 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 4.08 (1H, bs, N-H), 6.48 (1H, s, methine-H), 6.81 (2H, d, $J=8.0\text{Hz}$, Ar-H), 6.94 (1H, t, $J=8.1\text{Hz}$, Ar-H), 7.15-7.18 (6H, m, Ar-H), 7.29-7.38 (2H, t, m, Ar-H), 7.60 (1H, d, $J=8.5\text{Hz}$, Ar-H), 7.67-7.69 (1H, m, Ar-H), 7.79 (2H, d, $J=8.8\text{Hz}$, Ar-H), 11.68 (1H, bs, OH) ppm; ^{13}C NMR (100 MHz, CDCl_3): δ 60.8, 111.9, 115.6, 118.9, 120.5, 121.5, 121.9, 123.6, 126.2, 127.5, 127.5, 127.9, 128.2, 128.6, 129.5, 129.8, 130.5, 132.7, 138.2, 145.9, 155.9 ppm. Anal. Calcd. For $\text{C}_{23}\text{H}_{18}\text{BrNO}$: C,68.33; H,4.49; N,3.46; Found: C,68.08; H,4.45; N,3.38%

1-[(4-Methoxyphenyl)(phenylamino)methyl]naphthalen-2-ol (15h)

Light brown solid; Yield 87% ; mp: 272° C ; FT-IR (KBr) ν 3414, 3338, 3009, 2961, 1621, 1511, 1243, 822 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 3.69 (3H, s, $-\text{OCH}_3$), 4.14 (1H, bs, N-H), 6.14 (1H, s, methine-H), 6.75 (2H, d, $J=8.4\text{Hz}$, Ar-H), 6.82 (2H, d, $J=8.5\text{Hz}$, Ar-H), 6.98 (1H, t, $J=7.5\text{Hz}$, Ar-H), 7.13-7.18 (3H, m, Ar-H), 7.28-7.37 (4H, m, Ar-H), 7.75-7.85 (3H, m, Ar-H), 11.58 (1H, bs, OH) ppm; ^{13}C NMR (100 MHz, CDCl_3): δ 54.3, 61.4, 113.2, 113.6, 115.4, 118.9, 120.5, 120.8, 121.9, 125.8, 127.7, 128.1, 128.3, 128.6, 128.8, 130.5, 132.4, 145.8, 154.9, 158.8 ppm. Anal. Calcd. For $\text{C}_{24}\text{H}_{21}\text{NO}_2$: C,81.10; H,5.96; N,3.94; Found: C,80.86; H,5.94; N,3.90%

1-[(3-Chlorophenylamino)(phenyl)methyl]naphthalen-2-ol (15i)

Colourless Crystals; Yield 85% ; mp: 114-116° C ; FT-IR (KBr) ν 3375, 3338, 3069, 2940, 1597, 1479, 1240, 758 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 4.18 (1H, bs, N-H), 6.18 (1H, s, methine), 6.64 (1H, dd, $J=8.0\text{Hz}$, Ar-H), 7.06 (1H, t, $J=8.1\text{Hz}$, Ar-H), 7.15 (1H, d, $J=8.8\text{Hz}$, Ar-H), 7.31-7.35 (5H, m, Ar-H), 7.48 (2H, d, $J=7.1\text{Hz}$, Ar-H), 7.77-7.81 (3H, m, Ar-H), 10.89 (1H, bs, OH) ppm; ^{13}C NMR (100 MHz, CDCl_3): δ 61.8, 112.5, 113.6, 115.9, 118.9, 120.5, 120.8, 122.4, 125.9, 126.9, 127.5, 128.2, 128.3, 128.5, 129.2, 129.57, 129.7, 129.7, 130.7, 135.2, 139.6, 146.9, 154.9 ppm. Anal. Calcd. For $\text{C}_{23}\text{H}_{18}\text{ClNO}$: C,76.77; H,5.04; N,3.89; Found: C,76.54; H,5.02; N,3.84%

1-[(4-Methoxyphenylamino)(phenyl)methyl]naphthalen-2-ol (15j)

Light brown solid; Yield 75% ; mp: 119-122° C ; FT-IR (KBr) ν 3398, 3328, 3027, 2931, 1623, 1511, 1238, 822 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 3.71 (3H, s, $-\text{OCH}_3$), 4.24 (1H, bs, N-H), 6.14 (1H, s, methine-H), 6.71-6.77 (4H, m, Ar-H), 7.18 (1H, d, $J=8.8\text{Hz}$, Ar-H), 7.29-7.40 (5H, m, Ar-H), 7.50 (2H, d, $J=7.5\text{Hz}$, Ar-H), 7.75-7.85 (3H, m, Ar-H), 11.98 (1H, bs, OH) ppm; ^{13}C NMR (100 MHz, CDCl_3): δ 54.6, 62.6, 112.8, 113.9, 116.9, 119.2, 120.5, 121.8, 125.8, 126.9, 127.9, 128.1, 128.3, 128.8, 130.6, 139.4, 140.8, 153.9, 155.8 ppm. Anal. Calcd. For $\text{C}_{24}\text{H}_{21}\text{NO}_2$: C,81.10; H,5.96; N,3.94; Found: C,80.92; H,5.89; N,3.90%

1-[(4-Nitrophenylamino)(phenyl)methyl]naphthalen-2-ol (15k)

Yellow solid; Yield 88% ; mp: 154-156° C ; FT-IR (KBr) ν 3396, 3358, 3029, 2955, 1598, 1509, 1309, 1113, 768 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 5.08 (1H, bs, N-H, disappeared in presence of D_2O), 6.42 (1H, s, methine-H), 6.72 (2H, d, $J=8.8\text{Hz}$, Ar-H), 7.11 (1H, d, $J=8.8\text{Hz}$, Ar-H), 7.29-7.37 (4H, m, Ar-H), 7.42-7.45 (3H, m, Ar-H), 7.76-7.86 (3H, m, Ar-H), 8.01 (2H, d, $J=8.8\text{Hz}$, Ar-H), 11.67 (1H, bs, OH, disappeared in the presence of D_2O) ppm; ^{13}C NMR (100 MHz, CDCl_3): δ 58.1, 113.1, 113.6, 118.4, 120.6, 122.5, 125.0, 126.3, 126.5, 127.5, 128.2, 128.5, 128.6, 129.7, 130.7, 138.9, 139.7, 151.7, 153.2 ppm. Anal. Calcd. For $\text{C}_{23}\text{H}_{18}\text{N}_2\text{O}_3$: C,74.58; H,4.90; N,7.56; Found: C,74.34; H,4.85; N,7.48%

1-[(4-Fluorophenylamino)(phenyl)methyl]naphthalen-2-ol (15l)

White solid ; Yield 85% ; mp: 111-112°C ; FT-IR (KBr) ν 3397, 3338, 3023, 2970, 1607, 1505, 1230, 816, 753 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 4.15 (1H, bs, N-H), 6.12 (1H, s, methane-H), 6.69-6.73 (2H, m, Ar-H), 6.80-6.86 (2H, m, Ar-H), 7.46-7.47 (2H, m, Ar-H), 7.71-7.76 (3H, m, Ar-H), 11.69 (1H, bs, OH) ppm; ^{13}C NMR (100 MHz, CDCl_3): δ 61.8, 112.5, 114.6, 115.9, 118.9, 120.5, 120.8, 121.8, 125.9, 126.9, 127.2, 127.4, 127.9, 128.1, 128.4, 128.9, 129.2, 130.7, 137.4, 145.2, 155.6, 157.1, 159.6 ppm. Anal. Calcd. For $\text{C}_{23}\text{H}_{18}\text{FNO}$: C, 80.45; H, 5.28; N, 4.08; Found: C, 80.25; H, 5.20; N, 4.04%

1-[(3,4-Dichlorophenylamino)(phenyl)methyl]naphthalen-2-ol (15m)

Brown solid; Yield 87% ; mp: 124-126°C ; FT-IR (KBr) ν 3403, 3353, 3062, 2954, 1597, 1379, 1040, 748 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 4.28 (1H, bs, N-H), 6.14 (1H, s, methane-H), 6.54 (1H, dd, $J=8.8\text{Hz}$, $J=2.4\text{Hz}$, Ar-H), 6.81 (1H, d, $J=2.4\text{Hz}$, Ar-H), 7.09-7.14 (2H, m, Ar-H), 7.30-7.33 (4H, m, Ar-H), 7.35-7.44 (3H, m, Ar-H), 7.72-7.77 (3H, m, Ar-H), 10.55 (1H, bs, OH) ppm; ^{13}C NMR (100 MHz, CDCl_3): δ 60.8, 114.6, 116.9, 118.9, 120.8, 122.1, 122.4, 125.9, 126.8, 126.9, 127.7, 128.1, 128.2, 128.5, 129.2, 129.7, 130.7, 131.2, 139.2, 145.1, 154.9 ppm. Anal. Calcd. For $\text{C}_{23}\text{H}_{17}\text{Cl}_2\text{NO}$: C, 70.06; H, 4.35; N, 3.55; Found: C, 69.86; H, 4.29; N, 3.52%

1-[(4-Chlorophenyl)(3-chlorophenylamino)methyl]naphthalen-2-ol (15n)

Brown solid; Yield 86%; mp: 123-124°C; FT-IR (KBr) ν 3398, 3346, 3063, 2965, 1598, 1485, 751 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 4.12 (1H, bs, N-H), 6.08 (1H, s, methine-H), 6.74 (2H, t, $J=8.5\text{Hz}$, Ar-H), 7.05-7.09 (2H, m, Ar-H), 7.22 (2H, d, $J=8.4\text{Hz}$, Ar-H), 7.32-7.36 (3H, m, Ar-H), 7.42 (2H, d, $J=8.5\text{Hz}$, Ar-H), 7.32-7.36 (3H, m, Ar-H), 7.42 (2H, d, $J=8.5\text{Hz}$, Ar-H), 7.89 (3H, d, $J=8.9\text{Hz}$, Ar-H), 11.52 (1H, bs, OH) ppm; ^{13}C NMR (100 MHz, CDCl_3): δ 63.4, 111.6, 114.3, 114.6, 118.0, 121.9, 122.9, 126.0, 126.9, 127.3, 123.4, 127.8, 128.5, 128.6, 128.9, 129.5, 132.6, 132.8, 134.4, 140.1, 151.8, 155.0, 157.4, 159.9 ppm. Anal. Calcd. For $\text{C}_{23}\text{H}_{17}\text{Cl}_2\text{NO}$: C, 70.06; H, 4.35; N, 3.55; Found: C, 69.83; H, 4.30; N, 3.49%

1-[(4-Methoxyphenylamino)(p-tolyl)methyl] naphthalen-2-ol (15o)

Brown solid ; Yield 85%; mp: 115-117°C; FT-IR (KBr) ν 3401, 3339, 3013, 2959, 1605, 1509, 1252, 754 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 2.40 (3H, s, -CH₃), 3.83 (3H, s, -OCH₃), 4.11 (1H, bs, N-H), 6.08 (1H, s, methine-H), 6.69-6.76 (2H, m, Ar-H), 6.93 (2H, d, $J=8.9\text{Hz}$, Ar-H), 7.15 (2H, d, $J=8\text{Hz}$, Ar-H), 7.25-7.28 (3H, m, Ar-H), 7.36 (2H, d, $J=8.0\text{Hz}$, Ar-H), 7.75-7.79 (3H, m, Ar-H), 11.67 (1H, bs, OH) ppm; ^{13}C NMR (100 MHz, CDCl_3): δ 19.7, 64.4, 83.5, 112.4, 115.4, 118.2, 122.2, 122.5, 124.4, 125.7, 126.9, 127.4, 128.2, 128.3, 128.4, 131.4, 132.1, 136.4, 141.9, 143.5, 151.9 ppm. Anal. Calcd. For $\text{C}_{25}\text{H}_{23}\text{NO}_2$: C, 81.27; H, 6.27; N, 3.79; Found: C, 81.09; H, 6.25; N, 3.74%

1-[(4-Chlorophenyl)(4-fluorophenylamino)methyl] naphthalen-2-ol (15p)

Colourless crystals; Yield 88%; mp: 164-165°C; FT-IR (KBr) ν 3432, 3355, 3061, 2961, 1624, 1506, 1221, 816, 772 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3): δ 4.12 (1H, bs, N-H), 6.08 (1H, s, methine-H), 6.74 (2H, t, $J=8.4\text{Hz}$, Ar-H), 7.04-7.08 (2H, m, Ar-H), 7.21 (2H, d, $J=8.4\text{Hz}$, Ar-H), 7.32-7.36 (3H, m, Ar-H), 7.42 (2H, d, $J=8.4\text{Hz}$), 7.88 (3H, d, $J=8.8\text{Hz}$, Ar-H), 11.52 (1H, bs, OH) ppm; ^{13}C NMR (100 MHz, CDCl_3): δ 63.4, 111.6, 114.3, 114.5, 117.9, 121.8, 129.5, 132.5, 132.8, 134.4, 140.0, 151.7, 155.0, 157.4, 159.9 ppm. Anal. Calcd. For $\text{C}_{23}\text{H}_{17}\text{ClFNO}$: C, 73.11; H, 4.54; N, 3.71; Found: C, 72.94; H, 4.52; N, 3.69%

IV. CONCLUSION

In conclusion, we have developed a high yielding protocol for the synthesis of aminocatalyst. This method operates efficiently under solvent benzene and demonstrates excellent substrate versatility, providing a valuable approach for the synthesis of aminonaphthol derivatives in organic chemistry. A novel series of arylaminonaphthols was synthesized through a one-pot three component reaction involving an aldehyde, an aromatic amine, and 2-naphthol, facilitated by DABCO under solvent benzene. This efficient methodology allows for the straight forward assembly of the desired compounds, showcasing the effectiveness of DABCO as an organocatalyst and highlighting the potential for practical applications in organic synthesis.

REFERENCES

- [1] Knapp S (1995) Synthesis of complex nucleoside antibiotics. *Chem Rev* 95:1859-1876. doi:10.1021/cr00038a006.
- [2] Mannhold R, Kubinyi H (2006) Nucleic acid drugs. In: Dingermann T, Steinhilber D, Folkers G (eds) *Molecular biology in medicinal chemistry*. Wiley, Weinheim. doi: 10.1002/3527602666.

- [3] Wang YK, Izawa T, Kobayashi S, Ohno M (1982) Stereocontrolled synthesis of (+)- negamycin from an acyclic homoallylamine by 1,3-asymmetric induction. *J Am Chem Soc* 104:6465-6466. doi:10.1021/ja00387a060.
- [4] Cardellicchio C, Capozzi MAM, Naso F (2010) The Betti bases: the awakening of a sleeping beauty. *Tetrahedron* 21:507-517. doi: 10.1016/j.tetasy.2010.03.020.
- [5] Elmendorf HG, Walls CD, Wolf C (2012) Compositions and methods for the treatment of giardiasis. US Patent 20, 120, 283,267.
- [6] Gerlach M, Maul C (2007) Substituted 1 and 2- naphthol Mannich bases. US Patent 7, 202,242B2.
- [7] Gyemant N, Engi H, Schelz Z, Szatmari I, Toth D, Fulop F, Molnar J, de Witte P (2010) In vitro and in vivo multidrug resistance reversal activity by a Betti-base derivative of tylosin. *Br J Cancer* 103:178-185. doi:10.1038/sj.bjc.6605716.
- [8] Shen AY, Tsai CT, Chen CL (1999) Synthesis and cardiovascular evaluation of N-substituted 1-aminomethyl-2-naphthols. *Eur J Med Chem* 34:877-882. doi:10.1016/S0223-5234(99)00204-4.
- [9] Szatmari I, Fulop F (2013) Synthesis, transformations and applications of aminonaphthol derivatives prepared via modified Mannich reactions. *Tetrahedron* 69:1255-1278. doi:10.1016/j.tet.2012.11.055.
- [10] Salama TA (2013) Silicon - induced general, mild, and efficient one pot, three- component synthesis of amidoalkyl naphthol libraries. *Synlett* 24:713-718. doi:10.1055/s-0032-1318392
- [11] Safari J, Zarnegar Z (2013) A magnetic nanoparticle-supported sulfuric acid as a highly efficient and reusable catalyst for rapid Synthesis of amidoalkyl naphthols. *J Mol Catal A* 379:269-276. doi:10.1016/j.molcata.2013.08.028.
- [12] Rostamizadeh S, Abdollahi F, Shadjou N, Amani AM (2013) MCM- 41-SO₃H: a novel reusable nanocatalyst for synthesis of amidoalkyl naphthols under solvent - free conditions. *Monatsh Chem* 144:1191-1196. doi: 10.1007/s00706-013-0936-4.
- [13] Lei ZK, Xiao L, Lu XQ, Huang H, Liu CJ (2013) Graphite supported perchloric acid (HClO₄- C): an efficient and recyclable heterogeneous catalyst for the one - pot synthesis of amidoalkyl naphthols. *Molecules* 18:1653-1659. doi:10.3390/ molecules18021653.
- [14] Das VK, Borah M, Thakur AJ (2013) Piper-bottle-shaped nano- S-catalyzed synthesis of 1-amidoalkyl-2-naphthols under solvent -free reaction condition: A greener " Nanoparticle-Catalyzed Organic Synthesis Enhancement" approach. *J Org Chem* 78:3361-3366. doi:10.1021/jo302682k.
- [15] Csutortoki R, Szatmari I, Fulop F (2013) Syntheses of amidocarbamido- and carbamatoalkyl naphthols. *Curr Org Synth* 10:564-583.
- [16] Mulla SA, Salama TA, Pathan MY, Inamdar SM, Chavan SS (2012) Solvent-free, highly efficient one-pot multi-component synthesis of 1-amido and 1-carbamato-alkyl naphthols/phenols catalyzed by ethylammonium nitrate as reusable ionic liquid under neat reaction condition at ambient temperature. *Tetrahedron Lett* 54:672-675. doi:10.1016/j.tetlet.2012.12.004.
- [17] Kotadia DA, Soni SS (2012) Silica gel supported - SO₃H functionalised benzimidazolium based ionic liquid as a mild and effective catalyst for rapid Synthesis of 1-amidoalkyl-2-naphthols. *J Mol Catal A* 353:44-49. doi:10.1016/j.molcata.2011.11.003.
- [18] Zolfigol MA, Khazaei A, Moosavi - Zare AR, Zare A, Khakyzadeh V (2011) Rapid synthesis of 1-amidoalkyl-2-naphthols over sulfonic acid functionalized imidazolium salts. *Appl Catal A* 400: 70-81. doi:10.1016/j.apcata.2011.04.013.
- [19] Tamaddon F, Bistgani JM (2011) [MeC(OH)₂]⁺ClO₄⁻: a new efficient organocatalyst for the preparation of 1- amido- and 1-carbamato- alkyl naphthols. *Synlett* 2011:2947-2950. doi:10.1055/s-0031-1289906.
- [20] Mistry SR, Joshi RS, Maheria KC (2011) Zeolite H- BEA catalysed multicomponent reaction: one-pot synthesis of amidoalkyl naphthols -biologically active drug - like molecules. *J Chem Sci* 123:427-432. doi:10.1007/s12039-011-0095-2.
- [21] Shaterian HR, Yarahmadi H, Ghashang M (2008) Silica supported perchloric acid (HClO₄-SiO₂): an efficient and recyclable heterogeneous catalyst for the one - pot synthesis of . amidoalkyl-2-naphthols *Tetrahedron* 64:1263-1269. doi:10.1016/j.tet.2007.11.070.
- [22] Shaterian HR, Yarahmadi H, Ghashang M (2008) An efficient, simple and expedition synthesis of 1-amidoalkyl-2-naphthols as "drug like" molecules for biological screening. *Bioorg Med Chem Lett* 18:788-793. doi: 10.1016/j.tet.2007.11.070.
- [23] Karmakar B, Banerji J (2011) A competent pot and atom- efficient synthesis of Betti bases over nanocrystalline MgO involving a modified Mannich type reaction. *Tetrahedron Lett* 52:4957-4960. doi:10.1016/j.tetlet.2011.07.075.
- [24] Periasamy M, Anwar S, Reddy MN (2009) Simple and convenient methods for synthesis, resolution and application of aminonaphthols. *Indian J Chem B* 48: 1261- 1273.
- [25] Ghandi M, Olyaei A, Raoufmoghaddam S (2008) One - pot , three-component uncatalysed quantitative Synthesis of new aminonaphthols. *Indian J Chem B* 48: 1261-1273.
- [26] Oyaei A, Raoufmoghaddam S, Sadeghpour M, Ebadzadeh B (2010) Convenient and efficient method for the synthesis of N-heteroaryl aminonaphthols under solvent-free conditions. *Chin J Chem* 28:825-832. doi:10.1002/ cjoc.201090153.
- [27] Hosseinian A, Shaterian HR (2012) NaHSO₄. H₂O catalyzed multicomponent synthesis of 1-(benzothiazolylamino)methyl-2-naphthols under solvent - free conditions. *Phosphorus Sulfur Silicon* 187:1056-1063. doi:10.1080/10426507.2012.664221.
- [28] Buckley BR, Neary SP (2010) Organocatalysis. *Annu Rep Prog Chem Sect B Org Chem* 106:120-135. doi:10.1039/B927086H.
- [29] Grondal C, Jeanty M, Enders D (2010) Organocatalytic cascade reactions as a new tool in total synthesis. *Nat Chem* 2:167-178. doi:10.1038/nchem.539.
- [30] Marcelli T, van Maarseveen JH, Hiemstra H (2006) Cupreines and cupreidines: an emerging class of bifunctional cinchona organocatalysts. *Angew Chem Int Ed* 45:7496-7504. doi: 10.1002/anie. 200602318.
- [31] Liu G, Zhang S, Li H, Zhang T, Wang W (2011) Organocatalytic enantioselective Friedel-Crafts reactions of 1-naphthols with aldimines. *Org Lett* 13:838-831. doi:10.1021/ol02987n.
- [32] Ramachary DB, Babul Reddy G (2007) A new organocatalyst for Friedel-Crafts alkylation of 2- naphthols with isatins: application of an organo- click strategy for the cascade synthesis of highly functionalized molecules. *Tetrahedron Lett* 48: 7618-7623. doi:10.1016/j.tetlet.2007.08.129.
- [33] Shahriza A, Safa KD, Esmati S (2014) Synthesis, spectroscopic and DFT studies of novel fluorescent dyes: 3-aminoimidazo[1,2-a] pyridines processing 4-pyrone moieties. *Spectrochim Acta A* 117:614-621. doi:10.1016/j.saa.2013.09.056.
- [34] Shahriza A, Esmati S, Miri R, Firuzi I, Edraki N, Nejati M (2013) Cytotoxic activity assessment, QSAR and docking study of novel bis- carboxamide derivatives of 4- pyrones synthesized by Ugi four- component reaction. *Eur J Med Chem* 66:388-399. doi:10.1016/j.ejmech.2013.05.045.
- [35] Shahriza A, Esmati S (2013) Three novel sequential reactions for the facile synthesis of library of bisheterocycles possessing the 3-aminoimidazo[1,2-a] pyridine core catalysed by bismuth (III) chloride. *Synlett* 24:595-602. doi:10.1055/s-0032-1318221.



- [36] ShahrissaA,MiriR,Esmati S, Saraei M, MehdipourAR,Sharifi M(2012) Synthesis and calcium channel antagonist activity of novel 1,4-dihydropyridine derivatives possessing 4-pyrone moieties. Med Chem Res 21:284-292.doi:10.1007/s00044-010-9534-8.
- [37] ShahrissaA, Esmati S, Nazari MG(2012) Boric acid as a mild and efficient catalyst for one - pot synthesis of 1-amidoalkyl-2-naphthols under solvent -free conditions. J Chem Sci 124:927-931.doi:10.1007/s12039-012-0285-6.
- [38] Dindulkar SD, Puranik VG,Jeong YT(2012) Supported copper triflate as an efficient catalytic system for the synthesis of highly functionalized 2-naphthol Mannich bases under solvent free condition. Tetrahedron Lett 53:4376-4380.doi:10.1016/j.tetlet.2012.06/022.
- [39] Ganesan SS, Rajendran N, Sundarakumar SI, Ganesan A, PemiahB(2013) 2-naphthol in glycerol: a versatile pair for efficient and convenient synthesis of aminonaphthols, naphtho-1,3-oxazines,and benzoxanthene. Synthesis 45:1564-1568.doi:10.1055/s-0033-1338430.
- [40] Saidi MR, Azizi N, Naimi- Jamal MR(2001) Lithium perchlorate assisted one- pot three-component aminoalkylation of electron- rich aromatic compounds. Tetrahedron Lett 43:8111-8113.doi:10.1016/S0040-4039(01)01732-4.
- [41] Katritzky AR, Abdel - Fattah AA, Tymoshenko DO, Belyakov SA, Ghiviriga I, Steel PJ(1999) Amino (hetero) arylmethylation of phenols with N- [alpha-amino(hetero) arylmethyl] benzotriazoles. J Org Chem 64:6071-6075.doi: 10.1021/jo9903609.
- [42] Kumar A, Gupta MK, Kumar M(2010) Non - ionic surfactant catalyzed synthesis of Betti bases in water . Tetrahedron Lett 51:1582-1584. doi:10.1016/j.tetlet.2010.01.056.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)