



IJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 14 **Issue:** V **Month of publication:** May 2026

DOI: <https://doi.org/10.22214/ijraset.2026.82604>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Potential of Napier Grass for Biofertilizer Production: A Comprehensive Review

Khushi Undhad, Indar Kumar, Abre Alam, Dr. Reena Sheokand
BPT(A.H.C.S.) B.Sc.(Hons.) Agriculture Vivekananda Global University, Jaipur

Abstract: *This review looks into Napier grass's potential as a sustainable feedstock for the manufacturing of biofertilizer. The hunt for environmentally suitable substitutes has quickened due to growing environmental concerns related to the overuse of chemical fertilisers. Napier grass is a viable resource for biomass valorisation because of its quick growth, high biomass yield, and capacity to respond to a variety of agroclimatic situations. The literature on its biochemical makeup, breakdown processes (such as composting, fermentation, and anaerobic digestion), and function in improving soil fertility and plant growth is compiled in this study. Agronomic, environmental, and financial advantages of Napier-based biofertilizers are also covered in the review. Despite its potential, there is little research on its direct use as a feedstock for biofertilizer, which highlights important knowledge gaps in large-scale implementation, process optimisation, and nutrient characterisation. According to the study's findings, Napier grass has a lot of potential for sustainable agriculture, but more investigation is needed to validate field-scale uses and standardise production techniques.*

KEYWORDS: *Napier grass, Biofertilizer, Biomass valorization, Sustainable agriculture, Composting, Nutrient cycling*

I. INTRODUCTION

The usage of chemical fertilisers has significantly increased as a result of contemporary agriculture's intensification in order to meet the world's expanding food demand. Although these inputs have increased crop output, their overuse and uneven application have had a number of negative effects. Soil degradation, nitrogen imbalance, a decrease in soil organic matter, and a decrease in beneficial microbial activity have all been linked to continuous use of synthetic fertilisers [1,2]. Furthermore, eutrophication and groundwater contamination are two examples of environmental damage caused by chemical fertiliser leaking into water bodies [1]. These difficulties show how urgently better environmentally friendly and sustainable methods of managing nutrients are needed. Sustainable farming methods that reduce environmental impact while preserving soil fertility and crop output have gained popularity in response to these worries. Biofertilizers have become a viable substitute among these methods. Biofertilizers are organic formulations with decomposed organic matter or living microorganisms that improve soil structure, increase nutrient availability, and naturally promote plant development [2, 3]. By boosting microbial diversity, improving nutrient cycling, and promoting soil physical characteristics like water retention and aeration, biofertilizers—as opposed to chemical fertilizers—promote long-term soil health [4].

The creation of efficient biofertilizers depends critically on the choice of suitable biomass resources. Napier grass has drawn interest as a possible feedstock in this regard. Napier grass is a perennial plant that grows quickly and is commonly grown as a fodder crop in tropical and subtropical areas. It is distinguished by its high biomass output, quick growth rate, and capacity to flourish in a variety of environmental circumstances, such as low-input systems and marginal soils [5,6]. Because of these qualities, it is a plentiful and sustainable supply of organic material that can be used in biomass-based applications.

From a biochemical standpoint, Napier grass has large levels of lignocellulosic components, such as cellulose, hemicellulose, and lignin, which can be broken down by biological processes to release vital nutrients [7]. Because of its high organic matter content, it can be converted into biofertilizer using processes including anaerobic digestion, fermentation, and composting [8]. Napier grass is also more suitable for decentralised and small-scale biofertilizer production systems due to its broad availability and low production cost [6].

Despite these benefits, compared to other organic leftovers, the use of Napier grass especially for the creation of biofertilizer is still comparatively unexplored. Research on its direct usage as a feedstock for biofertilizer is lacking, as the majority of current studies concentrate on its use in fodder production or bioenergy generation [9]. By offering a cheap, environmentally benign substitute for chemical fertilisers, realising its promise in this area might greatly aid sustainable agriculture.

II. LITERATURE REVIEW

According to earlier research, soil fertility, nitrogen cycling, and microbial activity are all enhanced by organic and biofertilizers. It has been shown that compost made from agricultural waste can improve soil organic carbon and water-holding capacity. Beneficial bacteria found in biofertilizers enhance root development, phosphorus solubilisation, and nitrogen fixation.

Due to its high biomass production and nutritional value for cattle, napier grass has been extensively researched as a fodder crop. It has also been studied more recently for bioenergy uses including biomass fuel and biogas. However, only a small number of research have particularly examined its application in the manufacturing of biofertilizer. Researchers have discovered that fermentation and composting may break down lignocellulosic biomass, like Napier grass, to release nutrients. This suggests that Napier grass could develop into a significant renewable feedstock.

A. Overview of Napier Grass :

Elephant grass, or napier grass (*Pennisetum purpureum*), is a perennial C4 tropical grass that is extensively grown due to its remarkable biomass yield and adaptability to a variety of agroclimatic conditions ^[5,6]. Due to its high output potential and adaptability to a variety of environmental circumstances, this native African plant has been widely transplanted into Asia and Latin America ^[7]. It is a promising option for biomass valorisation due to its quick growth, high lignocellulosic content, and nutrient composition, especially in sustainable agricultural inputs as biofertilizers ^[8,9]. Napier grass has emerged as a promising renewable resource due to the growing interest in lignocellulosic biomass for agricultural and environmental uses ^[10,11]. Its importance in sustainable biomass production systems is further increased by its capacity to grow on marginal lands with little input ^[12,13].

B. Botanical and Agronomic Characteristics :

Napier grass is tall, grows quickly, and has a wide root system. It is a member of the Poaceae family. It usually reaches heights of 2-4 meters and yields dense tillers, which contribute to substantial biomass accumulation ^[5,14].

1) Growth Rate

Due to its C4 photosynthetic pathway, which enables effective carbon fixation and greater productivity than C3 plants, napier grass grows quickly ^[15]. It can be harvested several times a year, and its quick regrowth upon cutting guarantees a steady supply of biomass ^[6,16]. Research has demonstrated that when appropriate fertiliser management is maintained, frequent harvesting does not substantially lower productivity ^[17].

2) Climate Adaptability

Napier grass's broad adaptability is one of its main benefits. It can withstand high temperatures, low soil fertility, and drought, and it grows well in tropical and subtropical regions ^[7,18]. Additionally, it may grow on marginal and degraded soil, lessening competition with food crops and promoting sustainable land use practices ^[12,19].

3) Biomass Production

Under ideal circumstances, napier grass is known for its extraordinarily high biomass yield, which normally ranges from 40 to 80 tonnes of dry matter per hectare yearly ^[6,21]. Even greater yields have been documented in some intensive farming systems ^[22]. Because of its high productivity, it is a scalable and economically viable feedstock for the production of biofertilizer, particularly in areas with an abundance of land resources ^[23].

C. Chemical Composition:

Napier grass's biochemical makeup, especially its lignocellulosic structure and nutrient content, has a significant impact on its potential as a feedstock for biofertilizer.

1) Components of Lignocellulosic Materials

Lignocellulosic biomass, which includes cellulose (30–45%), hemicellulose (20–30%), and lignin (10–20%), makes up the majority of napier grass ^[10,24]. During the fermentation and composting processes, these structural elements offer a rich carbon source for microbial breakdown. Microorganisms break down these polymers, releasing simpler chemicals and nutrients necessary for plant growth ^[25, 26].

However, lignin might slow down the rate of breakdown, making microbial augmentation or pretreatment necessary for effective conversion^[27]. Microbial consortia and improvements in biomass processing technologies have demonstrated promise for increasing decomposition efficiency^[28].

2) NPK (Nutrient Content)

Essential macronutrients for plant development, such as potassium (K), phosphorus (P), and nitrogen (N), are found in napier grass^[9,29]. Its organic matter helps release nutrients gradually during decomposition, even if the concentration of nutrients may vary based on growth stage, soil conditions, and management techniques^[30]. Napier-derived biofertilizers are more sustainable than synthetic fertilisers because of their slow-release feature, which improves nutrient use efficiency and lowers leaching losses^[2,31]. Its high organic carbon content also enhances nutrient cycling and soil microbial activity^[12, 32].

3) Section Perspective

Napier grass is an excellent feedstock for the manufacturing of biofertilizer due to its fast growth, high biomass yield, environmental flexibility, and advantageous biochemical composition. Its potential in sustainable agriculture systems is reinforced by the lignocellulosic structure that promotes microbial breakdown and the nutritional profile that enhances soil fertility.

D. Napier grass transformation into biofertilizer :

Lignocellulosic biomass is converted into biofertilizer by physicochemical and biological processes that change complicated organic materials into nutrient-rich forms that are easily absorbed by plants. Napier grass is a good substrate for a number of biofertilizer production processes, such as composting, fermentation, anaerobic digestion, and microbial enrichment, because of its high cellulose and organic matter content^[10,24,25]. These activities support sustainable soil fertility management, increase microbial activity, and improve nutrient mineralisation^[2,12].

1) Composting Aerobically

One of the most popular techniques for turning plant biomass into stable organic fertiliser is aerobic composting. It involves the microbial breakdown of organic materials in the presence of oxygen, producing a nutrient-rich substance that resembles humus^[8,14]. Because of its fibrous structure, napier grass needs to be chopped in order to increase surface area and promote microbial activity. Microorganisms like bacteria and fungi break down cellulose and hemicellulose into simpler compounds during composting, releasing nutrients like potassium, phosphate, and nitrogen^[25, 26].

However, Napier grass's comparatively high lignin content may slow down the process of breakdown. Co-composting with nitrogen-rich materials, like animal dung or green waste, is frequently advised to get around this restriction in order to balance the carbon-to-nitrogen (C:N) ratio and boost microbial activity . Compost quality and nutrient availability are greatly improved by optimal composting conditions, such as adequate aeration, moisture content (50–60%), and temperature management .An efficient biofertilizer for sustainable agriculture, compost made from lignocellulosic biomass has been shown to improve soil structure, raise organic carbon content, and boost microbial diversity^[12,16].

2) Anaerobic Digestion

Another crucial process for turning Napier grass into biofertilizer is anaerobic digestion. In this process, organic matter is broken down in the absence of oxygen to produce digestate, a nutrient-rich byproduct, and biogas (methane and carbon dioxide)^[9,26]. To increase biogas generation and decomposition efficiency, napier grass can be co-digested with animal manure or other organic wastes . Rich in vital nutrients, the digestate can be directly applied to soil as a liquid or semi-solid biofertilizer. Due to partial mineralisation during digestion, digestate has more easily accessible nutrients than raw biomass^[36].

Anaerobic digestion is a dual-benefit method that also helps with waste management and the creation of renewable energy. However, effective conversion of lignocellulosic materials like Napier grass depends on issues like feedstock pretreatment and process optimisation .

3) Liquid Biofertilizers Based on Fermentation

Microbial cultures are used in fermentation operations to transform biomass into liquid biofertilizers. Sustainable agriculture systems frequently employ traditional formulations like jeevamrut or effective microorganism (EM) solutions .

This method allows for controlled fermentation by combining chopped Napier grass with water and microbial inoculants (such as molasses or cow manure). Organic matter is broken down by microbial activity, which also releases soluble nutrients and growth-promoting compounds such as phytohormones, amino acids, and enzymes^[39].

Fermentation-produced liquid biofertilizers have a number of benefits.

Quick availability of nutrients

Simple application using irrigation systems

Increased soil microbial activity

Fermented organic inputs greatly enhance plant growth, root development, and nutrient uptake efficiency, according to studies^[23, 40].

4) *Bioaugmentation and Microbial Enrichment*

In order to improve nutrient availability and speed up biomass degradation, microbial enrichment entails adding advantageous microorganisms. Among these microbes are:

Bacteria that fix nitrogen

Microbes that solubilise phosphate

Fungi that break down

In order to break down complex lignocellulosic substances and transform them into nutrients that plants can use, these microbial consortia are essential.

Microbial inoculation can greatly increase the rate of breakdown and shorten processing times for Napier grass. Compost quality and nutrient content have been proven to improve with bioaugmentation techniques employing specialised microbial strains.

5) *Systems of Integrated Conversion*

To optimise nutrient recovery and efficiency, integrated systems integrate several processes, including vermicomposting, anaerobic digestion, and composting. For instance:

Digestate is produced by anaerobic digestion.

Vermicomposting or composting are further methods of processing digestate.

These technologies improve the quality of the final biofertilizer and stabilise nutrients^[11,25].

In particular, vermicomposting uses earthworms to further break down organic debris, resulting in nutrient-rich vermicast with increased microbial activity and bioavailability^[11].

6) *Napier grass-derived biofertilizers' agronomic advantages*

The benefits of using biofertilizers made from lignocellulosic biomass, like Napier grass, for plant growth, soil health, and overall agricultural sustainability are becoming more widely acknowledged. These advantages result from greater soil physicochemical characteristics, increased microbial activity, and increased nutrient availability. Biomass-based biofertilizers support long-term soil fertility and ecological balance in contrast to synthetic fertilisers^[2,12,30].

7) *Enhancement of Soil Health*

The use of organic biofertilizers is essential for preserving and enhancing soil quality, which is a major factor in determining agricultural productivity. Napier-based biofertilizers have a major role in boosting soil organic matter, which improves soil porosity, structure, and water-holding capacity^[12,21].

Improved nutrient cycling and soil biological activity result from the addition of organic carbon through biomass degradation, which encourages microbial growth. Sustainable soil fertility depends on processes such as nitrogen fixation, phosphorus solubilisation, and organic matter mineralisation, all of which are supported by increased microbial diversity^[22,28].

Additionally, it has been demonstrated that organic amendments made from plant biomass improve aeration and lessen soil compaction, which promotes greater root penetration and growth.

8) *Plant Growth*

Plant growth indices, such as plant height, leaf area, root development, and total biomass yield, have shown notable benefits when biofertilizers made from Napier grass are used. The presence of growth-promoting compounds created during microbial breakdown and increased food availability are the main causes of these effects.

By ensuring a consistent supply of vital components like potassium, phosphorus, and nitrogen, organic biofertilizers increase the efficiency of nutrient uptake. Furthermore, the microbial activity linked to biofertilizers might increase the synthesis of phytohormones that support plant growth and development, such as auxins and gibberellins.

According to studies, crops treated with organic amendments show better physiological performance, such as increased photosynthetic efficiency and chlorophyll content.

9) Cycling and Availability of Nutrients

The role Napier-based biofertilizers play in nutrient cycle is one of their biggest benefits. Organic biofertilizers release nutrients gradually through microbial degradation processes, in contrast to synthetic fertilisers that supply minerals in easily soluble forms. By ensuring consistent nutrient availability throughout the crop growth cycle, this slow-release mechanism lowers the possibility of nutrient losses via volatilisation or leaching. Humic compounds, which improve soil's ability to retain nutrients and exchange cations, are formed as a result of the breakdown of lignocellulosic biomass.

Additionally, biofertilizers promote circular agricultural systems by facilitating the recycling of agricultural biomass and turning waste materials into useful inputs. This method lessens reliance on outside inputs while simultaneously increasing nutrient efficiency.

10) Benefits to the Environment and Sustainability

By lowering dependency on chemical fertilisers and minimising environmental effects, the use of biofertilizers generated from Napier is consistent with sustainable agriculture principles. It has been demonstrated that organic inputs improve soil carbon sequestration, decrease greenhouse gas emissions, and lessen soil and water pollution^[16].

Additionally, by transforming plentiful biomass into useful agricultural inputs, the use of Napier grass as a feedstock promotes resource efficiency. Without competing with food crops, its cultivation on marginal lands enhances sustainable land management^[7,19].

Napier-based biofertilizers provide a feasible route toward resilient and sustainable agricultural systems by enhancing soil health, increasing crop productivity, and minimising environmental degradation.

III. CHALLENGES AND LIMITATIONS

Although Napier grass has great potential as a feedstock for the manufacturing of biofertilizer, a number of obstacles prevent its widespread use. Its high lignin content, which slows down microbial breakdown and lowers the effectiveness of nutrient release during composting and other biological processes, is one of the main limitations^[25]. This frequently calls for co-composting techniques or pretreatment techniques, which can lengthen and complicate processing.

The absence of established procedures for turning Napier grass into biofertilizer is another drawback. Inconsistent product quality and nutritional content can result from variations in processing techniques, such as composting conditions, microbial inoculants, and moisture levels. Large-scale implementation and commercialisation are hampered by this discrepancy.

Additionally, farmers' and researchers' acceptance of Napier-derived biofertilizers is hindered by the lack of information on their nutrient profile, especially NPK values. There is insufficient agronomic validation under field settings because the majority of current studies concentrate on general biomass use rather than particular fertiliser applications^[12].

Logistical and economic considerations are also important. Napier grass is widely available, but not all areas have the infrastructure and labour needed for the harvesting, processing, and transportation of large amounts of biomass. To guarantee practical implementation, these limitations must be overcome.

IV. FUTURE PROSPECTS AND RESEARCH GAPS

There are still a number of significant gaps in the current research on using Napier grass to produce biofertilizer. The dearth of thorough research concentrating especially on Napier grass as a key biofertilizer feedstock is one of the biggest gaps. Its potential in nutrient management is understudied because the majority of literature focuses on its role in the production of bioenergy or feed. To optimise conversion processes, such as anaerobic digestion, fermentation, and composting, more study is needed. Decomposition rates and nutrient availability could be greatly increased by creating effective microbial consortia specifically designed for lignocellulosic biomass^[22].

Furthermore, incorporating Napier-based biofertilizers into contemporary agricultural systems—like hydroponics and precision farming—offers fresh chances for creativity.

Accessibility could be further improved via the creation of low-cost, decentralised production systems, particularly for smallholder farmers. In general, filling all these research gaps will be essential to turning Napier grass from a promising resource into a workable and extensively used sustainable agricultural option.

V. CONCLUSION

The hunt for sustainable and eco-friendly substitutes has accelerated due to the growing concerns about the usage of chemical fertilisers. Because of its high biomass yield, flexibility, and advantageous biochemical composition, Napier grass shows promise as a feedstock for the manufacturing of biofertilizer.

This review emphasises how composting, fermentation, and anaerobic digestion are some of the biological processes that can turn Napier grass into biofertilizer. The resultant biofertilizers have several agronomic advantages, including better soil health, increased plant growth, and effective nutrient cycling. However, issues with restricted field validation, delayed decomposition, and lack of standardisation need to be solved. To fully realise its potential, future research should concentrate on large-scale use, nutritional characterisation, and process optimisation. In summary, Napier grass is a feasible and sustainable source for the production of biofertilizer, and it has the potential to make a substantial contribution to environmentally friendly farming methods and long-term soil fertility management.

REFERENCES

- [1] Savci, S. (2012). Investigation of effect of chemical fertilizers on environment. APCBEE Procedia.
- [2] Vessey, J. K. (2003). Plant growth promoting rhizobacteria as biofertilizers. Plant and Soil.
- [3] Hanna, W. W. (2010). Napier grass: Production and uses. Agronomy Journal.
- [4] Anderson, W. F., et al. (2008). Bioenergy potential of Napier grass. Biomass and Bioenergy.
- [5] Isikgor, F. H., & Becer, C. R. (2016). Lignocellulosic biomass: A sustainable platform. Polymer Chemistry.
- [6] Ojedirani, J. O. (2021). Valorization of Napier grass biomass. Renewable Energy Reports.
- [7] FAO (2013). Grassland management and sustainability. Food and Agriculture Organization.
- [8] Bernal, M. P., et al. (2009). Composting of organic wastes. Waste Management.
- [9] Mshandete, A. M., & Parawira, W. (2009). Biogas technology review. African Journal of Biotechnology.
- [10] Zakarya, I. A. (2025). Bio-decomposer effects on Napier grass. ResearchGate Publication.
- [11] Edwards, C. A. (2011). Vermiculture technology. CRC Press.
- [12] Lal, R. (2015). Soil health and carbon management. Journal of Soil Science.
- [13] Diacono, M., & Montemurro, F. (2010). Organic fertilization effects on crops. Agronomy for Sustainable Development.
- [14] Pretty, J. (2008). Agricultural sustainability concepts. Philosophical Transactions of the Royal Society B.
- [15] FAO (2017). Organic farming benefits. Food and Agriculture Organization.
- [16] Smith, P. (2014). Climate change and agriculture. Global Change Biology.
- [17] Gupta, P. K. (2011). Composting challenges and solutions. Waste Management Research.
- [18] Ghimire, S., et al. (2015). Nutrient dynamics in compost systems. Bioresource Technology.
- [19] Singh, J., & Kalamdhad, A. S. (2013). Effects of composting on soil properties. Ecological Engineering.
- [20] Awasthi, M. K., et al. (2017). Advances in organic waste composting. Renewable and Sustainable Energy Reviews.
- [21] Agegnehu, G., et al. (2017). Benefits of organic amendments on soil. Soil and Tillage Research.
- [22] Bhattacharyya, P. N., & Jha, D. K. (2012). Plant growth-promoting rhizobacteria. World Journal of Microbiology.
- [23] Sharma, S. B., et al. (2013). Phosphate solubilizing microbes. SpringerPlus.
- [24] Chen, J. H. (2006). Organic fertilizers and biofertilizers. FAO Fertilizer Report.
- [25] Kibblewhite, M. G., et al. (2008). Soil health and sustainable agriculture. Philosophical Transactions B.
- [26] Tilman, D., et al. (2002). Agricultural sustainability and inputs. Nature.
- [27] Palm, C. A., et al. (2001). Organic inputs and nutrient cycling. Agriculture, Ecosystems & Environment.
- [28] Lehmann, J., & Joseph, S. (2009). Biochar for environmental management. Earthscan.
- [29] Sanchez, P. A. (2002). Soil fertility restoration in Africa. Nature.
- [30] Vanlauwe, B., et al. (2010). Integrated soil fertility management. Outlook on Agriculture.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)