



# **iJRASET**

International Journal For Research in  
Applied Science and Engineering Technology



---

# **INTERNATIONAL JOURNAL FOR RESEARCH**

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume: 10    Issue: VIII    Month of publication: August 2022**

**DOI: <https://doi.org/10.22214/ijraset.2022.46147>**

**[www.ijraset.com](http://www.ijraset.com)**

**Call:  08813907089**

**E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)**

# Properties of the Ternary Cubic Equation

$$5x^2 - 3y^2 = z^3 \sum$$

G. Janaki<sup>1</sup>, A. Gowri Shankari<sup>2</sup>

<sup>1</sup>Associate Professor, <sup>2</sup>Assistant Professor, PG & Research Department of Mathematics, Cauvery College for Women (Autonomous), Affiliated to Bharathidasan University, Trichy – 18

**Abstract:** To identify its different integral non-zero solutions, the ternary cubic equation  $5x^2 - 3y^2 = z^3$  is taken into consideration. Different integral solution patterns to the ternary cubic equation under consideration are obtained in each pattern by using the linear transformation and the method of factorization; interesting relationships between the solutions and some polygonal numbers, such as pyramidal and central pyramidal numbers, are also displayed.

**Keywords:** Diophantine equations, Ternary equation, Cubic Equation with Three Unknowns, Integral Solutions

## I. INTRODUCTION

Number theory, which is used to explain anything that can be quantified, is the language of patterns and relationships. A polynomial equation called a Diophantine equation can only have integers as solutions. Theories of numbers were featured in [1–3]. A unique Pythagorean triangle problem and its integral solutions are featured in [4, 5]. Higher order equations are taken into account for integral solutions in [6–10].

The non-homogeneous cubic equation with three unknowns represented by the equation is discussed in this communication, and in particular, a few intriguing relationships between the solutions are highlighted.

### A. Notations

$T_{m,a}$ : Polygonal Number of rank  $a$  with side  $m$

$Gno_a$ : Gnomonic Number of rank  $a$

$Star_a$ : Star Number of rank  $a$

$O_a$ : Octahedral Number of rank  $a$

$P_a^m$ : Pyramidal Number of rank  $a$  with sides  $m$

$SO_a$ : Stella Octangula Number of rank  $a$

$CC_a$ : Centered Cube Number of rank  $a$

$CS_a$ : Centered Square Number of rank  $a$

$RD_a$ : Rhombic Dodecagonal Number of rank  $a$

$TO_a$ : Truncated Octahedral Number of rank  $a$

## II. METHOD OF ANALYSIS

A non-zero integral solution to the Cubic equation can be found by

$$5x^2 - 3y^2 = z^3 \quad (1)$$

Upon switching the transformations,

$$x = X + 3T, y = X + 5T, z = 2Z \quad (2)$$

$$\text{in (1) leads to, } X^2 - 15T^2 = 4Z^3 \quad (3)$$

Below, we present illustration of distinct integer non-zero patterns (1)

### A. Pattern: 1

$$\text{Assume } Z = z(a,b) = a^2 - 15b^2 \quad (4)$$

Where a and b are positive integers.

$$\text{And write } 4 = (8 + 2\sqrt{15})(8 - 2\sqrt{15}) \quad (5)$$

Using the factorization approach, replacing (3) with (4) and (5),

$$(X + \sqrt{15}T)(X - \sqrt{15}T) = (8 + 2\sqrt{15})(8 - 2\sqrt{15})(a + \sqrt{15}b)^3(a - \sqrt{15}b)^3$$

Comparing real and imaginary elements while equating similar phrases,  $X = 8a^3 + 360ab^2 + 90a^2b + 450b^3$

$$T = 2a^3 + 90ab^2 + 24a^2b + 120b^3$$

The appropriate integer solutions of equation (1) are provided by substituting the above mentioned values of X and T into equation

$$x = x(a, b) = 14a^3 + 630ab^2 + 162a^2b + 810b^3$$

$$(2) y = y(a, b) = 18a^3 + 810ab^2 + 210a^2b + 1050b^3$$

$$z = z(a, b) = 2a^2 - 30b^2$$

*Properties:*

1.  $-7z(a, a)$  and  $-28z(a, a)$  are Perfect Squares
2.  $-x(1, 1) + y(1, 1) - z(1, 1)$  is a Harshad Number
3.  $y(a, 1) - x(a, 1) - 6P_a^6 - star_a - 2T_{29, a} - T_{26, a} \equiv 239(\text{mod } 123)$
4.  $y(a, 1) - x(a, 1) - 124z(a, 1) - 2SO_a - 91Gno_a \equiv 0(\text{mod } 4051)$
5.  $-y(a, 1) - z(a, 1) + 18P_a^3 + 4Star_a \equiv 1076(\text{mod } 852)$
6.  $x(a, 1) + y(a, 1) - 16z(a, 1) - 2TO_a - 29T_{30, a} \equiv 2352(\text{mod } 1769)$
7.  $-x(a, 1) + y(a, 1) - 23z(a, 1) + 12P_a^5 + T_{18, a} \equiv 930(\text{mod } 187)$

*B. Pattern: 2*

The modified form of equation (3) is

$$X^2 - 15T^2 = 4Z^3 * 1 \quad (6)$$

$$\text{Put 4 in as, } 4 = (8 + 2\sqrt{15})(8 - 2\sqrt{15}) \quad (7)$$

$$\text{and } 1 = (4 + \sqrt{15})(4 - \sqrt{15}) \quad (8)$$

When (7) and (8) are substituted in equation (6) and the process of factorization is used, as described in Pattern 1, the corresponding

$$x = x(a, b) = 110a^3 + 4950ab^2 + 1278a^2b + 6390b^3$$

integer solutions of (1) are represented by  $y = y(a, b) = 142a^3 + 6390ab^2 + 1650a^2b + 8250b^3$

$$z = z(a, b) = 2a^2 - 30b^2$$

*Properties:*

1.  $y(1, 1) - x(1, 1) + z(1, 1) - 22T_{3, 4}$  is a nasty number
2.  $x(1, 1) + y(1, 1) - 26T_{3, 4}$  is a perfect square number
3.  $y(1, 1) - x(1, 1) + 11z(1, 1) - T_{8, 3}$  Is a cubic number
4.  $y(a, 1) - x(a, 1) - 8RD_a - 72T_{15, a} - 886Gno_a \equiv 0(\text{mod } 2762)$
5.  $y(a, 1) - x(a, 1) - 186z(a, 1) - 48O_a - 712Gno_a \equiv 0(\text{mod } 8152)$

$$6. y(a,1) - x(a,1) - 2TO_a - 47T_{22,a} \equiv 1392(\text{mod}1815)$$

$$7. y(a,1) - 825x(a,1) - 213O_a \equiv 33000(\text{mod}6319)$$

$$8. 2x(a,1) - 426z(a,1) - 110SO_a - 284T_{26,a} - 6567Gno_a \equiv 0(\text{mod}6567)$$

C. Pattern: 3

$$\text{Write } 1 = (31 + 8\sqrt{15})(31 - 8\sqrt{15}) \quad (9)$$

By replacing (7) and (9) in equation (6) and factorising the result using the steps in Pattern 1, the corresponding integer solutions of

$$x = x(a,b) = 866a^3 + 38970ab^2 + 10062a^2b + 50310b^3$$

$$(1) \text{ are represented by } y = y(a,b) = 1118a^3 + 50310ab^2 + 12990a^2b + 64950b^3$$

$$z = z(a,b) = 2a^2 - 30b^2$$

Properties:

$$1. y(a,1) - 16z(a,1) - 2236P_a^5 - 1184T_{22,a} - 30483Gno_a \equiv 0(\text{mod}95913)$$

$$2. y(a,1) - x(a,1) + z(a,1) - 189HO_a - 827T_{10,a} - 5796Gno_a \equiv 0(\text{mod}23056)$$

$$3. \frac{1}{10}(x(1,1) - y(1,1)) \text{ is a square number}$$

$$4. x(1,b) + y(1,b) + 23z(1,b) - 138312P_b^7 - 3239Star_b - 44295Gno_b \equiv 0(\text{mod}42086)$$

$$5. y(1,1) - x(1,1) + 24T_{3,4} \text{ is a nasty number}$$

$$6. x(a,1) - 16z(a,1) - 4336CC_a - 11329CS_a + 22658T_{3,a} \equiv 39894(\text{mod}71658)$$

$$7. -x(a,1) + y(a,1) - 23z(a,1) - 63RD_a - 239T_{30,a} \equiv 14013(\text{mod}14195)$$

$$8. y(a,1) - x(a,1) + 19z(a,1) - 126SO_a - 1483T_{6,a} \equiv 14070(\text{mod}12949)$$

Note:

Additionally, 4 and 1 might be written as

$$4 = (62 + 16\sqrt{15})(62 - 16\sqrt{15})$$

$$1 = \frac{(8 + \sqrt{15})(8 - \sqrt{15})}{49}$$

In relation to these options, one may find many patterns of solutions of (1)

### III. CONCLUSION

Three unique patterns of non-zero distinct integer solutions to the given non-homogeneous problem  $5x^2 - 3y^2 = z^3$  are shown in this paper.

For various options of cubic Diophantine equations, additional patterns of non-zero integer unique solutions and their corresponding characteristics may be found.

### REFERENCES

- [1] Carmichael R.D., "The Theory of Numbers and Diophantine Analysis", Dover Publications, New York, 1959
- [2] Dickson L.E., "History of the theory of numbers", Chelsia Publishing Co., Vol II, New York, 1952
- [3] Mordell L.J., "Diophantine Equations", Academic Press, London 1969
- [4] Telang S. G., "Number Theory", Tata Mc Graw- Hill Publishing Company, New Delhi 1996

- [5] Gopalan M.A, Manju Somnath and Vanitha N, Parametric Solutions of  $x^2 - y^6 = z^2$ , Acta Ciencia Indica, Vol XXXIII, 3, 1083 – 1085, 2007
- [6] Gopalan M.A and Janaki G, “Integral solutions of  $(x^2 - y^2)(3x^2 + 3y^2 - 2xy) = 2(z^2 - w^2)p^3$ ”, Impact J. Sci, Tech., 4(1), 97-102, 2010
- [7] Gopalan M.A, Pandichelvi V, “Observations on the ternary cubic equation  $x^3 + y^3 + x^2 - y^2 = 4(z^3 + z^2)$ ”, Archimedes J. Math 1(1), 31 – 37, 2011
- [8] Gopalan M.A and Janaki G, “Integral solutions of  $(x^2 - y^2)(3x^2 + 3y^2 - 2xy) = 2(z^2 - w^2)p^3$ ”, Impact Journal of Science & Technology, Vol – 4, No. 97 – 102, 2010
- [9] Janaki G, Saranya C, “Observations on Ternary Quadratic Diophantine Equation  $6(x^2 + y^2) - 11xy + 3x + 3y + 9 = 72z^2$ ”, International Journal of Innovative Research and Science, Engineering and Technology, Volume 5, Issue 2, February 2016
- [10] Janaki G, Saranya P, “ On the Ternary Cubic Diophantine Equation  $5(x^2 + y^2) - 6xy + 4(x + y) + 4 = 40z^3$ ”, International Journal of Science and Research- online, Vol 5, Issue 3, Pg. No: 227 – 229, March 2016





10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)