



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 10 Issue: VIII Month of publication: August 2022 DOI: https://doi.org/10.22214/ijraset.2022.46147

www.ijraset.com

Call: 🕥 08813907089 🔰 E-mail ID: ijraset@gmail.com



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 10 Issue VIII Aug 2022- Available at www.ijraset.com

# **Properties of the Ternary Cubic Equation**

 $5x^2 - 3y^2 = z^3 \sum$ 

G. Janaki<sup>1</sup>, A. Gowri Shankari<sup>2</sup>

<sup>1</sup>Associate Professor, <sup>2</sup>Assistant Professor, PG & Research Department of Mathematics, Cauvery College for Women (Autonomous), Affiliated to Bharathidasan University, Trichy – 18

Abstract: To identify its different integral non-zero solutions, the ternary cubic equation  $5x^2 - 3y^2 = z^3$  is taken into consideration. Different integral solution patterns to the ternary cubic equation under consideration are obtained in each pattern by using the linear transformation and the method of factorization; interesting relationships between the solutions and some polygonal numbers, such as pyramidal and central pyramidal numbers, are also displayed.

Keywords: Diophantine equations, Ternary equation, Cubic Equation with Three Unknowns, Integral Solutions

#### I. INTRODUCTION

Number theory, which is used to explain anything that can be quantified, is the language of patterns and relationships. A polynomial equation called a Diophantine equation can only have integers as solutions. Theories of numbers were featured in [1-3]. A unique Pythagorean triangle problem and its integral solutions are featured in [4, 5]. Higher order equations are taken into account for integral solutions in [6-10].

The non-homogeneous cubic equation with three unknowns represented by the equation is discussed in this communication, and in particular, a few intriguing relationships between the solutions are highlighted.

A. Notations

 $T_{m, a}$ : Polygonal Number of rank *a* with side *m* 

Gno<sub>a</sub>: Gnomonic Number of rank a

Star<sub>a</sub>: Star Number of rank a

O<sub>a</sub>: Octahedral Number of rank *a* 

 $P_a^m$ : Pyramidal Number of rank *a* with sides *m* 

SO<sub>a</sub>: Stella Octangula Number of rank a

CC<sub>a</sub>: Centered Cube Number of rank *a* 

CS<sub>a</sub>: Centered Square Number of rank *a* 

RD<sub>a</sub>: Rhombic Dodecagonal Number of rank *a* 

TO<sub>a</sub>: Truncated Octahedral Number of rank *a* 

#### II. METHOD OF ANALYSIS

A non-zero integral solution to the Cubic equation can be found by

$$5x^2 - 3y^2 = z^3$$
Upon switching the transformations,
(1)

$$x = X + 3T, y = X + 5T, z = 2Z$$
 (2)

in (1) leads to, 
$$X^2 - 15T^2 = 4Z^3$$
 (3)

Below, we present illustration of distinct integer non-zero patterns (1)

A. Pattern: 1

Assume 
$$Z = z(a,b) = a^2 - 15b^2$$
 (4)



Where a and b are positive integers.

And write 
$$4 = (8 + 2\sqrt{15})(8 - 2\sqrt{15})$$
 (5)  
Using the factorization approach, replacing (3) with (4) and (5),  
 $(X + \sqrt{15}T)(X - \sqrt{15}T) = (8 + 2\sqrt{15})(8 - 2\sqrt{15})(a + \sqrt{15}b)^3(a - \sqrt{15}b)^3$ 

Comparing real and imaginary elements while equating similar phrases,  $X = 8a^3 + 360ab^2 + 90a^2b + 450b^3$ 

$$T = 2a^3 + 90ab^2 + 24a^2b + 120b^3$$

The appropriate integer solutions of equation (1) are provided by substituting the above mentioned values of X and T into equation  $x = x(a,b) = 14a^3 + 630ab^2 + 162a^2b + 810b^3$ 

(2) 
$$y = y(a,b) = 18a^3 + 810ab^2 + 210a^2b + 1050b^3$$
  
 $z = z(a,b) = 2a^2 - 30b^2$ 

Properties:

1. -7z(a,a) and -28z(a,a) are Perfect Squares

2. -x(1,1) + y(1,1) - z(1,1) is a Harshad Number

3.  $y(a,1) - x(a,1) - 6P_a^6 - star_a - 2T_{29,a} - T_{26,a} \equiv 239 \pmod{123}$ 

4. 
$$y(a,1) - x(a,1) - 124z(a,1) - 2SO_a - 91Gno_a \equiv 0 \pmod{4051}$$

2

5. 
$$-y(a,1) - z(a,1) + 18P_a^{-5} + 4Star_a \equiv 1076 \pmod{852}$$
  
6.  $x(a,1) + y(a,1) - 16z(a,1) - 2TO_a - 29T_{30,a} \equiv 2352 \pmod{1769}$ 

7. 
$$-x(a,1) + y(a,1) - 23z(a,1) + 12P_a^5 + T_{18,a} \equiv 930 \pmod{187}$$

B. Pattern: 2

The modified form of equation (3) is

$$X^{2} - 15T^{2} = 4Z^{3} * 1$$
(6)  
Put 4 in as,  $4 = (8 + 2\sqrt{15})(8 - 2\sqrt{15})$ 
(7)

and  $1 = (4 + \sqrt{15})(4 - \sqrt{15})$ 

When (7) and (8) are substituted in equation (6) and the process of factorization is used, as described in Pattern 1, the corresponding

$$x = x(a,b) = 110a^3 + 4950ab^2 + 1278a^2b + 6390b^3$$

(8)

integer solutions of (1) are represented by  $y = y(a,b) = 142a^3 + 6390ab^2 + 1650a^2b + 8250b^3$ 

$$z = z(a,b) = 2a^2 - 30b^2$$

Properties:

1.  $y(1,1) - x(1,1) + z(1,1) - 22T_{3,4}$  is a nasty number

2.  $x(1,1) + y(1,1) - 26T_{3,4}$  is a perfect square number

3.  $y(1,1) - x(1,1) + 11z(1,1) - T_{8,3}$  Is a cubic number

4.  $y(a,1) - x(a,1) - 8RD_a - 72T_{15,a} - 886Gno_a \equiv 0 \pmod{2762}$ 

5.  $y(a,1) - x(a,1) - 186z(a,1) - 48O_a - 712Gno_a \equiv 0 \pmod{8152}$ 



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 10 Issue VIII Aug 2022- Available at www.ijraset.com

(9)

6.  $y(a,1) - x(a,1) - 2TO_a - 47T_{22,a} \equiv 1392 \pmod{1815}$ 

7. 
$$y(a,1) - 825x(a,1) - 213O_a \equiv 33000 \pmod{6319}$$

8. 
$$2x(a,1) - 426z(a,1) - 110SO_a - 284T_{26,a} - 6567Gno_a \equiv 0 \pmod{6567}$$

Write  $1 = (31 + 8\sqrt{15})(31 - 8\sqrt{15})$ 

By replacing (7) and (9) in equation (6) and factorising the result using the steps in Pattern 1, the corresponding integer solutions of

$$x = x(a,b) = 866a^{3} + 38970ab^{2} + 10062a^{2}b + 50310b^{3}$$

(1) are represented by 
$$y = y(a,b) = 1118a^3 + 50310ab^2 + 12990a^2b + 64950b^3$$

$$z = z(a,b) = 2a^2 - 30b^2$$

Properties:

1. 
$$y(a,1) - 16z(a,1) - 2236P_a^{5} - 1184T_{22,a} - 30483Gno_a \equiv 0 \pmod{95913}$$
  
2.  $y(a,1) - x(a,1) + z(a,1) - 189HO_a - 827T_{10,a} - 5796Gno_a \equiv 0 \pmod{23056}$   
3.  $\frac{1}{10}(x(1,1) - y(1,1))$  is a square number

4. 
$$x(1,b) + y(1,b) + 23z(1,b) - 138312P_b^7 - 3239Star_b - 44295Gno_b \equiv 0 \pmod{42086}$$

5.  $y(1,1) - x(1,1) + 24T_{3,4}$  is a nasty number

6.  $x(a,1) - 16z(a,1) - 4336CC_a - 11329CS_a + 22658T_{3,a} \equiv 39894 \pmod{1658}$ 

7. 
$$-x(a,1) + y(a,1) - 23z(a,1) - 63RD_a - 239T_{30,a} \equiv 14013 \pmod{14195}$$

8.  $y(a,1) - x(a,1) + 19z(a,1) - 126SO_a - 1483T_{6,a} \equiv 14070 \pmod{12949}$ 

Additionally, 4 and 1 might be written as

$$4 = (62 + 16\sqrt{15})(62 - 16\sqrt{15})$$
$$1 = \frac{(8 + \sqrt{15})(8 - \sqrt{15})}{49}$$

In relation to these options, one may find many patterns of solutions of (1)

# III. CONCLUSION

Three unique patterns of non-zero distinct integer solutions to the given non-homogeneous problem  $5x^2 - 3y^2 = z^3$  are shown in this paper.

For various options of cubic Diophantine equations, additional patterns of non-zero integer unique solutions and their corresponding characteristics may be found.

### REFERENCES

- [1] Carmichael R.D., "The Theory of Numbers and Diophantine Analysis", Dover Publications, New York, 1959
- [2] Dickson L.E., "History of the theory of numbers", Chelsia Publishing Co., Vol II, New York, 1952
- [3] Mordell L.J., "Diophantine Equations", Academic Press, London 1969
- [4] Telang S. G., "Number Theory", Tata Mc Graw- Hill Publishing Company, New Delhi 1996



# International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538

Volume 10 Issue VIII Aug 2022- Available at www.ijraset.com

, Acta Ciencia Indica, Vol XXXIII, 3, 1083 – 1085, 2007

, Impact J. Sci, Tech., 4(1), 97-102, 2010  

$$x^{3} + y^{3} + x^{2} - y^{2} = 4(z^{3} + z^{2})$$
  
", Archimedes J. Math 1(1), 31 – 37,

[7] Gopalan M.A, Pandichelvi V, "Observations on the ternary cubic equation 2011

$$(x^{2} - y^{2})(3x^{2} + 3y^{2} - 2xy) = 2(z^{2} - w^{2})p^{3}_{,,}$$

, Impact Journal of Science &

[8] Gopalan M.A and Janaki G, "Integral solutions of Technology, Vol – 4, No. 97 – 102, 2010

$$6(x^2 + y^2) - 11xy + 3x + 3y + 9 = 72z^2$$
", International

- [9] Janaki G, Saranya C, "Observations on Ternary Quadratic Diophantine Equation Journal of Innovative Research and Science, Engineering and Technology, Volume 5, Issue 2, February 2016
- [10] Janaki G, Saranya P, "On the Ternary Cubic Diophantine Equation  $5(x^2 + y^2) 6xy + 4(x + y) + 4 = 40z^3$ ", International Journal of Science and Research- online, Vol 5, Issue 3, Pg. No: 227 229, March 2016











45.98



IMPACT FACTOR: 7.129







# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24\*7 Support on Whatsapp)