



# IJRASET

International Journal For Research in  
Applied Science and Engineering Technology



---

# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume:** 14    **Issue:** III    **Month of publication:** March 2026

**DOI:** <https://doi.org/10.22214/ijraset.2026.79234>

[www.ijraset.com](http://www.ijraset.com)

Call:  08813907089

E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)

# Role of Lifestyle Factors in Female Infertility: A Review of Current Evidence

Dr. Namrata Gabu<sup>1</sup>, Dr. Harsh Pandya<sup>2</sup>, Dr. Akshar Gabu<sup>3</sup>

**Abstract:** *Infertility is a growing global health concern affecting millions of couples worldwide. Female infertility may result from ovulatory dysfunction, tubal disease, uterine abnormalities, endometriosis, or unexplained causes. In recent years, increasing attention has been directed toward modifiable lifestyle factors that may contribute to infertility. Obesity, poor diet, sedentary behavior, smoking, alcohol consumption, psychological stress, sleep disturbances, and environmental toxin exposure have all been implicated in impaired fertility. Understanding these factors is important because many are potentially reversible and may improve reproductive outcomes without invasive interventions. This review summarizes the current evidence regarding lifestyle-related contributors to female infertility and highlights the importance of counseling and preventive strategies in reproductive medicine.*

**Keywords:** *infertility, female infertility, obesity, lifestyle, ovulation, reproductive health*

## I. INTRODUCTION

Infertility is defined as the inability to achieve pregnancy after 12 months of regular unprotected intercourse. It affects approximately 10–15% of couples globally and has significant psychological, social, and economic consequences.

Female infertility may arise due to several etiologies, including ovulatory disorders, tubal pathology, endometriosis, diminished ovarian reserve, uterine abnormalities, and unexplained infertility. While many causes are non-modifiable, lifestyle factors have emerged as important contributors to reproductive dysfunction.

With increasing urbanization and changes in dietary habits, physical activity, sleep patterns, and environmental exposures, lifestyle-related infertility is becoming more common. Recognition of these factors is essential because lifestyle modification may improve spontaneous conception rates as well as outcomes of assisted reproductive techniques.

## II. OBESITY AND INFERTILITY

Obesity is one of the most significant modifiable risk factors for infertility. Excess adipose tissue disrupts hormonal balance by increasing insulin resistance, androgen production, and inflammatory cytokines. These changes can impair follicular development, ovulation, and endometrial receptivity. Obese women are more likely to experience anovulation, menstrual irregularities, polycystic ovarian syndrome, poor oocyte quality, reduced implantation rates, and higher miscarriage rates. Body mass index greater than 30 kg/m<sup>2</sup> has been associated with lower spontaneous pregnancy rates and poorer outcomes following in vitro fertilization.

## III. POLYCYSTIC OVARIAN SYNDROME AND METABOLIC FACTORS

Polycystic ovarian syndrome is one of the leading causes of female infertility. Women with PCOS often have obesity, insulin resistance, hyperinsulinemia, and hyperandrogenism. Weight reduction in women with PCOS can improve ovulation, menstrual regularity, insulin sensitivity, and pregnancy rates. Even a modest reduction of 5–10% of body weight may significantly improve fertility outcomes.

## IV. DIET AND FERTILITY

Dietary habits influence reproductive health through their effects on metabolism, oxidative stress, and hormonal regulation.

Healthy dietary patterns rich in fruits, vegetables, whole grains, unsaturated fats, and plant-based proteins have been associated with better fertility outcomes.

In contrast, diets high in refined carbohydrates, saturated fats, sugary beverages, and processed foods may contribute to insulin resistance, obesity, and ovulatory dysfunction.

Micronutrients such as folic acid, vitamin D, iron, zinc, selenium, and omega-3 fatty acids may also play important roles in female fertility.

## V. PHYSICAL ACTIVITY

Regular physical activity improves insulin sensitivity, body composition, and cardiovascular health, all of which contribute to reproductive function.

Moderate exercise is associated with improved ovulation, reduced obesity-related infertility, and better fertility treatment outcomes. However, excessive strenuous exercise may suppress ovulation and lead to hypothalamic amenorrhea, particularly in women with low body weight. Thus, balanced physical activity is important for maintaining reproductive health.

## VI. PSYCHOLOGICAL STRESS AND INFERTILITY

Psychological stress is increasingly recognized as both a cause and consequence of infertility. Chronic stress may affect the hypothalamic-pituitary-ovarian axis, alter gonadotropin secretion, and impair ovulation.

Women experiencing infertility often report anxiety, depression, marital stress, and reduced quality of life.

Stress management techniques such as counseling, yoga, mindfulness, and cognitive behavioral therapy may help improve emotional well-being and fertility outcomes.

## VII. SLEEP AND REPRODUCTIVE HEALTH

Poor sleep quality and inadequate sleep duration may negatively affect reproductive hormones and ovulatory function.

Sleep disturbances can alter melatonin secretion, cortisol levels, gonadotropin release, and insulin sensitivity.

Women working night shifts or experiencing chronic sleep deprivation may therefore have increased risk of infertility.

## VIII. SMOKING, ALCOHOL, AND ENVIRONMENTAL TOXINS

Smoking has well-established negative effects on female fertility. It accelerates ovarian aging, reduces ovarian reserve, and increases the risk of miscarriage and ectopic pregnancy.

Alcohol intake may interfere with ovulation and hormone metabolism, while exposure to environmental toxins such as pesticides, plastics, heavy metals, and endocrine-disrupting chemicals has been linked with impaired fertility.

Common environmental toxins include bisphenol A, phthalates, lead, mercury, and pesticides. Reducing exposure to these agents may improve reproductive health.

## IX. LIFESTYLE MODIFICATION AND FERTILITY TREATMENT OUTCOMES

Lifestyle optimization before fertility treatment can improve success rates of ovulation induction, intrauterine insemination, and assisted reproductive techniques. Recommended interventions include weight reduction in overweight women, balanced diet, smoking cessation, regular exercise, adequate sleep, stress reduction, and limiting alcohol and caffeine.

Preconception counseling should therefore be considered an essential component of infertility management.

## X. CONCLUSION

Lifestyle factors play a major role in female infertility and may significantly influence natural conception as well as assisted reproductive outcomes. Obesity, poor diet, sedentary behavior, stress, smoking, sleep disturbances, and environmental toxins all contribute to impaired fertility. Since many of these factors are modifiable, early counseling and lifestyle intervention should be incorporated into routine infertility evaluation and treatment. Addressing these issues may improve reproductive outcomes, reduce the need for invasive treatments, and enhance overall health in women seeking pregnancy.

## REFERENCES

- [1] World Health Organization. Infertility prevalence estimates, 1990–2021.
- [2] Practice Committee of the American Society for Reproductive Medicine. Obesity and reproduction: a committee opinion.
- [3] Sharma R, Biedenharn KR, Fedor JM, Agarwal A. Lifestyle factors and reproductive health.
- [4] Palomba S, Santagni S, La Sala GB. Complications and challenges associated with polycystic ovary syndrome.
- [5] Gaskins AJ, Chavarro JE. Diet and fertility: a review.
- [6] Homan GF, Davies M, Norman R. The impact of lifestyle factors on reproductive performance.
- [7] Domar AD, Rooney KL, Wiegand B, Orav EJ, Alper MM. The impact of stress on fertility.
- [8] Ramlau-Hansen CH, et al. Sleep disturbances and fertility outcomes.
- [9] Augood C, Duckitt K, Templeton AA. Smoking and female infertility.
- [10] ACOG Committee Opinion on preconception counseling and infertility management.



10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)