



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 13 Issue: V Month of publication: May 2025

DOI: <https://doi.org/10.22214/ijraset.2025.70113>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Seasonal Variations in Physicochemical Characteristics of “Ram Mandir Pond”, Vasai, Dist. - Palghar, Maharashtra (INDIA)

Supriya Nitin Ghadi¹, Chandrakant Marathe²

¹Scholar, Shri J.J.T. University, Jhunjhunu, Rajasthan

²Principal, Sanskruti Arts, Science and Commerce college Lalonde, Boiser

Abstract: Ponds remain important resources of water from ancient time. Vasai - Virar city is suburban part of Mumbai city comes under district Palghar. In the city there are more than 100 ponds on record. Of these ponds ‘Ram Mandir Pond’ situated in the heart of city. 20 physicochemical parameters selected for studying pond water were Temperature, pH, Conductivity, Turbidity, Total solids, DS, TSS Dissolved Oxygen, Acidity, Alkalinity, BOD, COD, Nitrate, Nitrite, Total Phosphorus, Silica as SiO_2 , Sulphate, H_2S , Chlorides as Cl, and Total hardness as CaCO_3 . Physicochemical study of this pond reveals seasonal variation of pH of water ranging from 6.65 to the higher as 8.93. Seasonal variation in Total hardness ranging from 470 to 860 mg/l of carbonates. Nitrate values in water found highest during winter i.e. 1.11 mg/l and lowest during Monsoon i.e. 0.377 mg/l. Total phosphorus recorded in three seasons was 0.27, 0.16 and 0.12 mg/l in monsoon winter and summer respectively. Most of the parameters are in and around standard values. Physicochemical studies indicate pond is of mesotrophic category. Remedies for its conservation and maintenance are suggested.

Key words: Physicochemical parameters, Trophic level of pond, pond conservation pond maintenance.

I. INTRODUCTION

Temple ponds are the age old tradition of Indian culture. Temple ponds, often referred to as temple tanks or sacred ponds, hold significant cultural and religious importance in India. These ponds are typically located within or near Hindu temples and serve various purposes that are deeply rooted in tradition and spirituality. Temple ponds are considered sacred bodies of water. Before entering the temple for worship, devotees often ritually purify themselves by taking a dip in the temple pond. This act symbolizes the cleansing of both body and mind, preparing the devotee for spiritual communion with the deity. Beyond their religious significance, temple ponds play a practical role in local ecology. They often serve as reservoirs for rainwater harvesting, supporting groundwater recharge and maintaining biodiversity by providing habitats for various aquatic species.

Overall, temple ponds in India are not just bodies of water; they are repositories of cultural heritage, religious devotion, and ecological balance, embodying the interconnectedness of spirituality and sustainability in Indian traditions. In recent times, some temple ponds have faced challenges such as pollution, neglect, and encroachment. Physicochemical and biological parameters help to understand the quality of water and ecology of the pond (Arti & Saxena, 2012) (Amarendra Hari Chandan et al. 2016). Hence study of selected pond is planned to know the status of the pond and suggestions for preservation and restoration of this important cultural and ecological assets.

II. MATERIALS & METHODS

A. Study area:

Location Vasai Virar with a jurisdiction of about 311 sq. km. is geographically located at 19.3919° N, Latitude and 72.8397° E Longitude at about 36 feet above mean sea level. It is located in Maharashtra, India's Konkan region. Vasai-Virar is one of the twin cities that make up the Mumbai Metropolitan Region (MMR). Vasai-Virar city has been separated from Greater Mumbai and Mira-Bhayandar City by Vasai Creek.



Figure 1 Ram mandir Pond in Bolinj Virar a Satellite View.



Figure 2 Ram mandir Pond in Bolinj Virar- measurement.

Ram Mandir Pond is situated in Bolinj Division of Vasai Virar city. Bolinj is an old part of Present city having ancient temple of deity Shree Ram. Pond is on right side (south) of the temple (towards south). Pond is about 3 km from Virar railway station. Location of the pond is $19^{\circ}26'32.6''\text{N}$ $72^{\circ}47'43.0''\text{E}$. The temple pond is recently protected by constructed concrete walls and iron bars. Rain water is the only source of temple pond.

Physicochemical Analysis:Physicochemical analysis of water sample from the Ram Mandir Pond is carried about during the year 2020-21. Water samples collected in the Morning Between 7 am to 10 am. 20 Physicochemical parameters selected for the study were Temperature, pH, Conductivity, Turbidity, Total solids, DS, TSS Dissolved Oxygen, Acidity, Alkalinity, BOD, COD, Nitrate, Nitrite, Total Phosphorus, Silica as SiO₂, Sulphate, H₂S, Chlorides as Cl, and Total hardness as CaCO₃.All analysis was done following the standard method of APHA 22nd Edition (2012), Trivedi RK, Goel PK (1984) Chemical and Biological Methods for Water Pollution Studies and guide manual: water and wastewater analysis (CPCB).

III. RESULT & DISCUSSION

Results of Physicochemical parameters analysis is presented in the table 1. TemperatureDifference in ambient and water temperature is very negligible. Water temperature recoded were 28⁰ C in Monsoon and 27⁰ C in Winter and 29⁰ C during Summer.Amb. temperature remained slightly higher than the water temperature throughout the study period. Temperature is connected to other non-living factors such as sunlight, dissolved oxygen and turbidity. Warm water does not hold as much dissolved oxygen as cold water; therefore, dissolved oxygen levels drop as a pond heats up in the summer.

pH ranges from 6.05 to 8.93. Highest pH wasrecorded during the summer. During winter it was 8.05. pH from 6.5 to 8.5 is Permissible range. During Summer it slightly crossing the limits.In general, basic ponds with high pH values tend to be more productive than acidic ponds with low pH values. (Shanthi k et al,2002).

µmhos/cm. lowest in winter and high during monsoon and summer. According to Bhatt et al. (1999) EC in water is due to ionization of dissolved inorganic solids and is a measure of total dissolved solids and salinity. Salts that dissolved in water break in to positively charged and negatively charged ions.Shannon & Brezonick (1972) considered electrical conductivity as an indicator of trophic status.According to them Electrical conductivity is caused because of the presence of solids in water therefore these two parameters (EC & TDS) are closely related.

Table 1 Physicochemical parameters analysis of Ram Mandir Pond

SR. NO.	PARAMETERS	MONSOON	WINTER	SUMMER
1	Amb. Temperature	27	24	30
	Water Temperature	28	27	29
2	pH	6.65	8.05	8.93
3	Conductivity (m mho)	2.72	1.665	2.01
4	Turbidity (NTU)	2	5.2	9.5
5	Total solids (mg/L)	1740	1240	1520
6	TDS (mg/L)	840	1100	1290
7	TSS (mg/L)	900	140	230
8	Dissolved Oxygen (mg/L)	1.2	1.3	3.12
9	Acidity (mg/L)	100	50	100
10	Alkalinity (mg/L)	900	950	450
11	BOD	1	1	0.48
12	COD	88	44	88

13	Nitrate (mg/L)	0.377	1.11	1.15
14	Nitrite (mg/L)	0.14	0.17	0.12
15	Total Phosphorus (mg/L)	0.27	0.16	0.121
16	Silica as SiO ₂ (mg/L)	6.52	0.71	0.745
17	Sulphate (mg/L)	112	55.18	52.05
18	H ₂ S	36.14	12.75	8.5
19	Chlorides as Cl (mg/L)	603.5	383.4	447.3
20	Total hardness as CaCO ₃	480	860	470

Both water color and turbidity affect the penetration of light into lakes and ponds. Turbidity refers to water cloudiness and represents the concentration of suspended particles in the water. The most common types of suspended particles in ponds are sediments and phytoplankton. In present study turbidity was recorded 2NTU in Monsoon 5.2 NTU during winter and Highest value 9.5 NTU during summer. High value maybe due to evaporation of water and growth of Phytoplankton.

TDS varied from 840 mg/L to 1290 mg/L at Ram Mandir Pond whereas TDS. highest value recorded during summer. TSS (Total suspended solids) recorded in present study highest during monsoon 900 mg/L during monsoon and 230mg/L during summer and 140 mg/L during winter season. Total solids also exhibit the same trend like TSS i.e. 1740 mg/L in monsoon, 1240 mg/L in winter and highest 1520 mg/L in summer.

Dissolve oxygen important for the survival of aquatic organism. 3.5 to 5 Mg/L is limit for the DO. During monsoon and winter, it found low. Whereas, during summer it was found suitable 3.12. Acidity of water sample ranges from 50 to 100 mg/L and alkalinity ranging from 450 to 900 mg/L. which is suitable for aquatic life. Mean BOD recorded was 1. It lowered during summer to 0.48 and COD lowered to 0.48 during winter. Highest value of nitrate was 1.15 during summer and Nitrite found highest during winter i.e. 1.17 mg/L. Phosphorus showed high value 0.27 during monsoon and lower during winter and summer High value Si 6.52 found during monsoon. Similarly sulphate value recorded 112 mg/L during monsoon which was highest among three seasons. H₂S also high during monsoon i.e. 36.14. Chlorides in the same trend found high during monsoon. 603 and lowest 383.4 during winter. Total hardness values recorded ranges from 470 to 860mg/L. During winter hardness found to be at its peak.

IV. CONCLUSION

Most of Studied parameter are in good range indicating suitability for aquatic life. pH range is within permissible limit. Total Solids showed higher values. BOD, Phosphorus Nitrate nitrates are within permissible limit. Total hardness surpassed the limit during winter only. From above study it is concluded that it is suitable for the aquaculture, agriculture and other uses. However, suitability for drinking purpose must be depended on the further study of water sample for microbial content and presence of heavy metals.

REFERENCES

- [1] APHA (2012), Standard Methods for the Examination of Water and Waste Waters, 22nd Edition, American Public Health Association, Washington, DC.
- [2] Amarendra Hari Chandan, Himansu Sekhar, Patra, Kabir Mohan Sethy, Saswat Kumar Mohanty and Rabindra Panda (2016). Water Quality Index and Correlation Study of religious ponds in the temple city, Bhubaneswar, Odisha. American journal of scientific and industrial research. 7(2):22-31.
- [3] Arti Sharma & R.S. Saxena. 2012. limnological studies of Neem River with special reference to phytoplankton diversity. Ind.J. stud. Res vol.2(1)15-23.
- [4] Bhatt, L. R, Lacoul, P., Lekhal, H. D., and Jha, P. K. 1999. Physico-chemical characteristics and phytoplankton for Taudha lake, Kathmandu. Pollution Research, (18): pp. 353-358.
- [5] Shanthi, K., Ramasamy, K. and Laxmanaperumalsamy, P. 2002. Hydrological study of singanallur lake at Coimbatore, India Nature Env. And pollution Technology. 1(2): 97-101
- [6] Shannon, E. E. and Brezonick, P. 1972. Limnological characteristics of north and central Florida lakes. LimnolOceang. 17(1): 97-110.
- [7] Trivedi RK, Goel PK. Chemical and Biological Methods for Water Pollution Studies. Environmental Publication, Karad, India, 1984, 211-215.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)