iJRASET
International Journal For Research in Applied Science and Engineering Technology



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

**Volume:** 14 **Issue:** III **Month of publication:** March 2026

**DOI:** https://doi.org/10.22214/ijraset.2026.77871

www.ijraset.com

Call: 08813907089 | E-mail ID: ijraset@gmail.com

International Journal for Research in Applied Science & Engineering Technology (IJRASET)
*ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538*
*Volume 14 Issue III Mar 2026- Available at www.ijraset.com*

# Solutions of Pell's Equation Involving Wilson's Primes

C. Saranya[1], M. Shalini[2]
[1]*Associate Professor,* [2]*PG Student, PG and Research Department of Mathematics, Cauvery College for Women (Autonomous), (Affiliated to Bharathidasan University), Tiruchirappalli, Tamilnadu, India.*

***Abstract:*** ***The main goal of this study is to find non-trivial integer solutions to Pell's equation involving Wilson's primes. We obtain a series of unique integer solution and construct recurrence linkage between them using Brahmagupta's lemma. The derived solutions and their relationships are shown numerically, illustrating the function of Wilson's primes in producing these solutions.***

***Keywords: Brahmagupta lemma, Wilson's primes, recurrence relations, Diophantine equation, Pell's equation.***

## I. INTRODUCTION

A central theme in number theory is the study of Diophantine equations, which seek integer solutions to polynomial equations. Among them, Pell's equation $x^2 = Dy^2 + 1$. where $D$ is a positive square-free integer, occupies a fundamental position due to its deep connections with quadratic fields, continued fractions, and algebraic number theory [1,3-7,9&11]. The equation admits infinitely many integer solutions whenever a fundamental non-trivial solution exists [2,8,10 &12]. Over the centuries, several powerful techniques have been developed to determine these solutions, including continued fraction expansions and algebraic methods. One of the most elegant classical tools in generating solutions of Pell's equation is Brahmagupta's lemma, which provides a composition law for solutions. This lemma enables the recursive construction of infinitely many solutions from a known pair, revealing the multiplicative structure underlying the solution set. The lemma not only demonstrates the richness of the equation's algebraic framework but also establishes recurrence relations that govern the growth of its solutions.

In recent developments, special classes of prime numbers have been employed to construct and characterize solutions of Diophantine equations. In particular, Wilson's primes, defined by the congruence $(p-1)! \equiv -1 (\mathrm{mod}\ p^2)$ form a rare and remarkable subclass of primes arising from Wilson's theorem. Their exceptional modular properties and strong arithmetic constraints make them suitable candidates for generating structured integer solutions. The interplay between the factorial congruence condition of Wilson's primes and the quadratic form of Pell's equation provides a novel framework for deriving non-trivial solution sequences.

Motivated by these connections, the present study investigates Pell's equation through the combined application of Wilson's primes and Brahmagupta's lemma. By exploiting the arithmetic properties of Wilson's primes, we construct recurrence relations and establish the existence of distinct non-zero integer solutions. The theoretical results are substantiated with illustrative numerical examples, thereby contributing to the broader understanding of structured solution generation in classical Diophantine equations.

## II. METHOD OF ANALYSIS

*1) Theorem*

The sequence of non-zero distinct integer solution to the Pell's equation

$$x^2 - Dy^2 = z^2 \qquad (1)$$

where, $D = W^2 - 1$, $z$ is an integer and $W$ is a Wilson's primes given by

$$x_s = \frac{z}{2}\left(\sqrt{D+1} f_s + \sqrt{D} g_s\right)$$

$$y_s = \frac{z}{2\sqrt{D}}\left(\sqrt{D} f_s + \sqrt{D+1} g_s\right) \qquad s=0,1,2\ldots$$

©IJRASET: All Rights are Reserved 

and their recurrence relations are

$$x_{s+2} - 2\sqrt{D+1}x_{s+1} + x_s = 0$$

$$y_{s+2} - 2\sqrt{D+1}y_{s+1} + y_s = 0 \qquad s=0,1,2...$$

Proof:

The initial solution of (1) is $(x_0, y_0)$ is given by $x_0 = z\sqrt{D+1}, y_0 = z.$

To find the other solutions of (1), consider the Pell's equation

$$x^2 = D y^2 + 1$$

whose initial solution $(\tilde{x}_s, \tilde{y}_s)$ is given by

$$\tilde{x}_s = \frac{1}{2} f_s$$

$$\tilde{y}_s = \frac{1}{2\sqrt{D}} g_s \qquad s=0,1,2...$$

where $f_s = \left(\sqrt{D+1} + \sqrt{D}\right)^{s+1} + \left(\sqrt{D+1} - \sqrt{D}\right)^{s+1}$

$$g_s = \left(\sqrt{D+1} + \sqrt{D}\right)^{s+1} - \left(\sqrt{D+1} - \sqrt{D}\right)^{s+1},$$

Applying Brahmagupta's lemma between the solutions

$$(x_0, y_0) \text{ and } (\tilde{x}_s, \tilde{y}_s),$$

the sequence of non-zero distinct integer solutions to (1) are

$$x_s = \frac{z}{2}\left(\sqrt{D+1} f_s + \sqrt{D} g_s\right)$$

$$y_s = \frac{z}{2\sqrt{D}}\left(\sqrt{D} f_s + 2\sqrt{D+1} g_s\right) \qquad s=0,1,2...$$

and their recurrence relations are found to be

$$x_{s+2} - 2\sqrt{D+1}x_{s+1} + x_s = 0$$

$$y_{s+2} - 2\sqrt{D+1}\, y_{s+1} + y_s = 0, \qquad s=0,1,2\ldots$$

*2) Illustration*

The solutions of Pell's equation involving Wilson's Primes and its corresponding recurrence relations are presented in the table below:

| Wilson's primes $W$ | Integer $z$ | Pell's equation | Sequence of Integer Solutions | Recurrence relation |
|---|---|---|---|---|
| 5 | 2 | $x^2 = 24y^2 + 4$ | $x_s = 5f_s + \sqrt{24}g_s$<br>$y_s = \frac{1}{\sqrt{24}}\left(\sqrt{24}\, f_s + 5g_s\right)$ | $x_{s+2} - 10x_{s+1} + x_s = 0$<br>$y_{s+2} - 10y_{s+1} + y_s = 0$ |
| 13 | 4 | $x^2 = 168y^2 + 16$ | $x_s = 2\left(13f_s + \sqrt{168}g_s\right)$<br>$y_s = \frac{2}{\sqrt{168}}\left(\sqrt{168} f_s + 13g_s\right)$ | $x_{s+2} - 26x_{s+1} + x_s = 0$<br>$y_{s+2} - 26y_{s+1} + y_s = 0$ |

| | | | | |
|---|---|---|---|---|
| 563 | 6 | $x^2 = 316968y^2 + 36$ | $x_s = 3\left(563f_s + \sqrt{316968}g_s\right)$<br>$y_s = \frac{3}{\sqrt{316968}}\left(\sqrt{316968}f_s + 563g_s\right)$ | $x_{s+2} - 1126x_{s+1} + x_s = 0$<br>$y_{s+2} - 1126y_{s+1} + y_s = 0$ |


## III. CONCLUSION

In this paper, we have presented a structured method for generating non-zero and distinct integer solutions to Pell's equation by incorporating the arithmetic properties of Wilson's primes. By applying Brahmagupta's lemma as a fundamental tool, we established recurrence relations that systematically produce infinite sequences of solutions from an initial solution pair. The factorial congruence condition characterizing Wilson's primes plays a crucial role in constructing these solution families and in strengthening the underlying algebraic structure of the equation.

The results demonstrate that the classical theory of Pell's equation can be significantly enriched through the integration of special prime classes such as Wilson primes. This interplay between prime number theory and quadratic Diophantine equations not only deepens the understanding of solution structures but also opens new directions for further research in the study of Pell-type equations and related number-theoretic problems.

## REFERENCES

[1] Carmichael, R D., "History of Theory of numbers and Diophantine Analysis", Dover Publication, New york, 1959.
[2] Gopalan M A.,Vidhyalakshmi S., ÖZER ÖZEN., "A Collection of Pellian Equation (Solutions and Properties)", Akinik Publications, Publication No:1, 2018.
[3] Hardy, G H., Wright, E M., "An Introduction to the Theory of Numbers", 5th ed., Clarendon Press, 1979.
[4] Hua, L K., "Introduction to the Theory of Numbers", Springer-Verlag, Berlin-New York, 1982.
[5] Mordell,L J., "Diophantine equations", Academic press, London, 1969.
[6] Nagell,T., "Introduction to Number theory", Chelsea publishing company, New York, 1981.
[7] OZER ÖZEN., "A Certain Type of Regular Diophantine Triples and Their Non-Extendibility", Turkish Journal of Analysis and Number Theory, 7(2), 50-55, 2019.
[8] Saranya,C., "Solutions of Pell's Equation and Exponential Diophantine Equation Utilizing Ramanujan Prime Numbers of Order 2", Ganita, Volume 72, Issue 2, 455-460, 2024.
[9] Saranya,C., Janaki,G., "Solutions of Pell's Equation Involving Jarasandha Numbers", International Journal Of Scientific Research In Mathematical and Statistical Sciences, Vol 6, Issue 1, 234-236, Feb 2019.
[10] Saranya,C., Swethashree, A., "Solution of Exponential Diophantine Equation Involving Fermat Primes", International Journal for Research in Applied Science & Engineering Technology (IJRASET), Volume 13 Issue III, Mar 2025
[11] Shailesh Shirali., Yogananda, C.S., "Number Theory", University Press, Hyderabad, 2003.
[12] Vidhya, S., Janaki, G., "An Integral solution of Negative Pell's equation involving Two-digit Sphenic numbers", International Journal of Computer Sciences and Engineering, vol 6, Issue 7, 444-445, July 2018.



10.22214/IJRASET

45.98

IMPACT FACTOR: 7.129

IMPACT FACTOR: 7.429

# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)