



IJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 12 **Issue:** V **Month of publication:** May 2024

DOI: <https://doi.org/10.22214/ijraset.2024.62844>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Studies on Benzoxazine Derivatives

Mohamed Saad Muftah

Chemistry Department, Faculty of Sciences, Bani Walid University, Libya

Abstract: The 5, 6, 7 trimethoxy (1, 4) benzoxazine -2,3diones was synthesized by reaction of 2-amino-3,4,5 trimethoxyphenol with oxalyl chloride. The new compound of benzoxazine derivative was characterized by elemental analysis, mass spectroscopy, infrared and $^1\text{H-NMR}$. The compound exhibited antifungal and antibacterial activities.

Keywords: Benzoxazine, mass spectroscopy, $^1\text{H-NMR}$ and biological activities

I. INTRODUCTION

In recent years benzoxazine derivatives containing nitrogen have attracted increased attention due to the broad spectrum of their biological activities ^[1] like antibacterial ^[2,3] antifungal ^[4]

Benzoxazines show a wide range of biological activity which are key molecules for the synthesis of various pharmaceutical agents as antifungal ^[5], antimicrobial ^[6]. There are only few reports suggesting the antimicrobial properties of benzoxazine derivatives so far ^[7,8]. Benzoxazine derivatives also display various biological activities such as anticancer ^[9], antimicrobial ^[10], antifungi ^[11], antiplatelet ^[12], and antituberculosis activities ^[13].

II. EXPERIMENTAL PART

A. Instrumentation

All melting points were measured on a Gallenkamp melting apparatus and uncorrected. The IR spectra of compounds were recorded on Shimadzu IR Affinity FTIR spectrophotometer using KBr discs and the values are expressed in $\delta\text{ cm}^{-1}$. The $^1\text{H-NMR}$ spectra of compound were recorded on Bruker Avance 400 MHz NMR spectrophotometer using DMSO as an internal standard and the values are expressed in $\delta\text{ ppm}$. The elemental analysis of the compounds were recorded on a Perkin – Elmer 2400 CHN elemental analyzer.

The mass spectra were recorded on a GCMS-QP-1000EX mass spectrometer at (70eV). Synthesis of 5, 6, 7 trimethoxy {1, 4} benzoxazine -2, 3 diones: In a round bottom flask 250ml dissolve (16.3gm, 0.1mole) of 2-amino-3,4,5 trimethoxyphenol in 100ml ethanol then add (12.6gm, 0.1mole) of oxalyl chloride, stir the mixture at room temperature for 2hrs then add 2ml of piperidine. The mixture was heated to reflux for 10hrs and kept overnight. The solid was separated by filtration. The solid was recrystallized from ethanol.




Fig 1: 5, 6, 7 trimethoxy {1,4} benzoxazine -2,3 diones

III. RESULTS AND DISCUSSION

Spectroscopic studies of 5, 6, 7 trimethoxy (1, 4) benzoxazine -

2, 3 diones: The infrared spectrum of table (2) displayed a strong band at 1665cm^{-1} corresponding to $\nu(\text{C}=\text{O})$ (lactone), as well as the IR spectrum exhibited absorption band at 3218cm^{-1} due to (N-H). The $^1\text{H-NMR}$ spectrum of the (2) in deuterated DMSO- d_6 table (2) exhibited from low field to high field, the following signals (δ /ppm): 8.13(S,H,NH), 7.65(S,1H,Ar-H) and 3.81(3S,9H,3OCH₃). The mass spectrum of compound (1) exhibited the molecular ion peak $[\text{M}]^+$ at m/e 253 (67%) indicating the molecular formula C₁₁H₁₁NO₆, besides other fragments which are in accordance with the proposed structure 238(44%), 223(45%), 193(71%), 163(80%), 149(93%), 77(100%).

A. Biological Activity

Measurement of antimicrobial activity using diffusion disc method: Antibacterial and antifungal activities of some synthesized compound was screened using the disc diffusion method. All the tested compounds showed antibacterial and antifungal activity and these activities were compared to standard amoxicillin, the results of antimicrobial studies are given in table 3, the antimicrobial activity of newly synthesized benzoxazine was conducted against gram positive and gram negative groups namely staphylococcus aureus and Escherichia coli respectively as well as Aspergillus flavus and candida albicans as tested fungi by disc diffusion method. Amoxicillin was employed as reference standard to compare the results. Each test compare was dissolved in dimethyl sulphoxide (DMSO). The concentration of DMSO solutions was 0.1 mg/ml.

Table 1: Physical characterization of benzoxazine derivative

M.P/C°color	Solvent yield	MF(M.wt)	Elemental analysis calc/found			
			C%	H%	N%	S%
184-186	Ethanol	C ₁₁ H ₁₁ NO ₆	52,17	4,37	5,53	
Brown	90	253,208	51,82	3,95	4,79	

Table 2: Spectroscopic for benzoxazine derivative

Compound No.	IR(KBr) ν cm ⁻¹	¹ HNMR δ (PPm)	MS,Mm/z (%) relevan
(1)	ν N-H3218 ν C=O1665(Lacton)	8,13(S, H, NH) 7,65(S,1H, Ar-H) 3,81(3S,9H,3OCH ₃)	378(86,24%),359(32,19%) 352(49,23%),348(65,36%),332(54,10%), 266(83,29%),251(72,14%), 128(100%), 78(44,07%)

Table 3: The inhibition zone diameter of some benzoxazine derivative

Sample/ standard	Inhibition zone diameter (mm/mg sample)			
	<i>Staphylococcus aureus</i> (G+)	<i>Escherichia coli</i> (G-)	<i>Candida albicans</i> (fungus)	<i>Aspergillus</i>
benzoxazine	17	20	16	12
Amoxicillin	30	32	24	21

REFERENCES

- [1] Hossain M, Nada AK. A review on Heterocyclic synthesis and their application in medicinal chemistry of imidazole moiety, science journal of chemistry 2018;6(5):83-94.
- [2] Azab ME, Youssef MM, ELBodaany EA. Synthesis and Antibacterial Evaluation of Novel Heterocyclic compounds containing a sulfonamide moiety, Molecules. 2013;18:832-844.
- [3] Martins P, Jesus J, Santos S, Raposo LR, Roma-Rodrigues C, Baptista PV, Fernandes AR. Heterocyclic Anticancer Compounds; Recent Advances and the paradigm Shift towards the Use of nanomedicines Tool Box, Molecules, (2015);20:16852-16891.
- [4] George M, Joseph L, Sadanandan HR. A Research on Synthesis of Oxazine Derivatives & Screening of Oxazine Derivatives for certain pharmacological Activities, INTERNATIONAL Journal of pharmacy & pharmaceutical Research, 2016;3:14-42.
- [5] Fringuelli R, Pietrella D, Schiaffella F, Guarraci A, Perito S, Bistonib F, et al. Anti Candida albicans properties of novel benzoxazine analogues, Bioorg. Med. Chem. 2002; 10:1681-1686.
- [6] Alper-Hayta S, Aki-Sener E, Tekiner-Gulbas B, Yildiz I, Temiz-Arpaci O, Yalcin I, et al. Synthesis, antimicrobial activity and QSARs of new benzoxazine-3-one, Eur. J Med. Chem. 2006; 41:1398-1404.
- [7] Sun Y, Niu C, Wang H, Zhang B, Xie F, Xu H, et al. Discovery of 3-benzoxazine-2,4 dione analogues as allosteric mitogen activated kinase (MEK) inhibitors and anti-tuberculosis 71(EV71) agents, bioorg. Med. Chem. 2016; 24:3472-3482.
- [8] Itoni G, Vasam S, Guguloth V, Vadde R. One-pot multicomponent synthesis of [1,4] benzoxazine isoxazole hybrids and their antibacterial activity, IJPCBS. 2018;8:118-124
- [9] Rajitha C, Dubey PK, Sunku V, Piedrafita FJ, Veeramani VR, Pal M, et al. Eur. J Med Chem. 2011; 46(10):4887-4896.
- [10] Alper-Hayta S, Aki-Sener E, Tekiner-Gulbas B, Yildiz I, Temiz-Arpaci O, Yalcin I, Altanlar N, et al. Eur J Med. Chem. 2006; 41(12):1398-1404.
- [11] Shalaby AA, Elkhayry AM, Shiba SA, Ahmed AA, Hanafi AA, Arch. Pharm. 2000; 333(11):365-372.
- [12] Ihmaid SK, AlRawi JM, Bradley CJ, Angove MJ, Robertson MN, Eur. J Med Chem, 2012; 57:85-101.
- [13] Nemecek P, Mocak J, Lehotay J, Waissner K. Chem. Pep. 2013; 67(3):305-312.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)