



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 12 Issue: III Month of publication: March 2024 DOI: https://doi.org/10.22214/ijraset.2024.58881

www.ijraset.com

Call: 🕥 08813907089 🔰 E-mail ID: ijraset@gmail.com



Tackling Ternary Cubic Diophantine Equation $4(\alpha^2 + \beta^2) - 7(\alpha\beta) = 64\gamma^3$

P. Saranya¹, D. Keerthana²

¹Associate Professor, ²PG Student, PG and Research Department of Mathematics, Cauvery College for Women (Autonomous), Affiliated to Bharathidasan University, Trichy-18.

Abstract: A few interesting characteristics among the solutions and the patterns of non-zero integral solutions to the non-homogeneous cubic equation with three unknowns represented by the Diophantine equation $4(\alpha^2 + \beta^2) - 7(\alpha\beta) = 64\gamma^3$ are examined.

Keywords: Ternary cubic equation, Diophantine equation, Integer solution.

I. INTRODUCTION

Mathematics, conveys knowledge of number and is the global language of the world. Number theory is a field in pure mathematics that studies integral valued functions. In number theory, primes and prime factorization are highly important concepts. Fermat is recognized as the Father of modern number theory despite being lawyer by practice and an elementary "amateur" mathematician. Modern number theory, geometric number theory and probabilistic number theory. A Diophantine equation is a polynomial equation with two or more unknowns seeking only integer solutions. The word Diophantine refers to "Diophantus of Alexandria", a Hellenistic mathematician who lived in the century and is credited with being one of the first who introduced symbolism of algebra. A homogeneous degree 3 polynomial in three variables is called a Ternary Cubic forms. All cubic equation have either three real roots or one real root with two imaginary roots. Three degree polynomials are referred to as cubic equation. In [1-5] Elementary Number Theory concepts are studied, [6-9] examined quadratic Diophantine equation, [10-15] referred the cubic Diophantine equation and exponential Diophantine equation. A non-homogeneous ternary cubic equation with three unknowns $4(\alpha^2 + \beta^2) - 7(\alpha\beta) = 64\gamma^3$ is solved in this paper and using some interesting number properties.

II. NOTATIONS

1) $P_n = 2T_n = n(n+1) =$ Pronic number of rank n.

- 2) $TO_n = 16n^3 33n^2 + 24n 6 =$ Truncated octrahedral number of rank n.
- 3) $T_{8,n} = n(3n-2) = \text{Octagonal number of rank n.}$
- 4) $S_n = 6n(n-1) + 1 =$ Star number of rank n.
- 5) $T_{12,n} = n(5n 4) =$ Dodecagonal number of rank n.
- 6) $T_{26,n} = n(12n-11) =$ Icosihexagonal number of rank n.
- 7) $W_n = 2^n n 1 =$ Woodall number of rank n.
- 8) $Gno_n = (2n-1) =$ Gnomonic number of rank n.
- 9) $T_{30.n} = n(14n 13) =$ Triacontagonal number of rank n.
- 10) $T_{22,n} = n(10n 9) =$ Icosidigonal number of rank n.

III. METHOD OF ANALYSIS

(1)

Considering the Diophantine equation in ternary form $4(\alpha^{2} + \beta^{2}) - 7(\alpha\beta) = 64\gamma^{3}$



Substitution Taking,

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538

Volume 12 Issue III Mar 2024- Available at www.ijraset.com

The linear transformation used is,

$$\alpha = \lambda + \eta \text{ and } \beta = \lambda - \eta$$
(2)
By (2), (1) reflects as,

$$\lambda^2 + 15\eta^2 = 64\gamma^3$$
(3)
Below is the theorem derived from solving equation (3) using different substitutions.
Theorem: Multitude of solutions of $4(\alpha^2 + \beta^2) - 7(\alpha\beta) = 64\gamma^3$
Substitution: 1.1
Taking,

$$\gamma = \upsilon^2 + 15\omega^2$$
(4)
We can write 64 as,

$$64 = (7 + i\sqrt{15})(7 - i\sqrt{15})$$
(5)
Substitute the equation (4) and (5) in (3),

$$(\lambda + i\sqrt{15}\eta)(\lambda - i\sqrt{15}\eta) = (7 + i\sqrt{15})(7 - i\sqrt{15})(\upsilon + i\sqrt{15}\omega)^3(\upsilon - i\sqrt{15\omega})^3$$
Equating the real and imaginary terms,

Equating the

$$\lambda = 7\upsilon^3 - 315\upsilon\omega^2 - 45\upsilon^2\omega + 225\omega^3$$
$$\eta = \upsilon^3 - 45\upsilon\omega^2 + 21\upsilon^2\omega - 105\omega^3$$

Reducing equation (2) using (λ) and (η) gives,

$$\alpha = 8\upsilon^3 - 360\upsilon\omega^2 - 24\upsilon^2\omega + 120\omega^3$$
$$\beta = 6\upsilon^3 - 270\upsilon\omega^2 - 66\upsilon^2\omega + 330\omega^3$$
$$\gamma = \upsilon^2 + 15\omega^2$$

Inference: 1.1

- 1. $\gamma(c,c) 2T_{18,c} 7Gno_c \equiv 0 \pmod{7}$
- 2. $\beta(1,1) \alpha(1,1)$ is a perfect square

3.
$$3\alpha(c,1) - 4\beta(c,1) - 16T_{26,c} - 88Gno_c + 872 = 0$$

4. $\alpha(1,1) + 639$ is a woodall number

5. $5\gamma(c,1) - T_{12,c} - 2Gno_c \equiv 0 \pmod{77}$

Substitution: 1.2

Rewrite 64 as,
$$64 = (2 + i2\sqrt{15})(2 - i2\sqrt{15})$$
 (6)

From (4) and (6) in (3),

$$(\lambda + i\sqrt{15}\eta)(\lambda - i\sqrt{15}\eta) = (2 + i2\sqrt{15})(2 - i2\sqrt{15})(\upsilon + i\sqrt{15}\omega)^3(\upsilon - i\sqrt{15}\omega)^3$$

Equating the terms both real and imaginary,

$$\lambda = 2\upsilon^3 - 90\upsilon\omega^2 - 90\upsilon^2\omega + 450\omega^3$$
$$\eta = 2\upsilon^3 - 90\upsilon\omega^2 + 6\upsilon^2\omega - 30\omega^3$$

Reducing equation (2) using (λ) and (η) gives,

$$\alpha = 4\upsilon^3 - 180\upsilon\omega^2 - 84\upsilon^2\omega + 420\omega^3$$
$$\beta = -96\upsilon^2\omega + 480\omega^3$$
$$\gamma = \upsilon^2 + 15\omega^2$$

Inference: 1.2

- $1.\beta(1,1) 161$ is a carol number
- 2. $4\alpha(c,1) TO_c + 101T_{8,c} + 473Gno_c \equiv 0 \pmod{1213}$

3. $\frac{\gamma(c,c)}{c^2}$ - 3 is a star number



Volume 12 Issue III Mar 2024- Available at www.ijraset.com

4. $6\gamma(c,1) + \beta(c,1) + 18T_{12,c} + 36Gno_c \equiv 0 \pmod{534}$

1

5. $\alpha(1,1) - 50$ is a pronic number Substitution: 1.3 Write equation (3) as,

$$\lambda^2 + 15\eta^2 = 64\gamma^3 \times 1 \tag{7}$$

'1' can be written as,

$$=\frac{(1+i\sqrt{15})(1-i\sqrt{15})}{16}$$
(8)

Equating (4), (5) and (8) in (7) we have,

$$(\lambda + i\sqrt{15}\eta)(\lambda - i\sqrt{15}\eta) = (7 + i\sqrt{15})(7 - i\sqrt{15})(\upsilon + i\sqrt{15}\omega)^3(\upsilon - i\sqrt{15}\omega)^3\frac{(1 + i\sqrt{15})(1 - i\sqrt{15})}{16}$$

Equating the terms both real and imaginary,

$$\lambda = \frac{1}{4} (-8\upsilon^3 + 1800\omega^3 + 360\upsilon\omega^2 - 360\upsilon^2\omega)$$
$$\eta = \frac{1}{4} (8\upsilon^3 + 120\omega^3 - 360\upsilon\omega^2 - 24\upsilon^2\omega)$$

Reducing equation (2) using (λ) and (η) gives,

$$\alpha = 480\omega^3 + 84\upsilon^2\omega - 180\upsilon\omega^2$$
$$\beta = -4\upsilon^3 + 420\omega^3 + 96\upsilon^2\omega$$
$$\gamma = \upsilon^2 + 15\omega^2$$

Inference: 1.3

1. $\alpha(1,1) - 9\gamma(1,1)$ is a pronic number

- 2. $\alpha(c,1) 6T_{30,c} + 51Gno_c \equiv 0 \pmod{429}$
- 3. $\frac{32\beta(c,c)}{c^3}$ 257 is a carol number
- 4. $\alpha(c,1) 4\gamma(c,1) 8T_{22,c} + 54Gno_c \equiv 0 \pmod{366}$
- 5. $3\gamma(1,1) 3$ is a hexagonal number

IV. CONCLUSION

In this paper, we've employed diverse substitutions to seek integral solutions for the non-homogeneous ternary cubic equation. These same substitutions can be applied to solve analogous equations.

REFERENCES

- [1] David M. Burton "Elementary Number Theory", Sixth edition, Tata mc graw hill edition.
- [2] Neville Robbins "Beginning Number Theory", Second edition, Naroa Publishing House.
- [3] Gareth A. Jones and J. Marry Jones "Elementary Number Theory", Springer international edition.
- [4] Dr. Sudhir K. Pundir and Dr. Rimple Pundir "Theory of Numbers", Second edition, Published by: K.K. Mittal prakashan, meerut-250001, Laser typesetting.
- [5] Thomas Koshy "Elementary Number Theory with Applications", Published by Elsevier, a division of Reed Elsevier India private limited, 17-A/1, Main Ring Road, Lajpat Nagar-Iv, New Delhi-110 024, India.
- [6] M.A.Gopalan, P.Shanmuganadham and A.Vijayashankar, (2008), "On Binary Quadratic Equation $x^2 5xy + y^2 + 8x 20y + 15 = 0$ ", Acta Ciencia Indica, Vol .XXXIVM.NO.4, pp.1803-1805.
- [7] P.Saranya, G. Janaki, "On the Exponential Diophantine Equation $36^x + 3^y = z^2$ ", International Research Journal of Engineering and Technology, Vol 4, Issue 110, pp: 1042-1044, 2017.
- [8] C.Saranya, G, Janaki, "Observations on the Ternary Quadratic Diophantine Equation $5x^2 + 7y^2 = 972z^2$ ", Jamal Academic Research Journal- An Interdisciplinary, Special Issue, February 2016, pg no.305-308, Impact Factor, ISSN No: 0973-0303.



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.538 Volume 12 Issue III Mar 2024- Available at www.ijraset.com

- [9] G. Janaki, P. Saranya, "On the Ternary Quadratic Diophantine Equation $5(x^2 + y^2) 6xy = 4z^2$ ", Imperial Journal of Interdisciplinary Research, Vol 2, Issue 3, pp: 396-397, 2016.
- [10] M.A.Gopalan, S.Vidhyalakshmi, A.Kavitha, "Observation on the Ternary Cubic Equation $x^2 + y^2 + xy = 12z^3$ ", Antartica J.Math 10(5),2013,453-460.
- [11] S.Vidhyalakshmi, T.R. Usha Rani and M.A. Gopalan "Integral Solution of Non-Homogenous Ternary Cubic Equation $ax^2 + by^2 = (a + b)z^3$ ", Diophantus J.Math., 2(1), 31-38, 2013.
- [12] G. Janaki, P. Saranya, "On the Ternary Cubic Diophantine Equation $5(x^2 + y^2) 6xy + 4(x + y) + 4 = 40z^3$ ", Imperial Journal of Interdisciplinary research. Vol 5, Issue 3, pp: 227-229, 2016.
- [13] C. Saranya, G. Janaki, "Integral Solutions of the Ternary Cubic Equation $3(x^2 + y^2) 4xy + 2(x + y + 1) = 972z^3$ ", International Journal of Engineering and Technology, Vol 4, Issue 3, March 2017.pg no.665-669.
- [14] P.Saranya, G. Janaki, On the Exponential Diophantine Equation $36^x + 3^y = z^2$, International Research Journal of Engineering and Technology, Vol 4, Issue 11, Nov 2017, Pg No: 1042-1044.
- [15] P.Saranya, G. Janaki, Ascertainment on The Exponential Equation $p^{3c} a(p^{2c} p^b) = p^{b+c}$, Compliance Engineering Journal, Vol 10, Issue 8, August 2019, Pg No:225-229.
- [16] G.Janaki, P.Saranya, On the Ternary Cubic Diophantine Equation $5(x^2 + y^2) 6xy + 4(x + y) + 4 = 40z^3$, International Journal of Science and Research-Online, Vol 5, Issue 3, March 2016, Pg No: 227-229.
- [17] P.Saranya, G. Janaki, Explication of The Transcendental Equation $p + \sqrt{p^3 + q^3 pq} + \sqrt[3]{r^2 + s^2} = (m^2 + 1)t^3$, Adalya Journal, Vol 8, Issue 9, September 2019, Pg No: 70-73.
- [18] Saranya, P., G. Janaki, and M. Shri Padmapriya. "Elucidation of the Transcendence Equation $j + \sqrt{(j^2 + k^3 jk) + \sqrt[3]{(l^2 + m^2)}} = h^3 (2^{2n} + 1)$." (2023): Pg No: 28-36.
- [19] Saranya, P., G. Janaki, and K. Poorani. "On the Ternion Cubical Diophantine Equation $5(m^2 + n^2)-6(mn)+8(m+n)+16=370p^3$." (2023): Pg No: 13-21.
- [20] Saranya, P., and M. Priya. "Transcendental Equation Involving Palindrome Number." Arya Bhatta Journal of Mathematics and Informatics 15, no. 1 (2023): 61-66.











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24*7 Support on Whatsapp)