



IJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 13 **Issue:** II **Month of publication:** February 2025

DOI: <https://doi.org/10.22214/ijraset.2025.67075>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Terpenoids Associating Medicinal Plants and Its Values in Pharmacognosy: An Overview

Somenath Bhattacharya¹, Soumallya Chakraborty²

^{1,2}Assistant Professor, Department of Pharmaceutical Chemistry, Global College of Pharmaceutical Technology, Nadia, West Bengal, India

Abstract: Terpenoids are important chemical constituents of medicinal plant. These are basically hydrocarbons. This type of constituent is usually found in many plants like Dill, Caraway, Fennel, Nutmeg, Cinnamon, Tulsi, Eucalyptus, Chenopodium, Ajowan, Coriander, Cardamom, etc across the World. Various types of terpenoids like Myrene, Ocimene, Nerol, Geraniol, Nerol, Geraniol, Phellandral, Perillaldehyde, Santonin, Cadalene, Manool, Lanosterol, Squalene, Lupeol, Betulin, etc are observed in various plants. These constituents can be used carminative, stimulant, flavoring agent, anthelmintics, expectorant, antiseptic, rubefacient, antispasmodic agent, etc. Terpenoids are found in India, China, Europe, USA, Japan, Russia, Indonesia, Caribbean islands, etc. This review article deals with the classifications of terpenoids and role of medicinal plants with terpenoids in details.

Keywords: Terpenoids, Dill, Caraway, Fennel, Nutmeg, Cinnamon, Tulsi, Eucalyptus, Chenopodium, Ajowan, Coriander, Cardamom.

I. INTRODUCTION

Terpenoids are present in many species of plants like Dill, Caraway, Cardamom, Fennel, Ajowan, Nutmeg, Black Pepper, Tulsi, Cinnamon, Coriander, Clove, etc across the different parts of World like Asia, Europe, USA, etc. Chemically it is hydrocarbon in where it should follow the isoprene unit rule (C_5H_8). It is identified by Sudan III chemical in where plant drug is treated with alcoholic Sudan III. Red color shows the presence of volatile oils in terpenoids. Terpenoids are insoluble in water but dissolve in organic solvents. These are colorless solid or liquid with a pleasant, aromatic and characteristic odor. It has optical activity. The terpenoids are having the portions of terpenes, ketones, alcohols, acids, aldehydes and esters. Terpenoids are oxidized easily. It also can be identified by treating the plant part with tincture alkane drop and showing red color. Red color indicates the presence of volatile oils. Different types of family members like Umbelliferae, Liliaceae, Labiatae, Lauraceae, Zingiberaceae, Rutaceae, Myristicaceae, Myrtaceae, etc are observed among the plants belonging from terpenoids. Chemical constituents like Nerol, Phellandral, Geraniol, Betulin, Manool, Lupeol, etc are showing different activities like anthelmintics, antiseptic, antispasmodic, stimulant, carminative, expectorant, flavoring agent, rubefacient, etc. Terpenoids are present in volatile oil. [1-3].

II. CLASSIFICATION OF TERPENES

The classification of terpenes is given in the following table 1.

Table 1: Descriptions of Terpenoids [1, 3-4]

Sl. No.	Types	Molecular Formula	No. of Carbon Atoms	No. of Hydrogen Atoms	No. of Isoprene Units
1	Hemiterpene	C_5H_8	5	8	1
2	Monoterpene	$C_{10}H_{16}$	10	16	2
3	Sesquiterpene	$C_{15}H_{24}$	15	24	3
4	Diterpene	$C_{20}H_{32}$	20	32	4
5	Sesterterpenes	$C_{25}H_{40}$	25	40	5
6	Triterpenes	$C_{30}H_{48}$	30	48	6
7	Tetraterpenes	$C_{40}H_{64}$	40	64	8
8	Pentaterpenes	$C_{50}H_{80}$	50	80	10
9	Polyterpenes	$(C_5H_8)_n$	>50	>80	n

III. CHEMICAL CONSTITUENTS OF TERPENOIDS

The chemical constituents of terpenoids are given in the following table 2.

Table 2: Chemical Constituents of Terpenoids [1, 3-8]

Sl. No.	Terpenes	Sub-divisions	Chemical Constituents
1	Monoterpenes	Acyclic Monoterpenes	Myrcene, Nerol
		Monocyclic Monoterpenes	Menthol, Phellandral
2	Sesquiterpenes	Monocyclic Sesquiterpenes	Zingiberene
		Bicyclic Sesquiterpenes	Cadalene, Santonin
3	Diterpene	Acyclic Diterpenes	Phytol
		Monocyclic Diterpenes	Vitamin A ₁
		Bicyclic Diterpenes	Manool
		Tricyclic Diterpenes	Abietic Acid, Podocarpic Acid
4	Triterpenes	Acyclic Triterpenes	Squalene
		Tetracyclic Triterpenes	Agnosterol, Lanosterol
		Pentacyclic Triterpenes	Asiatic Acid, Ursolic Acid, Glycyrrhetic Acid, Betulin, Lupeol

IV. ALLOCATION OF TERPENOIDS ALONG WITH MEDICINAL PLANTS

Table 3: Medicinal Plants with Terpenoids in Details

Sl. No.	Plant Name	Scientific Name	Family Name	Parts Used	Chemical Constituents	Uses	References
1	Coriander	<i>Coriandrum sativum</i>	Umbelliferae	Dried ripe fruits	Coriandrol, Geraniol, Pinene, L-borneol	Carminative, Stimulant, Aromatic, Flavoring agent	[1, 9-10]
2	Dill	<i>Anethum graveolens</i>	Umbelliferae	Dried ripe fruits	Carvone, D-limonene, Dihydrocarvone, Phellandrene	Carminative, Stimulant, Aromatic, Also used in gastric problems	[1, 11-12]
3	Ajowan	<i>Trachyspermum ammi</i>	Umbelliferae	Dried ripe fruits	Thymol, Pinene, p-cymene	Carminative, Stimulant, Antispasmodic, Antiseptic, Insecticide, Antifungal, Anthelmintic, Flavoring agent Also used in bronchitis	[1, 13-14]
4	Fennel	<i>Foeniculum vulgare</i>	Umbelliferae	Dried ripe fruits	Anethole, Fenchone, Phellandrene, Limonene, Methyl chavicol	Carminative, Stimulant, Aromatic, Flavoring agent, Expectorant	[1, 15-16]
5	Cardamom	<i>Elettaria cardamomum</i>	Zingiberaceae	Dried ripe fruits	Borneol, α -terpineol, Cineole	Carminative, Stimulant, Aromatic, Flavoring agent	[1, 17-18]
6	Caraway	<i>Carum carvi</i>	Umbelliferae	Dried ripe fruits	Carvacrol, Carvone, Limonene, Dihydro carvone	Carminative, Flavoring agent, Stimulant, Aromatic, Spice	[1, 19-20]

7	Spearmint	<i>Mentha spicata</i>	Labiatae	Dried leaves and flowering tops	Pinene, Phellandrene, Cineole, Linalool, l-carvone	Carminative, Flavoring agent	[1, 21-22]
8	Nutmeg	<i>Myristica fragrans</i>	Myristicaceae	Dried kernels from the seeds	Saffrole, Elimicin, Myristicin, Geraniol, d-camphene	Carminative, Flavoring agent, Stimulant, Aromatic	[1, 23-24]
9	Anise	<i>Pimpinella anisum</i>	Umbelliferae	Dried ripe fruits	Anisaldehyde, Chavicol, Methyl chavicol	Carminative, Expectorant, Stimulant, Flavoring agent, Used in production of dentifrices	[1, 25-26]
10	Celery	<i>Apium graveolens</i>	Umbelliferae	Dried ripe fruits	d-limonene, d-selinine, sedanolide, sedanoic acid	Carminative, Stimulant, Tonic, Sedative, Flavoring agent, also used for the treatment of rheumatism	[1, 27]
11	Black Pepper	<i>Piper nigrum</i>	Piperaceae	Dried unripe fruit	Piperine, Piperidine, Caryophyllene, l-phellandrene	Carminative, Stimulant, Aromatic, Stomachic, Spice	[1, 28-30]
12	Cumin	<i>Cuminum cyminum</i>	Umbelliferae	Dried ripe fruits	Cuminaldehyde, Pinene, Hydrocuminine, Cuminic alcohol, Phellandrene	Carminative, Stimulant, Used in the treatment of dyspepsia and diarrhoea	[1, 31-32]
13	Tulsi	<i>Ocimum Sanctum</i>	Lamiaceae	Fresh as well as dried leaves	Eugenol, Ccarvacrol, Methyl eugenol, Caryophylline	Antibacterial, Spasmolytic, Stimulant, Insecticidal, Expectorant, Aromatic, Anti-inflammatory, Stomachic	[1, 33-36]
14	Rasna	<i>Alpinia officinarum</i>	Zingiberaceae	Dried rhizomes	Pinene, Cineole, Alpinol, Galangol, Dioxylavnlol, Galangin	Antibacterial, Aromatic, Carminative, Stimulant, Anti-rheumatic	[1, 37-38]
15	Shati	<i>Hedychium spicatum</i>	Zingiberaceae	Dried rhizomes (sliced)	8-caryophyllene, Cineole, Limonene	Stomachic, Carminative, Stimulant, Tonic, Aromatic	[1, 39-42]
16	Garlic	<i>Allium sativum</i>	Liliaceae	Bulb	Allicin, Alliin	Aphrodisiac, Carminative,	[1, 43-44]

						Stimulant, Expectorant, Anti-dysentery, Antibacterial, Condiment, Anthelmintics, Disinfectant, Rubefacient	
17	Arnica	<i>Arnica montana</i>	Compositae	Dried flowering tops and heads	Helenalin, Arnifolin, Epoxyhelenalin, 13-dehydrohelenalin	Cosmetic, Anti-rheumatic, Anthelmintics, Antitumor, Antihyperlipidemic, Antifungal, Anti-inflammatory	[1, 45-46]
18	Jatamansi	<i>Nardostachys jatamansi</i>	Valerianaceae	Dried rhizomes	Nardostachone, Jatamansone	Antispasmodic, Sedative, Diuretic, Stomachic, Emmenagogue, Stimulant, Anti-epileptic, Anti-arrhythmic, Flavoring agent, Hair tonic	[1, 47-48]
19	Cinnamon	<i>Cinnamomum zeylanicum</i>	Lauraceae	Dried inner bark of the shoot	Eugenol, Caryophyllene, Cymene, Pinene, Phellandrene	Carminative, Mild astringent, Stomachic, Antiseptic, Aromatic, Stimulant, Condiment, Spice, Also used in the preparation of perfumes and dentifrices	[1, 49-50]
20	Clove	<i>Eugenia caryophyllus</i>	Myrtaceae	Dried flower buds	Eugenol, Caryophyllenes	Carminative, Flavoring agent, Stimulant, Antiseptic, Dental analgesic, Aromatic	[1, 51-52]
21	Crocus	<i>Crocus sativus</i>	Iridaceae	Dried stigmas	Crocetin, Picrocrocin	Antispasmodic, Stimulant, Emmenagogue, Coloring agent, Flavoring agent	[1, 53-54]
22	Annatto	<i>Bixa orellana</i>	Bixaceae	Dried seeds	Bixin	Antioxidant	[1, 55-56]
23	Valerian	<i>Valeriana wallichii</i>	Valerianaceae	Dried rhizomes and roots	Valeric acid, Valtrate, Camphene, Borneol acetate, Valerenic acid	Antispasmodic, Carminative, Stimulant, Antidepressant, Tonic	[1, 57]

24	Cubeb	<i>Piper cubeba</i>	Piperaceae	Dried unripe fruits	Cubebin, Cubebic acid, Cadinene	Aromatic, Stimulant, Antiseptic, Used in urinary and cough treatment	[1, 58]
25	Saussurea	<i>Saussurea lappa</i>	Compositae	Dried roots	Camphene, Dihydrolactone, Costuslactone	Expectorant and used for the treatment of asthma	[1, 59-60]
26	Acorus	<i>Acorus calamus</i>	Araceae	Dried rhizomes	Eugenol, Asarone, Acorine, Asaraldehyde	Carminative, Vermifuge, Stimulant, Insect repellent, Flavoring agent, Sedative, Antidepressant, Antiepileptic	[1, 61-63]
27	Feverfew	<i>Tanacetum parthenium</i>	Asteraceae	Dried as well as fresh leaves	Camphor, Borneol, Farnesene, Partholide, Chrysanthemolide, Chrysanthemonin, Parthenolide, Chrysanthemine A, Chrysanthemine B	Used for the treatment of arthritis, migraine, fever, vertigo	[1, 64-67]
28	Coleus	<i>Coleus forskohlii</i>	Labiatae	Roots	Forskolin, Deoxycoleonol, Coleonol B, Coleonol C, Coleon F, Coleon E	Vasodilator, Used in high blood pressure and glaucoma treatment	[1, 68]
29	Vidang	<i>Embelia ribes</i>	Myrsinaceae	Dried fruits	Vilangin, Embelin	Carminative, Astringent, Anthelmintics, Stimulant	[1, 69-70]
30	Bitter Orange Peel	<i>Citrus aurantium</i>	Rutaceae	Dried as well as fresh ripe fruits outer part (Pericarp)	Limonene, Citronellal, Citral	Carminative, Aromatic, Stomachic, Flavoring agent, Perfuming agent, Antioxidant, Antimicrobial	[1, 71-72]
31	Rosemary Oil	<i>Rosmarinus officinalis</i>	Labiatae	Fresh flowering tops	Borneol, Camphor, Pinene, d-camphene	Carminative, Flavoring agent, Stimulant, Refrigerant, Natural insecticides	[1, 73-76]
32	Oil of Vetiver	<i>Vetiveria zizanioides</i>	Gramineae	Roots	Vetiverol, Vetivenol, α -vetivone, β -vetivone	Flavoring agent, Aromatic, Stimulant, Refrigerant, Stomachic, Antibacterial	[1, 77-78]
33	Hops	<i>Humulus lupulus</i>	Cannabaceae	Dried female	Lupulone, Humulone	Spasmolytic, Sedative	[1, 79-81]

				flowers			
34	Artemisia	<i>Artemisia annua</i>	Asteraceae	Herb	Caryophyllene, Camphor, Myrcene, Artemisinin, Arteether, Artemisinic acid, Arteannuin B	Anti-malarial, Antioxidant	[1, 82-83]
35	Oil of Sandal Wood	<i>Santalum album</i>	Santalaceae	Heart wood	β -santalol, Santenone, Santene, Santalene, Santalone	Used for the treatment of dysurea	[1, 84-86]
36	Davana Oil	<i>Artemisia pallens</i>	Compositae	Flowering herbs	Davanone, Cineol, Geraniol, Borneol, Eugenol, Linalool	Flavoring agent, Perfuming gent, Anti-diabetic, Anti-helmintic, Antipyretic, Antibacterial, Antifungal, Used to treat wound healing, Immuno-modulator	[1, 87-88]
37	Citronella Oil	<i>Cymbopogon nardus</i>	Graminae	Fresh leaves	Citronellal, Geraniol, d-camphene, Borneol, Methyl eugenol, Linalool, Limonene, Elmicin, Cadinene	Flavoring agent, Perfuming agent, Used in mosquito repellent cream	[1, 89]
38	Lavender Oil	<i>Lavandula officinalis</i>	Labiatae	Fresh flowering tops	Linalool, Geraniol, Pinene, Cineol, Linalyl acetate	Carminative, Aromatic, Flavoring agent, Perfuming agent, Antibacterial	[1, 90]
39	Palmarosa Oil	<i>Cymbopogon martini</i>	Graminae	Fresh leaves with tops	Geraniol, Linalool, Citronellal	Perfuming agent, Insect repellent, Anti-rheumatic, Bronchodilator, Vasodilator, Spasmolytic	[1, 91-92]
40	Gaultheria Oil	<i>Gaultheria procumbens</i>	Ericaceae	Leaves	Gaultherin	Vermicide, Anti-rheumatic, Flavoring agent, Perfuming agent, Antibacterial, Antioxidant	[1, 93-94]
41	Peppermint Oil	<i>Mentha piperita</i>	Labiatae	Fresh flowering tops and leaves	l-limonene, Cineole, Pinene, Isopulegone, Camphene, Menthone,	Antiseptic, Carminative, Flavoring agent, Stimulant	[1, 95-96]

					Pulegone, Menthofuran		
42	Lemon Grass Oil	<i>Cymbopogon flexuosus</i>	Graminae	Leaves with aerial parts	Citronellal, Citral, Nerol, Geraniol, Methylheptenol	Flavoring agent, Perfuming agent	[1, 97]
43	Geranium Oil	<i>Pelargonium graveolens</i>	Geraniaceae	Fresh stems and leaves	Geraniol, β -citronellol	Flavoring agent	[1, 98]
44	Camphor Oil	<i>Cinnamomum camphora</i>	Lauraceae	Wood	d-pinene, Camphor, Eucalyptol, Cineole, Eugenol, Phellandrene	Flavoring agent, Rubefacient, Anti-inflammatory, Antioxidant	[1, 99-101]
45	Eucalyptus Oil	<i>Eucalyptus globulus</i>	Myrtaceae	Fresh leaves	Eucalyptol, Geranyl acetate, Pinene, Cineole, Phellandrene, Citronellal, Camphene, Pinene	Expectorant, Antiseptics, Used for the treatment of bronchitis and cough	[1, 102-103]
46	Chenopodium Oil	<i>Chenopodium ambrosioides</i>	Chenopodiaceae	Fresh flower, leaves, roots and fruit	Ascaridole, Cymene, l-limonene, Myrcene, Camphor	Anthelmintics, Antifungal, Antioxidant, Anti-malarial, Antimicrobial, Anti-inflammatory	[1, 104-112]

V. CONCLUSION

Lots of chemical constituents are present in medicinal plants to show different pharmacological activities. Terpenoids are important constituents among all of constituents. Different types of terpenoids like Geraniol, Nerol, Phellandral, Manool, Lupeol, etc, are showing different roles like anthelmintics, antispasmodic, carminatives, antiseptics, expectorant, etc. The terpenoid plant name like Dill, Caraway, Cardamom, Fennel, Ajowan, Coriander, etc from different plant families like Umbelliferae, Liliaceae, Lauraceae, Zingiberaceae, Rutaceae, Myristicaceae, Labiatae, Myrtaceae, Santalaceae, Piperaceae, etc, are present in all over the world. Presently different research activities are continuing in this topic to show important and newer cum potent pharmacological activities from terpenoids. In near future, we hope that terpenoids may be a promising target for treating so many diseases and problems across the World.

Conflicts of Interest: Nil


REFERENCES

- [1] Kokate CK, Purohit AP, Gokhale SB. Pharmacognosy. Nirali Prakashan. 2012;47:1.1-1.104.
- [2] Evans WC. Trease and Evans Pharmacognosy. Saunders Elsevier. 2009;16:263-298.
- [3] Sunar P, Dasgupta RK, Roy SD, Shil D. A Review on Terpenoids. Indian Research Journal of Pharmacy and Science. 2023 March;10(1):2807-2816.
- [4] Graßmann J. Terpenoids as plant antioxidants. Vitamins & hormones. 2005 Jan 1;72:505-35.
- [5] Bisht BS, Bankoti H, Bharti T. A review on therapeutic uses of terpenoids. Journal of Drug Delivery and Therapeutics. 2021 Jan 2;11(1-s):182-5.
- [6] Urabe D, Asaba T, Inoue M. Convergent strategies in total syntheses of complex terpenoids. Chemical Reviews. 2015 Sep 9;115(17):9207-31.
- [7] Cheng AX, Lou YG, Mao YB, Lu S, Wang LJ, Chen XY. Plant terpenoids: biosynthesis and ecological functions. Journal of integrative plant biology. 2007 Feb;49(2):179-86.
- [8] Bohlmann J, Keeling CI. Terpenoid biomaterials. The Plant Journal. 2008 May;54(4):656-69.
- [9] Nadeem M, Muhammad Anjum F, Issa Khan M, Tehseen S, El-Ghorab A, Iqbal Sultan J. Nutritional and medicinal aspects of coriander (*Coriandrum sativum* L.) A review. British Food Journal. 2013 May 10;115(5):743-55.

- [10] Laribi B, Kouki K, M'Hamdi M, Bettaieb T. Coriander (*Coriandrum sativum* L.) and its bioactive constituents. *Fitoterapia*. 2015 Jun 1;103:9-26.
- [11] Huopalahti R, Lahtinen R, Hiltunen R, Laakso I. Studies on the essential oils of dill herb, *Anethum graveolens* L. *Flavour and fragrance journal*. 1988 Sep;3(3):121-5.
- [12] Brunke EJ, Hammerschmidt FJ, Koester FH, Mair P. Constituents of dill (*Anethum graveolens* L.) with sensory importance. *Journal of Essential Oil Research*. 1991 Jul 1;3(4):257-67.
- [13] Goyal S, Chaturvedi V, Dhingra G, Tanwar S, Sharma K, Singh S. *Trachyspermum ammi*: a review on traditional and modern pharmacological aspects. *Biological Sciences*. 2022 Nov 8;2(4):324-37.
- [14] Goyal S, Chaturvedi V, Dhingra G, Tanwar S, Sharma K, Singh S. *Trachyspermum ammi*: a review on traditional and modern pharmacological aspects. *Biological Sciences*. 2022 Nov 8;2(4):324-37.
- [15] Kooti W, Moradi M, Ali-Akbari S, Sharafi-Ahvazi N, Asadi-Samani M, Ashtary-Larky D. Therapeutic and pharmacological potential of *Foeniculum vulgare* Mill: a review. *Journal of HerbMed Pharmacology*. 2015 May 26;4(1):1-9.
- [16] Rahimi R, Ardekani MR. Medicinal properties of *Foeniculum vulgare* Mill. in traditional Iranian medicine and modern phytotherapy. *Chinese journal of integrative medicine*. 2013 Jan;19:73-9.
- [17] Nair TR, Kutty MG. Cardamom (*Elettaria cardamomum*) in Kerala, India. *Forest products, livelihoods and conservation*. 2004;1:130-47.
- [18] Govindarajan VS, Narasimhan S, Raghuvveer KG, Lewis YS, Stahl WH. Cardamom—Production, technology, chemistry, and quality. *Critical Reviews in Food Science & Nutrition*. 1982 Mar 1;16(3):229-326.
- [19] Miraj S, Kiani S. Pharmacological activities of *Carum carvi* L. *Der Pharmacia Lettre*. 2016;8(6):135-8.
- [20] Agrahari P, Singh DK. A review on the pharmacological aspects of *Carum carvi*. *J. Biol. Earth Sci*. 2014;4(1):1-3.
- [21] Ebrahimzadeh MA, Nabavi SM, Nabavi SF. Biological activities of *Mentha spicata* L. *Pharmacology online*. 2010;1:841-8.
- [22] Nour-Eddine B, Belkacem Z, Abdellah K. Experimental study and simulation of a solar dryer for spearmint leaves (*Mentha spicata*). *International Journal of Ambient Energy*. 2015 Mar 4;36(2):50-61.
- [23] Ashokkumar K, Simal-Gandara J, Murugan M, Dhanya MK, Pandian A. Nutmeg (*Myristica fragrans* Houtt.) essential oil: A review on its composition, biological, and pharmacological activities. *Phytotherapy Research*. 2022 Jul;36(7):2839-51.
- [24] Ha MT, Vu NK, Tran TH, Kim JA, Woo MH, Min BS. Phytochemical and pharmacological properties of *Myristica fragrans* Houtt.: an updated review. *Archives of Pharmacol Research*. 2020 Nov;43:1067-92.
- [25] Tepe AS, Tepe B. Traditional use, biological activity potential and toxicity of *Pimpinella* species. *Industrial Crops and Products*. 2015 Jul 1;69:153-66.
- [26] Romdhane M, Tizaoui C. The kinetic modelling of a steam distillation unit for the extraction of aniseed (*Pimpinella anisum*) essential oil. *Journal of Chemical Technology & Biotechnology: International Research in Process, Environmental & Clean Technology*. 2005 Jul;80(7):759-66.
- [27] Al-Asmari AK, Athar MT, Kadasah SG. An updated phytopharmacological review on medicinal plant of Arab region: *Apium graveolens* linn. *Pharmacognosy reviews*. 2017 Jan;11(21):13-18.
- [28] Takooree H, Aumeeruddy MZ, Rengasamy KR, Venugopala KN, Jeewon R, Zengin G, Mahomoodally MF. A systematic review on black pepper (*Piper nigrum* L.): from folk uses to pharmacological applications. *Critical reviews in food science and nutrition*. 2019 Jun 27;59(sup1):S210-43.
- [29] Meghwal M, Goswami TK. *Piper nigrum* and piperine: an update. *Phytotherapy Research*. 2013 Aug;27(8):1121-30.
- [30] Damanhoury ZA, Ahmad A. A review on therapeutic potential of *Piper nigrum* L. *Black Pepper*: The King of Spices. *Med. Aromat. Plants*. 2014;3(3):1-6.
- [31] Gohari AR, Saaidnia S. A review on phytochemistry of *Cuminum cyminum* seeds and its standards from field to market. *Pharmacognosy Journal*. 2011 Sep 1;3(25):1-5.
- [32] Mnif S, Aifa S. Cumin (*Cuminum cyminum* L.) from traditional uses to potential biomedical applications. *Chemistry & biodiversity*. 2015 May;12(5):733-42.
- [33] Bhadra P, Sethi L. A review paper on the Tulsi plant (*Ocimum sanctum*). *Indian Journal of Natural Sciences*. 2020;10(60):20854-60.
- [34] Gupta BM, Gupta R, Agarwal A, Goel S. *Ocimum Sanctum* (medicinal plant) research: A scientometric assessment of global publications output during 2008–17. *International Journal of Information Dissemination and Technology*. 2018;8(2):67-73.
- [35] Kumar A, Rahal A, Chakraborty S, Tiwari R, Latheef SK, Dhama K. *Ocimum sanctum* (Tulsi): a miracle herb and boon to medical science-A Review. *Int J Agron Plant Prod*. 2013;4(7):1580-9.
- [36] Pandey G, Madhuri S. Pharmacological activities of *Ocimum sanctum* (tulsi): a review. *Int J Pharm Sci Rev Res*. 2010 Nov;5(1):61-6.
- [37] Abubakar IB, Malami I, Yahaya Y, Sule SM. A review on the ethnomedicinal uses, phytochemistry and pharmacology of *Alpinia officinarum* Hance. *Journal of ethnopharmacology*. 2018 Oct 5;224:45-62.
- [38] Bitari A, Oualdi I, Touzani R, Elachouri M, Legssyer A. *Alpinia officinarum* Hance: A mini review. *Materials Today: Proceedings*. 2023 Jan 1;72:3869-74.
- [39] Rasool S, Maqbool M. An overview about *Hedychium spicatum*: a review. *Journal of Drug Delivery and Therapeutics*. 2019 Jan 2;9:476-80.
- [40] Rawat S, Jugran AK, Bhatt ID, Rawal RS. *Hedychium spicatum*: a systematic review on traditional uses, phytochemistry, pharmacology and future prospectus. *Journal of Pharmacy and Pharmacology*. 2018 Jun;70(6):687-712.
- [41] Sravani T, Paarakh PM. *Hedychium spicatum* Buch. Ham.-an overview. *Pharmacologyonline Newsletter*. 2011;2:633-42.
- [42] Singh M, Kumar R, Sharma S, Kumar L, Kumar S, Gupta G, Dua K, Kumar D. *Hedychium spicatum*: A comprehensive insight into its ethnobotany, phytochemistry, pharmacological and therapeutic attributes. *South African Journal of Botany*. 2023 Oct 1;161:638-47.
- [43] Pittler MH, Ernst E. Clinical effectiveness of garlic (*Allium sativum*). *Molecular nutrition & food research*. 2007 Nov;51(11):1382-5.
- [44] Lee JY, Gao Y. Review of the application of garlic, *Allium sativum*, in aquaculture. *Journal of the World Aquaculture Society*. 2012 Aug;43(4):447-58.
- [45] Kriplani P, Guarve K, Baghael US. *Arnica montana* L.–a plant of healing. *Journal of Pharmacy and Pharmacology*. 2017 Aug;69(8):925-45.
- [46] Bellavite P, Marzotto M, Bonafini C. *Arnica montana* experimental studies: confounders and biases?. *Journal of integrative medicine*. 2018 Mar 1;16(2):72-6.
- [47] Singh J, Arunima PS, Tiwari R, Byadgi PS. Standardization & Identification of *Jatamansi* (*Nardostachys jatamansi* DC.). *Korean Journal of Physiology and Pharmacology*. 2023;27(3):353-358.
- [48] Bhagat M, Pandita RM, Saxena AK. In vitro and in vivo biological activities of *Nardostachys jatamansi* roots. *Med Aromat Plants*. 2013;2(142):2167-0412.
- [49] Mallavarapu GR, Ramesh S. Essential oil of the fruits of *Cinnamomum zeylanicum* Blume. *Journal of Essential Oil Research*. 2000 Sep 1;12(5):628-30.
- [50] Saleem M, Bhatti HN, Jilani MI, Hanif MA. Bioanalytical evaluation of *Cinnamomum zeylanicum* essential oil. *Natural product research*. 2015 Oct 2;29(19):1857-9.

- [51] Ghai R, Nagarajan K, Singh J, Swarup S, Kesheri M. Evaluation of antioxidant status in-vitro and in-vivo in hydro-alcoholic extract of *Eugenia caryophyllus*. *Int J Pharmacol Toxicol*. 2016;4(1):19-24.
- [52] Rajkumar V, Berwal JS. Inhibitory effect of clove (*Eugenia caryophyllus*) on toxigenic molds. *JOURNAL OF FOOD SCIENCE AND TECHNOLOGY-MYSORE*. 2003;40(4):416-8.
- [53] Srivastava R, Ahmed H, Dixit RK, Saraf SA. *Crocus sativus* L.: a comprehensive review. *Pharmacognosy reviews*. 2010 Jul;4(8):200-208.
- [54] Kumar R, Singh V, Devi K, Sharma M, Singh MK, Ahuja PS. State of art of saffron (*Crocus sativus* L.) agronomy: A comprehensive review. *Food Reviews International*. 2008 Dec 22;25(1):44-85.
- [55] Shilpi JA, Taufiq-Ur-Rahman M, Uddin SJ, Alam MS, Sadhu SK, Seidel V. Preliminary pharmacological screening of *Bixa orellana* L. leaves. *Journal of Ethnopharmacology*. 2006 Nov 24;108(2):264-71.
- [56] Satyanarayana A, Prabhakara Rao PG, Rao DG. Chemistry, processing and toxicology of annatto (*Bixa orellana* L.). *J Food Sci Technol*. 2003 Mar 1;40(2):131-41.
- [57] Bos R, Woerdenbag HJ, Hendriks H, Smit HF, Wikström HV, Scheffer JJ. Composition of the Essential Oil from Roots and Rhizomes of *Valeriana wallichii* DC. *Flavour and Fragrance Journal*. 1997 Mar;12(2):123-31.
- [58] Kumar P. A review on medicinal plant *Piper cubeba* L. and its pharmaceutical properties. *Int J Sci Food Agric*. 2021;5:174-9.
- [59] Amara U, Khan A, Laraib S, Wali R, Sarwar U, Ain QT, Shakeel S. Conservation status and therapeutic potential of *Saussurea lappa*: An overview. *American Journal of Plant Sciences*. 2017 Feb 4;8(3):602-14.
- [60] Wei H, Yan LH, Feng WH, Ma GX, Peng Y, Wang ZM, Xiao PG. Research progress on active ingredients and pharmacologic properties of *Saussurea lappa*. *J Asian Nat Prod Res*. 2014;10(11):1045-53.
- [61] Balakumbahan R, Rajamani K, Kumaran K. *Acorus calamus*: An overview. *Journal of Medicinal Plants Research*. 2010 Dec 29;4(25):2740-5.
- [62] Muthulakshmi T, SALEH AM, KUMARI NV, Palanichamy V. *Acorus Calamus*. *Int. J. Drug Dev. & Res*. 2015 Jan;7(1):0975-9344.
- [63] Subha TS, Gnanamani A, Mandal AB. Pharmacognostic evaluation of *Acorus calamus* L. *Pharmacognosy Journal*. 2011 Jul 1;3(23):24-7.
- [64] Ernst E, Pittler MH. The efficacy and safety of feverfew (*Tanacetum parthenium* L.): an update of a systematic review. *Public health nutrition*. 2000 Dec;3(4a):509-14.
- [65] Hendricks H, Anderson-Wildeboer Y, Engels G, Bos R, Woerdenbag HJ. The content of parthenolide and its yield per plant during the growth of *Tanacetum parthenium*. *Planta medica*. 1997 Aug;63(04):356-9.
- [66] Brown AM, Lowe KC, Davey MR, Power JB. Feverfew (*Tanacetum parthenium* L.): tissue culture and parthenolide synthesis. *Plant Science*. 1996 May 3;116(2):223-32.
- [67] Hendriks H, Bos R, Woerdenbag HJ. The Essential Oil of *Tanacetum parthenium* (L.) Schultz-Bip. *Flavour and Fragrance Journal*. 1996 Nov;11(6):367-71.
- [68] Swamy KN, Rao SS. Effect of brassinosteroids on the performance of coleus (*Coleus forskohlii*). *Journal of herbs, spices & medicinal plants*. 2011 Mar 16;17(1):12-20.
- [69] Singh G, Ahmad M, Nazim M. A bibliometric study of *Embelia ribes*. *Library review*. 2008 Apr 18;57(4):289-97.
- [70] Dhavala A, Rathore TS. Micropropagation of *Embelia ribes* Burm f. through proliferation of adult plant axillary shoots. *In Vitro Cellular & Developmental Biology-Plant*. 2010 Apr;46:180-91.
- [71] Divya PJ, Jamuna P, Jyothi LA. Antioxidant properties of fresh and processed *Citrus aurantium* fruit. *Cogent Food & Agriculture*. 2016 Dec 31;2(1):1184119.
- [72] Hsouna AB, Hamdi N, Halima NB, Abdelkafi S. Characterization of essential oil from *Citrus aurantium* L. flowers: antimicrobial and antioxidant activities. *Journal of Oleo Science*. 2013;62(10):763-72.
- [73] Neves JA, Neves JA, Oliveira RD. Pharmacological and biotechnological advances with *Rosmarinus officinalis* L. Expert opinion on therapeutic patents. 2018 May 4;28(5):399-413.
- [74] Jahanian H, Kahkeshani N, Sanei-Dehkordi A, Isman MB, Saeedi M, Khanavi M. *Rosmarinus officinalis* as a natural insecticide: a review. *International Journal of Pest Management*. 2024 Oct 1;70(4):818-63.
- [75] Atti-Santos AC, Rossato M, Pauletti GF, Rota LD, Rech JC, Pansera MR, Agostini F, Serafini LA, Moyna P. Physico-chemical evaluation of *Rosmarinus officinalis* L. essential oils. *Brazilian archives of biology and technology*. 2005;48:1035-9.
- [76] Serin ZO, Ateş N, Cavunt A. Utilization of rosemary (*Rosmarinus officinalis* L.) stalks in pulp and paper production. *Turkish Journal of Forestry*. 2017 Dec 20;18(2):155-9.
- [77] Srivastava J, Kayastha S, Jamil S, Srivastava V. Environmental perspectives of *Vetiveria zizanioides* (L.) Nash. *Acta Physiologiae Plantarum*. 2008 Jul;30:413-7.
- [78] Luqman S, Srivastava S, Darokar MP, Khanuja SP. Detection of Antibacterial Activity in Spent Roots of Two Genotypes of Aromatic Grass *Vetiveria zizanioides*. *Pharmaceutical biology*. 2005 Jan 1;43(8):732-6.
- [79] Almaguer C, Schönberger C, Gastl M, Arendt EK, Becker T. *Humulus lupulus*—a story that begs to be told. A review. *Journal of the Institute of Brewing*. 2014 Oct;120(4):289-314.
- [80] Zanolli P, Zavatti M. Pharmacognostic and pharmacological profile of *Humulus lupulus* L. *Journal of ethnopharmacology*. 2008 Mar 28;116(3):383-96.
- [81] Korpeläinen H, Pietiläinen M. Hop (*Humulus lupulus* L.): Traditional and present use, and future potential. *Economic botany*. 2021 Dec;75(3):302-22.
- [82] Garcia LC. A Review of *Artemisia annua* L.: Its genetics, biochemical characteristics, and anti-malarial efficacy. *Int. J. Sci. Technol*. 2015 Feb;5(2):38-46.
- [83] Zhigzhitzhapova SV, Dylenova EP, Gulyaev SM, Randalova TE, Taraskin VV, Tykheev ZA, Radnaeva LD. Composition and antioxidant activity of the essential oil of *Artemisia annua* L. *Natural Product Research*. 2020 Sep 16;34(18):2668-71.
- [84] Sindhu RK, Upma KA, Arora S. *Santalum album* linn: a review on morphology, phytochemistry and pharmacological aspects. *International Journal of PharmTech Research*. 2010 Jan;2(1):914-9.
- [85] Ma GH, Bunn E, Zhang JF, Wu GJ. Evidence of Dichogamy in *Santalum album* L. *Journal of Integrative Plant Biology*. 2006 Mar;48(3):300-6.
- [86] Viswanath S, Purushothaman S. Sandal (*Santalum album* L.) conservation in southern India: a review of policies and their impacts. *Journal of Tropical Agriculture*. 2010;48(1-2):1-10.
- [87] Yogendra ND, Kumara RR, Prakash TA, Mohanty RP, Singh S, Prakhyath KM, Keerthi PE, Devanathan K, Pragadheesh VS, Sundaresan V, Verma RS. *Artemisia pallens* Wall. ex. DC: A comprehensive review. *Journal of Herbmmed Pharmacology*. 2024 Oct 1;13(4):501-22.

- [88] Suresh J, Singh A, Vasavi A, Ihsanullah M, Mary S. Phytochemical and pharmacological properties of *Artemisia pallens*. *International Journal of Pharmaceutical Sciences and Research*. 2011 Dec 1;2(12):3081-3090.
- [89] Kaur H, Bhardwaj U, Kaur R. *Cymbopogon nardus* essential oil: a comprehensive review on its chemistry and bioactivity. *Journal of Essential Oil Research*. 2021 May 4;33(3):205-20.
- [90] Diass K, Merzouki M, Elfazazi K, Azzouzi H, Challioui A, Azzaoui K, Hammouti B, Touzani R, Depeint F, Ayerdi Gotor A, Rhazi L. Essential oil of *lavandula officinalis*: Chemical composition and antibacterial activities. *Plants*. 2023 Apr 6;12(7):1571(1-16).
- [91] Sen S, Israr M, Singh S, Singh MK, Verma RS, Bawankule DU. Pharmaceutical, cosmeceutical, food additive and agricultural perspectives of *Cymbopogon martini*: A potential industrial aromatic crop. *South African Journal of Botany*. 2023 Jul 1;158:277-91.
- [92] Janbaz KH, Qayyum A, Saqib F, Imran I, Zia-Ul-Haq M, De Feo V. Bronchodilator, vasodilator and spasmolytic activities of *Cymbopogon martinii*. *J Physiol Pharmacol*. 2014 Dec 1;65(6):859-66.
- [93] Nikolić M, Marković T, Mojević M, Pejin B, Savić A, Perić T, Marković D, Stević T, Soković M. Chemical composition and biological activity of *Gaultheria procumbens* L. essential oil. *Industrial crops and products*. 2013 Aug 1;49:561-7.
- [94] Jintanasirinurak S, Manichart N, Somala N, Laosinwattana C, Teerarak M. Chemical analysis and bioactivities of *Gaultheria procumbens* L. essential oil: Herbicidal, antioxidant and antibacterial activities. *Current Applied Science and Technology*. 2023 Mar 29;23(5):1-13.
- [95] Trevisan SC, Menezes AP, Barbalho SM, Guiguer ÉL. Properties of *Mentha piperita*: a brief review. *World J Pharm Med Res*. 2017 Feb;3(1):309-13.
- [96] Dai J, Orsat V, Raghavan GV, Yaylayan V. Investigation of various factors for the extraction of peppermint (*Mentha piperita* L.) leaves. *Journal of food engineering*. 2010 Feb 1;96(4):540-3.
- [97] Tanu B, Harpreet K. Benefits of essential oil. *Journal of Chemical and Pharmaceutical Research*. 2016;8(6):143-9.
- [98] Van der Walt JJ, Demarne F. *Pelargonium graveolens* and *P. radens*: A comparison of their morphology and essential oils. *South African Journal of Botany*. 1988 Dec 1;54(6):617-22.
- [99] Hamidpour R, Hamidpour S, Hamidpour M, Shahleri M. Camphor (*Cinnamomum camphora*), a traditional remedy with the history of treating several diseases. *Int. J. Case Rep. Images*. 2013 Feb 1;4(2):86-9.
- [100] Malabadi RB, Kolkar KP, Meti NT, Chalannavar RK. Camphor tree, *Cinnamomum camphora* (L.); Ethnobotany and pharmacological updates. *Biomedicine*. 2021 Jul 7;41(2):181-4.
- [101] Lee HJ, Hyun EA, Yoon WJ, Kim BH, Rhee MH, Kang HK, Cho JY, Yoo ES. In vitro anti-inflammatory and anti-oxidative effects of *Cinnamomum camphora* extracts. *Journal of ethnopharmacology*. 2006 Jan 16;103(2):208-16.
- [102] Carrillo I, Vidal C, Elissetche JP, Mendonça RT. Wood anatomical and chemical properties related to the pulpability of *Eucalyptus globulus*: a review. *Southern Forests: a Journal of Forest Science*. 2018 Jan 1;80(1):1-8.
- [103] Arroja L, Dias AC, Capela I. The role of *Eucalyptus globulus* forest and products in carbon sequestration. *Climatic Change*. 2006 Jan;74(1):123-40.
- [104] Kumar R, Mishra AK, Dubey NK, Tripathi YB. Evaluation of *Chenopodium ambrosioides* oil as a potential source of antifungal, antiaflatoxigenic and antioxidant activity. *International journal of food microbiology*. 2007 Apr 30;115(2):159-64.
- [105] Jiménez-Osornio FJ, Kumamoto J, Wasser C. Allelopathic activity of *Chenopodium ambrosioides* L. *Biochemical Systematics and Ecology*. 1996 Apr 1;24(3):195-205.
- [106] Cysne DN, Fortes TS, Reis AS, de Paulo Ribeiro B, dos Santos Ferreira A, do Amaral FM, Guerra RN, Marinho CR, Nicolete R, Nascimento FR. Antimalarial potential of leaves of *Chenopodium ambrosioides* L. *Parasitology research*. 2016 Nov;115:4327-34.
- [107] Ajaib M, Hussain T, Farooq S, Ashiq M. Analysis of antimicrobial and antioxidant activities of *Chenopodium ambrosioides*: An ethnomedicinal plant. *Journal of Chemistry*. 2016;2016(1):4827157.
- [108] Ouadja B, Katawa G, Toudji GA, Layland L, Gbékley EH, Ritter M, Anani K, Ameyapoh Y, Karou SD. Anti-inflammatory, antibacterial and antioxidant activities of *Chenopodium ambrosioides* L. (*Chenopodiaceae*) extracts. *Journal of Applied Biosciences*. 2021 Jun 30;162(1):16764-94.
- [109] Holm T. *Chenopodium ambrosioides* L.; a morphological study. *American Journal of Science*. 1923 Aug 1;5(32):157-67.
- [110] Pandey AK, Singh P, Palni UT, Tripathi NN. Application of *Chenopodium ambrosioides* Linn. essential oil as botanical fungicide for the management of fungal deterioration in pulses. *Biological agriculture & horticulture*. 2013 Sep 1;29(3):197-208.
- [111] Wirth EH. A study of *Chenopodium ambrosioides* var. *anthelminticum* and its volatile oil. *Journal of the American Pharmaceutical Association*. 1920 Feb;9(2):127-41.
- [112] Shah H, Nisar M, Suhail M, Bacha N. Antimicrobial studies of the crude extracts from the roots of *Chenopodium ambrosioides* Linn. *African journal of microbiology research*. 2014;8(21):2099-104.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)