



IJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 13 **Issue:** V **Month of publication:** May 2025

DOI: <https://doi.org/10.22214/ijraset.2025.70294>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

The Contribution of Microplastics to Marine Pollution

M L Sharma¹, Sunil Kumar², Sunil Maggu³, Lakshya Goswami⁴, Riya Kalra⁵

^{1, 2, 3}Maharaja Agrasen Institute of Technology, Delhi

^{4, 5}Research Scholar, Maharaja Agrasen Institute of Technology, Delhi

Abstract: *The entry of microplastics into the world's oceans is now a critical global environmental concern. Microplastics, or man-made polymer particles with diameters of less than five millimeters, are the byproduct of a diverse array of human activities, ranging from industrial operations, consumer product use, to the breakdown of larger plastic materials through physical, chemical, and biological processes. The tiny plastic particles now cover all marine ecosystems — from shallow coastal waters to the deepest depths of the oceanic trenches. Their durability and tiny size make them easily consumed by a wide array of marine species, ranging from plankton, mollusks, and fish, to seabirds and marine mammals. Upon ingestion, microplastics can produce physical obstruction, internal injury, and false feelings of satiation, with grave impacts on the growth, reproductive success, and survival rates of marine species.*

Additionally, microplastics serve as carriers for a variety of toxic chemicals, such as persistent organic pollutants (POPs) and heavy metals, which adsorb on their surfaces and find their way into marine food chains. Through biomagnification and bioaccumulation, these toxic chemicals can be spread throughout the food chain, and have significant implications for human health, especially through the consumption of contaminated seafood. This research aims to provide a comprehensive analysis of microplastic sources and pathways, assess their multidimensional impacts on marine ecosystems and human populations, and review existing and emerging solutions to counter their prevalence. Countermeasures to counter microplastic pollution must be a concerted global effort with effective policy-making, technological innovation, improved waste management strategies, and mass public education campaigns. A comprehensive understanding of the role of microplastics in oceanic pollution is not only crucial for marine biodiversity conservation and the maintenance of ecological balance but also for the health, food security, and economic prosperity of future generations.

Keywords: *Microplastics, ocean pollution, marine environments, plastic pollution, environmental considerations, human health, have sustainable options, marine life, pollution, plastic decay*

I. INTRODUCTION

The rapid growth of plastics has characterized our age of industrialization by driving transformation across multiple sectors including packaging, construction, healthcare and travel; indeed, since the mid-20th century the production of plastic has increased many-fold, with now well over 400 million tons produced annually. Plastics are favorable due to their versatility, durability, lightweight and low costs for production, yet these same characteristics help contribute to plastic presence in the environment, where degradation takes much longer and dire future effects become apparent throughout ecosystems globally.

Within all types and forms of plastic pollution, microplastics, which are plastic fragments that are less than five mm in size, represent an almost-brilliant level of threat. Microplastics enter into marine ecosystems through two primary pathways: 1) as primary microplastics engineered at a microscopic scale for use in products such as cosmetics, cleaning products and industrial abrasives; and 2) as secondary plastic materials produced from the degradation of larger plastic material by the effects of sunlight, abrasion, even metabolism by microbes.

Once released into the marine environment, microplastics disperse very easily due to their size and buoyancy; almost anything in the water column consumes microplastics, and edible to both bottom and pelagic fishes, but also, fish that are larger and are therefore upper or mid-level predators.

Microplastics may not only alter biological functions in the ocean but also adversely affect the ecological health of oceans, biodiversity and ecosystem services that are also important to human wellbeing (e.g., fisheries and climate regulation). Despite increased recognition and awareness of the widespread sources of this pollution, and the resultant problems, the full extent of the issue is difficult to correctly quantify due to the difficulties of detection, monitoring and analysis.

Given the scale and complexity of microplastic pollution, it necessitates an immediate, coordinated global response. The intent of this paper is to provide a broad examination of the sources and pathways of microplastics, their environmental behaviours, their effects on marine organisms and human health, and to evaluate existing and possible means of mitigation and prevention. Identifying and understanding the many and varied challenges posed by microplastics is not only essential for ocean health but also in order to ensure a sustainable future for humanity.

II. SOURCE OF MICROPLASTIC

Microplastics enter the marine environment through many sources, and are broadly classified into two sources: primary and secondary microplastics - and understands their source is important in determining how to limit and reduce the release of microplastics.

A. Primary Microplastics

Primary microplastics are intentionally manufactured and sold for specifications of size ranging in a few micrometers to a few millimeters. Primary microplastics are manufactured as a known sizing for certain industrial and commercial use, and upon manufacturing and out of the hands of the manufacturer can enter the environment directly.

1) Personal Care Products

One of the largest contributors for our primary microplastics enter the environment is the cosmetics and personal care industry. The industry uses microbeads, or very small spherical plastic-balls, in cosmetics and personal care products such as facial scrubs, body washes, exfoliation products, toothpastes etc. The consumer rinses the product off during use, and its despite the size of the particle, microbeads are not cleaned out in the traditional wastewater treatment systems, instead enter rivers, lakes, and oceans.

2) Industrial

Industry uses microplastics as plastic pellets, also called nurdles, standard industry name to denote the pellet as the raw plastic product that becomes larger goods. During manufacturing, spillages during loading and transportation, and in the responsibility of the controller of the raw product before manufacturing it is possible for nurdles to enter the environment. Their small, light, and in some case buoyant nature allows them to be moved vertically via wind, rain, water, and we find them littering our environments and entering microplastic pollution.

3) Synthetic Textiles

Synthetic fibers include polyester, nylon, acrylic, and spandex. These fabrics shed microfibers in the wash. Each wash releases thousands of microscopic plastic fibers into the wastewater that are not easily trapped by wastewater treatment plants. Consequently, these microfibers then enter freshwater and marine ecosystems. Synthetic microfibers have become one of the most common forms of microplastic pollution worldwide.

B. Secondary Microplastics

These secondary microplastics are by-products from the process of breaking down existing larger plastic waste in the environment. The degradation takes that break down progressively the plastics into smaller particles.

1) Mechanical Interactions

Mechanical processes, including wave dynamics, sand abrasion, and impact with buoyant detritus, play a pivotal role in the physical degradation of plastic items. Plastic bags, bottles, fishing equipment, and packaging items become increasingly brittle and break apart into fragments over time as a result of the incessant pounding by waves and currents.

2) Chemical Decomposition

The exposure to sun rays, particularly ultraviolet (UV) radiation, triggers photo-degradation, a chemical process that weakens plastic polymers. UV radiation alters the molecular structure of plastics to make them more susceptible to degradation. The process is particularly prevalent on coastlines and on surface waters where plastics are subjected to direct sunlight.

3) Biological Mechanisms

Certain microorganisms such as bacteria and fungi are the agents responsible for the breakdown of plastics by their biological activity. Although the biodegradation rate of all plastics is much lower, they are capable of inducing small weakening and fragmenting of structure, thus giving rise to secondary microplastics on long time scales.

C. Common Examples

- 1) Secondary microplastics result from the disintegration of a variety of ubiquitous plastic items. These are:
- 2) Plastic Bags: Break into tiny film-like pieces.
- 3) Plastic bottles and packaging: Shatter into pieces and splinters.
- 4) Fishing lines and nets emit particles and fibers into the marine ecosystem. Packaging materials break down into foam-like granules and irregular fragments. Secondary microplastics are a distinct issue since they are constantly generated as bulk plastic waste remains in the environment. While primary microplastics develop sources that can be managed and addressed, secondary microplastics are generated by current pollution and thus are harder to manage and remove.

III. EFFECTS OF MICROPLASTICS

The presence of microplastics in marine ecosystems has far-reaching impacts, affecting the smallest marine animals and also humans and the ecosystem as a whole. Microplastics are not biologically inert and interact with living organisms and the physical environment in complex and often harmful ways. The impact of microplastics on marine life, human health, and ecological systems is discussed in this section.

A. Impact on Marine Life

Microplastics are ingested continuously by a wide range of marine animals, including plankton, mollusks, crustaceans, fish, seabirds, and marine mammals. Such extensive exposure causes direct and indirect biological effects.

1) Gastrointestinal Obstruction and Consumption

Marine species tend to misidentify microplastics as food due to their size, shape, color, and movement within waters. When ingested, microplastics can be lodged within gastrointestinal tracts, increasing physical obstructions. The condition produces a false sense of satiety, which reduces food intake and can therefore lead to malnourishment, reduced growth, decreased energy, and in extreme cases, death. Filter-feeding organisms like mussels and baleen whales are particularly vulnerable, as they unknowingly consume large amounts of water with microplastics.

2) Chemical Toxicity and Bioaccumulation

Microplastics possess a high surface-area-to-volume ratio and hydrophobic nature, which enable them to adsorb and concentrate poisonous chemicals from surrounding seawater. These encompass persistent organic pollutants (POPs) like PCBs, DDT, and heavy metals like lead and mercury. Upon consumption by aquatic animals, such contaminants may penetrate tissues and organs, impacting physiological processes and potentially reaching upper trophic levels through bioaccumulation and biomagnification. Such chemical transfer may interfere with metabolic processes, hormone regulation, and immune function.

3) Reproductive and Developmental Damage

Experiments have shown that microplastics have negative impacts on reproductive success in many marine species. In particular, studies using oysters and zebrafish have found reduced egg production, delayed larval development, and deformity of offspring following adult exposure to microplastics. These disruptions pose serious long-term threats to population viability and biodiversity, especially for species already at risk from overfishing or habitat loss.

B. Impacts on Human Health

As microplastics get incorporated into marine food webs, they are bound to reach human consumers through seafood and other marine products consumption. Exposure to these microplastics has raised major alarm regarding public health.

Food Chain Contamination Microplastics have been detected in a wide variety of foods including shellfish, fish, sea salt, and bottled and tap water. Since human beings eat these products regularly, the consumption of microplastics is now regarded as a normal phenomenon. Microplastic particles have been calculated to be consumed by an individual tens of thousands of times a year; however, the long-term health impact of this consumption is a subject of ongoing research.

1) Toxicological Health Hazards

The ingestion health risk of microplastics is not just caused by the particles but also by the types of chemicals they carry. Many microplastics contain or adsorb endocrine-disrupting chemicals (EDCs), carcinogens, and plasticizers such as phthalates and bisphenol A (BPA).

Such chemicals are implicated in developmental malformations, hormonal interference, cancer, and neurological effects in animal models. In addition, nanoplastics—microscopic particles that are smaller than microplastics—grow more and more concerning due to the potential of the tiny particles to traverse cell membranes and build up in tissues.

2) *Research Uncertainty and Gap*

While growing concern, the total health effects of microplastics are not yet fully understood. Current evidence is limited by heterogeneity of study design, lack of standardized sampling protocols, and ethical constraints in studying the long-term effect in human populations. However, the precautionary principle supports taking preventive action, as waiting for evidence to be definitive may result in irreversible damage.

C. *Ecological and Environmental Disturbance*

Beyond individual organisms, microplastics have the ability to alter the physical, chemical, and biological processes of entire ecosystems. Habitat Modification Microplastics settle in ocean sediments, particularly in estuaries, coastal environments, and deep-sea communities. Microplastics may alter sediment texture, porosity, and nutrient content, degrading habitat quality for bottom-dwelling organisms. Microplastics suspended in the water column may impact light penetration and gas exchange, modifying photosynthesis and oxygen levels essential for aquatic organisms. Spread of Non-Native Species and Pathogenic Agents Free-floating microplastics are a vector for the transport of invasive species, algae, and microbial pathogens, allowing them to travel long oceanic distances. These "plastic rafts" provide stable surfaces for colonization and can introduce non-native species into vulnerable ecosystems, where they can outcompete native flora and fauna. Similarly, pathogens that are attached to plastics can allow for the transmission of marine disease, affecting both wild and farmed animals. Finally, the worldwide impact of microplastics on marine life, human well-being, and ecosystem harmony is one of the Anthropocene's greatest challenges. The cumulative and interconnected character of the effects calls for a rapid, multi-disciplinary action to research, regulate, and restore.



IV. MITIGATION STRATEGIES AND SOLUTIONS

To tackle the intricate and compound issue of microplastic pollution needs to be addressed through an integrated and multi-faceted approach. Solutions need to come from concerted efforts by governments, industries, scientists, and citizens. This section discusses existing and evolving measures towards minimizing the discharge and occurrence of microplastics in seas through policy measures, technological innovations, better waste management, and public awareness.

A. Policy and Regulation

Stringent laws are at the heart of curbing production and distribution of microplastics. Governments have started paying attention to the threat and enforcing regulatory measures to cut plastic pollution at the source.

1) Bans on Microbeads

Arguably the most effective policy has been the blanket ban on microbeads in cleaning products. The United States, United Kingdom, Canada, and New Zealand have all enacted legislation to ban the sale and manufacture of cosmetics and cleaners containing microbeads. The ban significantly reduced the amount of primary microplastics entering water systems, demonstrating the effectiveness of targeted regulation in preventing pollution.

2) Extended Producer Responsibility (EPR)

The Extended Producer Responsibility principle assigns the waste disposal liability to producers and holds them responsible for the whole life of their products from the time of production, use to post-consumer disposal. Through EPR schemes, producers are forced to produce more recyclable products, fund recycling and collect used products. The application of EPR to plastic packaging and synthetic textiles would cause industries to adopt more eco-friendly production practices and decrease the production rate of microplastics.

3) International Agreements and Treaties

At a larger level, international coordination is required. The United Nations has paved the way for an internationally legally binding plastics treaty that would aim to address the full plastics life cycle from production to design to end-of-life waste disposal. Such a treaty would be able to coordinate efforts on a global scale, adopting similar standards to combat microplastics on a worldwide level.

B. Technological Innovations

Scientific and engineering advances bring hopeful prospects in ending microplastic contamination, particularly in designing filtration technologies and large-scale cleaning operations.

1) Filtration Technologies

Maybe the most effective way of reducing microplastic release from synthetic clothing is filtration. Future washing machine technologies include built-in microfiber filters and accessories like Guppyfriend bags and Cora Balls that trap fibers before they enter wastewater systems. Integration of more advanced filtration systems in municipal wastewater treatment plants can also help in stopping microplastics from entering natural water bodies.

2) Ocean Cleanup Initiatives

Several non-governmental organizations have initiated the removal of plastic pollution from the ocean. Notable among them is The Ocean Cleanup, which employs large-scale passive collection systems in ocean gyres—rotating zones in the ocean where plastics become concentrated. The systems are intended to capture and remove macro- and microplastics, dispersing the concentration of pollutants in open water.

3) Biodegradable Alternatives

Parallel to that, scientific research keeps developing biodegradable plastics and green commodities that break down faster in the natural world. While not the perfect solution, these technologies are a step in the direction of overcoming the long-term persistence of plastic pollution.

C. Improved Waste Management

Strong and effective waste management is significant in reducing plastic litter—and, as a consequence, microplastics—access into the environment.

1) Enhanced Recycling Plant

Most nations have limitations in recycling capability due to limited infrastructure, material contamination, and economics. Investment in sortation plants for materials, collection facilities, and establishment of local markets makes recycling efforts stronger. Standardization of plastics to improve the recyclability of materials is another where the process needs to be initiated.

2) Waste Reduction Campaigns

Governments and NGOs are encouraging reduction of waste that seeks to lower the use of plastic at the source. These strategies encourage the use of reusable cutlery, containers, and bags, and consumers are being encouraged to opt for plastic-free or low-waste products. Reducing the use of plastic in general is a major step towards preventing the formation of microplastics in the first place.

4.4 Behaviour Change and Public Education

Greater awareness and alteration of public attitude are key to the success of any long-term solution to the problem of microplastic pollution. Awareness Campaigns Education initiatives work to make individuals aware of microplastic's health and environmental effects. Through school education, media, and public service announcements, and also at community events, individuals are made aware of the source of microplastics and how they can be part of the solution. Awareness of the correlation between individual consumer behavior and ocean pollution has the potential to drive mass behavior change.

Behavioral Changes Towards Sustainability

Consumers have a major role to play in the generation of demand for sustainable goods. Promoting the adoption of green practices, such as the use of one-time plastics, reducing the use of packaging, purchasing sustainable products, and political support for the environment, can all combine to ease the strain on ecosystems. Social movements, such as zero-waste lifestyles, are also on the rise and are promoting cultural change around consumption. To sum up, notwithstanding the enormity of the problem of microplastic pollution, a combined package of regulatory response, technological measures, improved infrastructure, and informed public participation provides an implementable pathway forward. The success of these measures depends on cross-sectoral action and global commitment to sustainable development.

V. DISCUSSION

Worldwide, microplastic pollution is causing widespread concern and discussion, but action on the issue presents a very real and serious challenge. This is largely due to the fact that microplastics differ greatly from larger plastic debris and are more challenging to monitor, identify, document, and retrieve. Microplastics are commonly small, measurable at less than 5 millimetres and typically invisible to the naked eye, which allows them to disperse thousands of miles across environmental barriers; they can float on ocean currents or be transported by atmospheric weather patterns. Microplastics have been retrieved from remote marine areas, from Arctic sea ice, to the Mariana Trench, which demonstrates the difficulty with the global scale contamination.

A. The Platform for Source Prevention

Source prevention is the best way to prevent microplastic pollution. Source prevention includes reducing plastic production (especially single-use plastics) and using biodegradable and eco-friendly materials. Source prevention matters because once microplastics are introduced to marine environments, there is no practical way to remove them without disturbing aquatic life and the ecosystem. The focus must be on redesigning products, including packaging and consumption, using the opportunities in the circular economy, and transitioning society into an experimental stage of adopting new sustainable alternatives.

B. Limitations of Clean Up Technologies

While ocean clean-up technologies and methods (floating barriers, plastic interceptors, and concentrated waste collection, etc.) does provide some hope, in reality, the limitations of present cleanup technologies are considerable. Cleanup technologies and methods generally only affect larger debris (i.e., macroplastics) and while they may be managed by cleanup methods, microplastics eventually*** end up in the ocean and urban canals.

VI. CONCLUSION

Microplastics are a major and growing risk to marine ecosystems, global biodiversity, and human health. Plastic production continues to rise, with the yearly total exceeding 400 million tons worldwide. Microplastics have created pollution that has entered almost all of the world's oceans because plastic is resistant to degradation and fragmentation.

Microplastics may be tiny and often not seen by the human eye, but they have disproportionately large effects on marine organisms, including disrupting feeding, reproduction, and physiological functions. Now, there are emerging risks to human health from bioaccumulation in seafood and drinking water supplies.

The reach of microplastics and their known resistance to breakdown highlights the need for a global coordinated course of action. Failing to act now will result in impacts that may be irreversible for critical ecosystems like coral reefs, seagrass beds, and the deep-sea environment. Not to mention the existential risks posed as microplastics enter the food chain and humans consume these pollutants with toxic additives, along with chemical and microbial transfer processes.

Microplastic pollution needs to be addressed at multiple levels, via a holistic approach. Key to any management approach must be source prevention with sustainable reductions in plastic production, eliminating unnecessary single-use products, and enabling the transition to biodegradable alternatives, including bio-based materials. In addition, technological Innovation—through improved filtration technology, waste management systems, and ocean cleanup efforts—needs to be supported and scaled. Nationally and internationally, legislative frameworks and policy will need to hold car manufacturers accountable, regulate plastic use, and incentivize a circular economy. In addition to technical innovations, public education and behavioral change can influence consumer-driven demand for plastics, and encourage participation in sustainable behaviours.

REFERENCES

- [1] Andrady, A. L. (2011). Microplastics in the marine environment. *Marine Pollution Bulletin*, 62(8), 1596–1605. <https://doi.org/10.1016/j.marpolbul.2011.05.030>
- [2] Cole, M., Lindeque, P., Halsband, C., & Galloway, T. S. (2011). Microplastics as contaminants in the marine environment: A review. *Marine Pollution Bulletin*, 62(12), 2588–2597. <https://doi.org/10.1016/j.marpolbul.2011.09.025>
- [3] GESAMP. (2015). Sources, fate and effects of microplastics in the marine environment: A global assessment (Kershaw, P. J., Ed.; Reports and Studies No. 90). IMO/FAO/UNESCO-IOC/UNIDO/WMO/IAEA/UN/UNEP Joint Group of Experts on the Scientific Aspects of Marine Environmental Protection. <https://www.gesamp.org/publications/microplastics-in-the-marine-environment>
- [4] Browne, M. A., Galloway, T. S., & Thompson, R. C. (2010). Spatial patterns of plastic debris along beaches in the North Atlantic Ocean. *Marine Environmental Research*, 70(3), 257–263. <https://doi.org/10.1016/j.marenvres.2010.05.007>
- [5] Jambeck, J. R., Geyer, R., Wilcox, C., Siegler, T. R., Perryman, M., Andrady, A., & Law, K. L. (2015). Plastic waste inputs from land into the ocean. *Science*, 347(6223), 768–771. <https://doi.org/10.1126/science.1260352>
- [6] UNEP. (2018). Single-use plastics: A roadmap for sustainability. United Nations Environment Programme. <https://www.unep.org/resources/report/single-use-plastics-roadmap-sustainability>
- [7] Rochman, C. M., Hoh, E., Kaye, S., & Sivan, A. (2013). Ingested plastic transfers hazardous chemicals to fish and induces hepatic stress. *Scientific Reports*, 3, 3263. <https://doi.org/10.1038/srep03263>



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)