



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 5 Issue: IX Month of publication: September 2017

DOI: <http://doi.org/10.22214/ijraset.2017.9195>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Certain Summation and Transformation Formulae for Basic Hypergeometric Series

Rajesh Pandey

Maharishi University of Information Technology Lucknow 226013, India.

Abstract: In this paper, making use of certain known summation formulae an attempt has been made to establish certain very interesting summation and transformation formulae for basic hypergeometric series.

Keywords: Hypergeometric functions, Summation, Transformation, Polybasic, Converges

I. INTRODUCTION

In 1972, Verma [2] established the following very general transformation formulae:

$$\sum_{n=0}^{\infty} A_n B_n \frac{(x\omega)^n}{(q; q)_n} = \sum_{n=0}^{\infty} \frac{(-x)^n q^{\binom{n}{2}}}{(q, \gamma q^n; q)_n} \sum_{k=0}^{\infty} \frac{(\alpha, \beta; q)_{n+k} B_{n+k} x^k}{(q; q)_k (\gamma q^{2n+1}; q)_k} \sum_{j=0}^n \frac{(q^{-n}, \gamma q^n; q)_j A_j (\omega q)^j}{(q, \alpha, \beta; q)_j} \quad (1.1)$$

Making use of certain known summation formulae due to Verma and Jain [4] and the expansion formulae (1.1), an attempt has been made to establish certain very interesting summation and transformation formulae for basic hypergeometric series.

II. DEFINITIONS AND NOTATIONS

The following result will be required in our analysis:-

$${}_2\Phi_1 \left[\begin{matrix} a, b; q; \frac{c}{ab} \\ cq \end{matrix} \right] = \frac{\left(\frac{cq}{a}, \frac{cq}{b}; q \right)_{\infty}}{\left(cq, \frac{cq}{ab}; q \right)_{\infty}} \left\{ \frac{ab(1+c) - (a+b)c}{ab-c} \right\} \quad (2.1)$$

$${}_4\Phi_3 \left[\begin{matrix} q^{-n}, x^2 y^2 q^{1+n}, x, -xq; q; q \\ xyq, -xyq, x^2 q \end{matrix} \right] = \frac{x^n (q; q)_n (x^2 q^2; q^2)_m (y^2 q^2; q^2)_m}{(x^2 q; q)_n (x^2 y^2 q^2; q^2)_m (q^2; q^2)_m} \quad (2.2)$$

Where m is the greatest integer $\leq n/2$

$${}_3\Phi_2 \left[\begin{matrix} q^{-n}, x^2 q^{1+n}, 0; q; q \\ xy, -xy \end{matrix} \right] = \begin{cases} 0 & \text{if } n \text{ is odd} \\ (-1)^n q^{n(n+1)} x^{2n} \frac{(q; q^2)_n}{(x^2 q^2; q^2)_n} & \text{if } n \text{ is even} \end{cases} \quad (2.3)$$

$${}_5\Phi_4 \left[\begin{matrix} x, aq^{1+n}, \left(\frac{aq}{b} \right)^{1/2}, -\left(\frac{aq}{b} \right)^{1/2}, q^{-n}; q; q \\ (aq)^{1/2}, -(aq)^{1/2}, \frac{aq}{b}, xq \end{matrix} \right] = \frac{x^n (q; q)_n (aq/x; q)_n}{(aq; q)_n (xq; q)_n}$$

$${}_6\Phi_5 \left[\begin{matrix} a, q^2 \sqrt{a}, -q^2 \sqrt{a}, b, x, xq; q^2; \frac{aq}{bx^2} \\ \sqrt{a}, -\sqrt{a}, \frac{aq^2}{b}, \frac{aq^2}{x}, \frac{aq}{x} \end{matrix} \right] \text{ to } (n+1) \text{ terms,} \quad (2.4)$$

Where m is the greatest integer $< n/2$

$${}_4\Phi_3 \left[\begin{matrix} x, -xq, bx^2 q^{2+n}, q^{-n}; q; q \\ x^2 q^2, xq\sqrt{b}, -xq\sqrt{b} \end{matrix} \right] = \frac{x^n (q; q)_n (bxq^2; q)_n (bx^2 q^3; q^2)_m (bq^2; q^2)_m (xq^2; q)_{2m}}{(xq; q)_n (bx^2 q^2; q)_n (q^2; q^2)_m (x^2 q^3; q^2)_m (bxq^2; q)_{2m}} \quad (2.5)$$

Where m is the greatest integer $< n/2$

$${}_5\Phi_4 \left[\begin{matrix} x, aq^{1+n}, \left(\frac{x}{q}\right)^{1/2}, -\left(\frac{x}{q}\right)^{1/2}, q^{-n}; q; q \\ (aq)^{1/2}, -(aq)^{1/2}, \frac{x}{q}, xq \end{matrix} \right] = \frac{x^{n-m} \left(\frac{aq}{x}; q\right)_n (q; q)_n (aq^2; q^2)_m (xq; q^2)_m}{q^m (aq; q)_n (xq; q)_n (q^2; q^2)_m \left(\frac{aq}{x}; q^2\right)_m} \quad (2.6)$$

Where m is the greatest integer $< n / 2$

III.MAIN RESULTS

In this section we shall establish our main results :

$${}_4\Phi_3 \left[\begin{matrix} \alpha, \beta, x, -xq; q, \frac{x^2y^2q}{\alpha\beta} \\ xyq, -xyq, x^2q \end{matrix} \right] = \frac{\left(\frac{x^2y^2}{\alpha}q^2, \frac{x^2y^2}{\beta}q^2; q\right)_\infty}{\left(x^2y^2q^2, \frac{x^2y^2}{\alpha\beta}q^2; q\right)_\infty} \sum_{n=0}^{\infty} \frac{(\alpha, \beta; q)_n (x^2y^2q^2; q)_{2n} (x^2q^2, y^2q^2; q^2)_m}{\left(x^2y^2q^{n+1}, \frac{x^2y^2}{\alpha}q^2, \frac{x^2y^2}{\beta}q^2, x^2q; q\right)_n (x^2y^2q^2, q^2; q^2)_m} \\ \times \left(-\frac{x^3y^2}{\alpha\beta}\right)^n q^{n(n+1)/2} \left\{ \frac{\alpha\beta(1+x^2y^2q^{2n+1}) - x^2y^2q^{n+1}(\alpha+\beta)}{\alpha\beta - x^2y^2q} \right\} \quad (3.1)$$

Where m is the greatest integer $\leq n / 2$

$${}_3\Phi_2 \left[\begin{matrix} \alpha, \beta, 0; q; \frac{x^2q}{\alpha\beta} \\ xq, -xq \end{matrix} \right] = \frac{\left(\frac{x^2}{\alpha}q^2, \frac{x^2}{\beta}q^2; q\right)_\infty}{\left(x^2q^2, \frac{x^2}{\alpha\beta}q^2; q\right)_\infty} \sum_{n=0}^{\infty} \frac{(\alpha, \beta; q)_n (x^2q^2; q)_{2n}}{\left(q, x^2q^{n+1}, \frac{x^2}{\alpha}q^2, \frac{x^2}{\beta}q^2; q\right)_n} \left(-\frac{x^2}{\alpha\beta}\right)^n q^{n(n+1)/2} \\ \left\{ \frac{\alpha\beta(1+x^2q^{2n+1}) - x^2q^{n+1}(\alpha+\beta)}{\alpha\beta - x^2q} \right\} \times \begin{cases} 0, & \text{if } n \text{ is odd} \\ (-1)^n q^{n(n+1)} \frac{x^{2n}(q; q^2)_n}{(x^2q^2; q^2)_n}, & \text{if } n \text{ is even} \end{cases} \quad (3.2)$$

$${}_5\Phi_4 \left[\begin{matrix} \alpha, \beta, x \left(\frac{aq}{b}\right)^{1/2}, -\left(\frac{aq}{b}\right)^{1/2}; q; \frac{aq}{\alpha\beta} \\ (aq)^{1/2}, -(aq)^{1/2}, \frac{aq}{b}, xq \end{matrix} \right] = \frac{\left(\frac{a}{\alpha}q^2, \frac{a}{\beta}q^2; q\right)_\infty}{\left(aq^2, \frac{a}{\alpha\beta}q^2; q\right)_\infty} \\ \times \sum_{n=0}^{\infty} \frac{\left(\alpha, \beta, \frac{aq}{x}; q\right)_n (aq^2; q)_{2n}}{\left(aq^{n+1}, \frac{a}{\alpha}q^2, \frac{a}{\beta}q^2, aq, xq; q\right)_n} \left(-\frac{ax}{\alpha\beta}\right)^n q^{n(n+1)/2} \left\{ \frac{\alpha\beta(1+aq^{2n+1}) - aq^{n+1}(\alpha+\beta)}{\alpha\beta - aq} \right\} \\ \times {}_6\Phi_5 \left[\begin{matrix} a, q^2\sqrt{a}, -q^2\sqrt{a}, b, x, xq; q^2; \frac{aq}{bx^2} \\ \sqrt{a}, -\sqrt{a}, \frac{aq^2}{b}, \frac{aq^2}{x}, \frac{aq}{x} \end{matrix} \right] \text{ to } (m+1) \text{ terms,} \quad (3.3)$$

Where m is the greatest integer $\leq n / 2$

$${}_4\Phi_3 \left[\begin{matrix} \alpha, \beta, x, -xq; q; \frac{bx^2q^2}{\alpha\beta} \\ x^2q^2, xq\sqrt{b}, -xq\sqrt{b} \end{matrix} \right] = \frac{\left(\frac{b}{\alpha}x^2q^3, \frac{b}{\beta}x^2q^3; q\right)_\infty}{\left(bx^2q^3, \frac{b}{\alpha\beta}x^2q^3; q\right)_\infty} \sum_{n=0}^{\infty} \frac{(\alpha, \beta, bxq^2; q)_n (bx^2q^3; q)_{2n}}{\left(bx^2q^{n+2}, \frac{b}{\alpha}x^2q^3, \frac{b}{\beta}x^2q^3, bx^2q^2, xq; q\right)_n}$$

$$\times \frac{(bx^2q^3, bq^2; q^2)_m (xq^2; q)_{2m}}{(x^2q^3, q^2; q^2)_m (bxq^2; q)_{2m}} \left(\frac{-bx^3}{\alpha\beta} \right) q^{n(n+3)/2} \times \left\{ \frac{\alpha\beta(1+bx^2q^{2n+2}) - bx^2q^{n+2}(\alpha+\beta)}{\alpha\beta - bx^2q^2} \right\} \quad (3.4)$$

Where m is the greatest integer $\leq n/2$

$${}_5\Phi_4 \left[\begin{matrix} \alpha, \beta, x, \sqrt{\frac{x}{q}}, -\sqrt{\frac{x}{q}}; q; \frac{aq}{\alpha\beta} \\ \sqrt{aq}, -\sqrt{aq}, \frac{x}{q}, xq \end{matrix} \right] = \frac{\left(\frac{a}{\alpha}q^2, \frac{a}{\beta}q^2; q \right)_\infty}{\left(aq^2, \frac{a}{\alpha\beta}q^2; q \right)_\infty} \times$$

$$\times \sum_{n=0}^{\infty} \frac{\left(\alpha, \beta, \frac{aq}{x}; q \right)_n (aq^2; q)_{2n} \left(-\frac{ax}{\alpha\beta} \right)^n (xq)^{-m} q^{n(n+1)/2}}{\left(aq^{n+1} \frac{a}{\alpha}q^2, \frac{a}{\beta}q^2, aq, xq; q \right)_n} \times$$

$$\times \frac{(aq^2, xq; q^2)_m}{\left(\frac{aq}{x}, q^2, q^2 \right)_m} \left\{ \frac{\alpha\beta(1+aq^{2n+1}) - aq^{n+1}(\alpha+\beta)}{\alpha\beta - aq} \right\} \quad (3.5)$$

Where m is the greatest integer $\leq n/2$.

$${}_6\Phi_5 \left[\begin{matrix} \alpha, \beta, x, a^{1/3}, ba^{1/3}, b^2a^{1/3}; q; \frac{aq}{\alpha\beta} \\ \sqrt{a}, -\sqrt{a}, \sqrt{aq}, -\sqrt{aq}, xq \end{matrix} \right] = \frac{\left(\frac{a}{\alpha}q^2, \frac{a}{\beta}q^2; q \right)_\infty}{\left(aq^2, \frac{a}{\alpha\beta}q^2; q \right)_\infty} \times$$

$$\times \sum_{n=0}^{\infty} \frac{\left(\alpha, \beta, \frac{aq}{x}; q \right)_n (aq^2; q)_{2n} \left(-\frac{ax}{\alpha\beta} \right)^n q^{n(n+1)/2}}{\left(aq^{n+1} \frac{a}{\alpha}q^2, \frac{a}{\beta}q^2, aq, xq; q \right)_n} \times$$

$$\times \left\{ \frac{\alpha\beta(1+aq^{2n+1}) - aq^{n+1}(\alpha+\beta)}{\alpha\beta - aq} \right\}$$

$$\times {}_6\Phi_5 \left[\begin{matrix} a, q^3\sqrt{a}, -q^3\sqrt{a}, x, xq, xq^2; q^3; \frac{a}{x^3} \\ \sqrt{a}, -\sqrt{a}, \frac{aq^3}{b}, \frac{aq^2}{x}, \frac{aq}{x} \end{matrix} \right] \text{ to } (m+1) \text{ terms,} \quad (3.6)$$

Where m is the greatest integer $\leq n/3$.

Proof of (3.1) - (3.6)

In this section we shall give the outline of the proof of (3.1) – (3.6)

1) In order to prove (3.1), let us suppose that

$$A_n = \frac{(\alpha, \beta, x, -xq; q)_n}{(xyq, -xyq, x^2q; q)_n} (\omega)^n, B_n = \left(\frac{xy^2q}{\alpha\beta} \right)^n \text{ and } \gamma = x^2y^2q$$

In (1.1), we get;

$${}_4\Phi_3 \left[\begin{matrix} \alpha, \beta, x, -xq; q; \frac{x^2y^2q}{\alpha\beta} \\ xyq, -xyq, x^2q \end{matrix} \right] = {}_2\Phi_1 \left[\begin{matrix} \alpha q^n, \beta q^n; q, \frac{x^2y^2q}{\alpha\beta} \\ x^2y^2q^{2n+2} \end{matrix} \right] \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} (\alpha, \beta; q)_n}{(q, x^2y^2q^{n+1}; q)_n} \left(-\frac{x^2y^2q}{\alpha\beta} \right)^n \times$$

$$\times {}_4\Phi_3 \left[\begin{matrix} q^{-n}, x^2 y^2 q^{1+n}, x, -xq; q; q \\ xyq, -xyq, x^2 q \end{matrix} \right]$$

Now making use of (2.1) and (2.2), we get (3.1) after some simplifications.

2) In order to prove (3.2), let us suppose that

$$A_n = \frac{(\alpha, \beta, 0; q)_n}{(xq, -xq; q)_n (\omega)^n}, B_n = \left(\frac{xq}{\alpha\beta} \right)^n \text{ and } \gamma = x^2 q$$

In (1.1), we get;

$${}_3\Phi_2 \left[\begin{matrix} \alpha, \beta, 0; q; \frac{x^2 q}{\alpha\beta} \\ xq, -xq \end{matrix} \right] = \sum_{n=0}^{\infty} q^{\binom{n}{2}} (\alpha, \beta; q)_n \left(-\frac{x^2 q}{\alpha\beta} \right)^n \times {}_2\Phi_1 \left[\begin{matrix} \alpha q^n, \beta q^n; q, \frac{x^2 q}{\alpha\beta} \\ x^2 q^{2n+2} \end{matrix} \right] \times {}_3\Phi_2 \left[\begin{matrix} q^{-n}, x^2 q^{n+1}, 0; q; q \\ xq, -xq \end{matrix} \right]$$

Now making use of (2.1) and (2.3), we get (3.2) after some simplifications.

3) In order to prove (3.3), let us suppose that

$$A_n = \frac{(\alpha, \beta, x, \sqrt{\frac{aq}{b}}; -\sqrt{\frac{aq}{b}}; q)_n}{(\sqrt{aq}, -\sqrt{aq}, \frac{aq}{b}, xq; q)_n (\omega)^n}, B_n = \left(\frac{aq}{x\alpha\beta} \right)^n \text{ and } \gamma = aq$$

In (1.1), we get ;

$${}_5\Phi_4 \left[\begin{matrix} \alpha, \beta, x, \sqrt{\frac{aq}{b}}, -\sqrt{\frac{aq}{b}}; q; \frac{aq}{\alpha\beta} \\ \sqrt{aq}, -\sqrt{aq}, \frac{aq}{b}, xq \end{matrix} \right] = \sum_{n=0}^{\infty} q^{\binom{n}{2}} (\alpha, \beta; q)_n \left(-\frac{aq}{\alpha\beta} \right)^n \times {}_2\Phi_1 \left[\begin{matrix} \alpha q^n, \beta q^n; q, \frac{aq}{\alpha\beta} \\ aq^{2n+2} \end{matrix} \right] \times$$

$$\times {}_5\Phi_4 \left[\begin{matrix} x, aq^{n+1}, \sqrt{\frac{aq}{b}}, -\sqrt{\frac{aq}{b}}, q^{-n}; q; q \\ \sqrt{aq}, -\sqrt{aq}, \frac{aq}{b}, xq \end{matrix} \right]$$

Now making use of (2.1) and (2.4), we get (3.3) after some simplifications.

4) In order to prove (3.4), let us suppose that

$$A_n = \frac{(\alpha, \beta, x, -xq; q)_n}{(x^2 q^2, xq\sqrt{b} - xq\sqrt{b}; q)_n (\omega)^n}, B_n = \left(\frac{bxq^2}{\alpha\beta} \right)^n \text{ and } \gamma = bx^2 q^2$$

In (1.1), we get;

$${}_4\Phi_3 \left[\begin{matrix} \alpha, \beta, x, -xq; q; \frac{bx^2 q^2}{\alpha\beta} \\ x^2 q^2, xq\sqrt{b}, -xq\sqrt{b} \end{matrix} \right] = \sum_{n=0}^{\infty} q^{\binom{n}{2}} (\alpha, \beta; q)_n \left(-\frac{bx^2 q^2}{\alpha\beta} \right)^n \times {}_2\Phi_1 \left[\begin{matrix} \alpha q^n, \beta q^n; q, \frac{bx^2 q^2}{\alpha\beta} \\ bx^2 q^{2n+3} \end{matrix} \right] \times$$

$$\times {}_4\Phi_3 \left[\begin{matrix} x, -xq, bx^2 q^{2+n}, q^{-n}; q; q \\ x^2 q^2, xq\sqrt{b}, -xq\sqrt{b} \end{matrix} \right]$$

Now making use of (2.1) and (2.5), we get (3.4) after some simplifications.

5) In order to prove (3.5), let us suppose that

$$A_n = \frac{(\alpha, \beta, x, \sqrt{\frac{x}{q}}; -\sqrt{\frac{x}{q}}; q)_n}{(\sqrt{aq}, -\sqrt{aq}, \frac{x}{q}, xq; q)_n (\omega)^n}, B_n = \left(\frac{aq}{x\alpha\beta} \right)^n \text{ and } \gamma = aq$$

In (1.1), we get;

$${}_5\Phi_4 \left[\begin{matrix} \alpha, \beta, x, \sqrt{\frac{x}{q}}, -\sqrt{\frac{x}{q}}; q; \frac{aq}{\alpha\beta} \\ \sqrt{aq}, -\sqrt{aq}, \frac{x}{q}, xq \end{matrix} \right] = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} (\alpha, \beta; q)_n \left(-\frac{aq}{\alpha\beta}\right)^n}{(q, aq^{n+1}; q)_n} {}_2\Phi_1 \left[\begin{matrix} \alpha q^n, \beta q^n; q; \frac{aq}{\alpha\beta} \\ aq^{2n+2} \end{matrix} \right] \times \\
 \times {}_5\Phi_4 \left[\begin{matrix} x, aq^{n+1}, \sqrt{\frac{x}{q}}, -\sqrt{\frac{x}{q}}, q^{-n}; q; q \\ \sqrt{aq}, -\sqrt{aq}, \frac{x}{q}, xq \end{matrix} \right]$$

Now making use of (2.1) and (2.6), we get (3.5) after some simplifications.

6) In order to prove (3.6), let us suppose that

$$A_n = \frac{(\alpha, \beta, x, a^{1/3}, ba^{1/3}; q)_n}{(\sqrt{a}, -\sqrt{a}, \sqrt{aq}, -\sqrt{aq}, xq; q)_n} (\omega)^n, B_n = \left(\frac{aq}{x\alpha\beta}\right)^n \text{ and } \gamma = aq$$

In (1.1), we get ;

$${}_6\Phi_5 \left[\begin{matrix} \alpha, \beta, x, a^{1/3}, ba^{1/3}, b^2 a^{1/3}; q; \frac{aq}{\alpha\beta} \\ \sqrt{a}, -\sqrt{a}, \sqrt{aq}, -\sqrt{aq}, xq \end{matrix} \right] = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} (\alpha, \beta; q)_n \left(-\frac{aq}{\alpha\beta}\right)^n}{(q, aq^{n+1}; q)_n} {}_2\Phi_1 \left[\begin{matrix} \alpha q^n, \beta q^n; q; \frac{aq}{\alpha\beta} \\ aq^{2n+2} \end{matrix} \right] \times \\
 \times {}_6\Phi_5 \left[\begin{matrix} a^{1/3}, ba^{1/3}, b^2 a^{1/3}, x, aq^{n+1}, q^{-n}; q; q \\ \sqrt{a}, -\sqrt{a}, \sqrt{aq}, -\sqrt{aq}, xq \end{matrix} \right].$$

Now making use of (2.1) and (2.7), we get (3.6) after some simplifications.

IV. CONCLUSION

In this paper, an attempt has been made to establish six certain transformation formulae for basic hypergeometric series by making use of the identity (1.1)

V. ACKNOWLEDGEMENT

My thanks are due to Dr. G.C Chaubey Ex Associate Professor & Head department of Mathematics TDPG College Jaunpur and Professor B. Kunwar Department of Mathematics IET, Lucknow for their encouragement and for providing necessary support. I am extremely grateful for their constructive support.

REFERENCES

- [1] Slater, L.J.: Generalized Hypergeometric Functions, Cambridge University Press, London (1996)
- [2] Verma, A : Some transformation of series with arbitrary terms, Institute Lambrado (Rend. Sc.), A 106 (1972), 342-353.
- [3] Verma, A. : On identities of Rogers-Ramanujan type, Indian J. Pure and Appl. Math, 11(6), (1980), 770-90.
- [4] Verma, A. and Jain V.K. : Some summation formulae of basic hypergeometric series, Indian J. Pure and Appl. Math, 11(8), (1980), 1021-38.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)