



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 5 Issue: X Month of publication: October 2017 DOI: http://doi.org/10.22214/ijraset.2017.10026

www.ijraset.com

Call: 🕥 08813907089 🔰 E-mail ID: ijraset@gmail.com



# **Biological Monitoring Of Heavy Metals in Human**

G.Manimegalai<sup>1</sup> and Dr.A.Thaminum Ansari<sup>2</sup>

<sup>1</sup> Reasearch Scholar, Research and Development Centre, Bharathiyar University, Coimbatore, Tamilnadu. <sup>2</sup>Assistant Professor, Muthurangam Govt. Arts College, Vellore, Tamilnadu

Abstract: Air pollution in India is mainly caused by three sources, namely, vehicles, industrial and domestic sources. The objective of this paper is to provide an analysis of particulate metal emissions. The aim of this study was to investigate blood and urine levels of some heavy metals such as of chromium (Cr) and lead (Pb) in workers. These results have indicated that further studies are warranted for investigation of possible roles of heavy metals in industry workers. The majority of airborne metal emissions originate from metal industries and electric utility facilities, environmental and worker assessments of exposures as well as air pollution control research will be presented. The high morbidity and mortality among industry workers and dwellers in nearby industrial areas may be due to elevated levels of urinary and blood chromium and lead levels and resulting from increased air levels of chromium and lea.

Keywords: Heavy metals, Human serum, urine, AAS.

#### I. INTRODUCTION

WHO pointed out that air pollution is sixth-leading cause of death, causing over 2.4million premature deaths worldwide (WHO, 2002). The long-term effect of studies in several large cities predicts 60,000 deaths each year due to cardiac and respiratory problems which caused by air pollutants. It is generally accepted that geographical features of the study area may affect the concentrations of air pollutants and play an important role in the variation of meteorological conditions. Highly elevated concentrations of particulate matter with mean aerodynamic diameter less than 10  $\mu$ m (PM<sub>10</sub>) have been reported during low-level

temperature inversion in the Utah Valley as a result of local emissions being trapped in a stagnant air mass near the valley floor (Pope, 1996). Some of the important heavy metals include lead (Pb), cadmium (Cd),zinc (Zn), mercury (Hg), arsenic (As), chromium(Cr), silver (Ag), iron (Fe),and copper(Cu).

The major findings of the study are: (1) Air pollutants can be classified as suspended particulate matter, gases and vapours that are present in the atmosphere in abnormally high concentrations. The main components of suspended particulate matter are coarse particles such as soil and mineral ash or fine particles found in wood smoke or coming from engine exhausts. Gaseous air pollutants are pricipally oxides of Nitrogen, Ozone, Carbon monoxide, Sulphur dioxide, Ammonia and Volatile organic compounds.

#### **II. MATERIALS AND METHODS**

#### A. Sample Collection

The blood samples were collected from chemical industry workers and patients from nearby industries in heparinized 5 mL, glass test-tubes containing 1 mg of sodium heparin[83]. The hospital staff served as control group.

#### B. Detection of Chromium and Lead levels in Human blood and urine samples

The measurements of chromium and lead levels were performed using atomic absorption

spectrophotometry. The urine samples were collected in 250 mL polyethylene bottles which were thoroughly washed and cleaned at the laboratory and to which 0.1 g of Na<sub>4</sub>EDTA was added before used. New pyrex glass equipment and reagent –grade chemicals contained an acceptable amounts of chromium when we operated at the parts - per – billion level. The glass equipment was boiled twice in 1:1 aqua regia (and kept overnight in double – distilled reagent –grade nitric acid(1:1). Before use the glass equipment was thoroughly washed with distilled water. 50 mL of urine samples were added to 5 mL of double distilled nitric acid and evaporated in 100 mL glass beakers to an approximate volume of 7 mL, when nitrous gases appear. The beaker was then covered with a watch glass and the solution was evaporated to dryness. When the colour of the residue became yellowish white, the residue was dissolved in 2 mL of nitric acid and the glass equipment was rinsed with distilled water to a total volume of 10 mL. When the colour became brownish, 2 mL of acid was added, and the solution was evaporated a second time and then dissolved as described previously.

A series of urine specimens was analyzed with the standard addition method. From one sample to another the interferences seemed to be approximately constant, and a constant calibration factor could well be used. The data of chromium levels in urine samples of workers and dwellers were recorded.



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor:6.887 Volume 5 Issue X, October 2017- Available at www.ijraset.com

### **III. RESULT AND DISCUSSION**

An important health risk factor for the cement industry workers is occupational exposure to chromium, which is used as a basic tanning pigment. The workers on exposure to cement dust, which contains chromium and lead in the protein bound form, exhibited a higher mean concentration of urinary and blood chromium and lead than the reference values. The personal sampling conducted at different work sites exhibited higher levels of total chromium. The high morbidity (15%) observed in the workers and dwellers in comparison to the reference values (5%) could be due to high levels of chromium and lead in the biological samples of the exposed workers and dwellers and air samples collected at the work site. The higher biological values of chromium and lead among the workers and dwellers could be explained by atmospheric pollution caused by the liberated cement dust at the work place and near the residential areas. In the literature that it is the metabolism distribution and transport of the chromium and lead in the blood that is a casual factor for increased respiratory morbidity. The hexavalent chromium is rapidly absorbed by the lungs into the blood and easily penetrates the cellular membranes and binds to the hemoglobin in the red blood cells thereby affecting the oxygen carrying capacity and impairing the lung function status. Our study confirmed the, earlier reports that the biological levels of chromium (in urine and blood) are good indicators of the exposure levels of chromium and lead at the work place.

We observed that blood chromium levels of the workers, nearby dwellers and controls were found to be 0.7248 mg/L, 0.7580mg/L, 0.6391 mg/L, and 0.5251mg/L respectively and the urine chromium levels were found to be  $35 - 45 \ \mu g / L$ , and  $25-35 \ \mu g / L$ , and  $4.5 - 12.5 \ \mu g / L$  respectively. Similarly, we observed, the blood lead levels were found to be  $20 \ \mu g / L$ ,  $62 \ \mu g / L$ , and  $38 \ \mu g / L$  respectively and the urine lead levels were found to be  $32 \ \mu g / L$ ,  $25 \ \mu g / L$ , and  $1 - 5 \ \mu g / L$  respectively.

Workers and dwellers Serum samples	Chromium in serum (µg /L)	Chromium in serum (µg /L) (Control)	Lead in serum (µg /L)	Lead in serum (µg /L) (Control)
01 02 03 04 05 06 07 08 09 10	$\begin{array}{c} 0.5251 \\ 0.5825 \\ 0.6240 \\ 0.6805 \\ 0.7248 \\ 0.4589 \\ 0.7580 \\ 0.6391 \\ 0.7045 \\ 0.6582 \end{array}$	0.0119 0.0138 0.0145 0.0128 0.0153 0.0110 0.0147 0.0142 0.0141 0.0124	$\begin{array}{c} 0.2421\\ 0.2648\\ 0.3468\\ 0.2065\\ 0.3867\\ 0.4219\\ 0.4965\\ 0.5824\\ 0.5710\\ 0.6210\\ \end{array}$	$\begin{array}{c} 0.0124\\ 0.0148\\ 0.0119\\ 0.0105\\ 0.0128\\ 0.0135\\ 0.0139\\ 0.0126\\ 0.0117\\ 0.0145\end{array}$

Table – 1. Concentration	of	Chromium and Lead level	in serum	of Cement industry workers
	· · ·	ein einam and Bead ie iei		







International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor:6.887 Volume 5 Issue X, October 2017- Available at www.ijraset.com

Workers and dwellers Urine samples	Chromium in blood (µg/L)	Chromium in blood (µg/L) (Control)	Lead in urine (µg /L)	Lead in urine (µg /L) (Control)
01	35.0	4.5	25.2	1.0
02	35.5	5.2	26.3	2.2.
03	36.4	7.4	27.1	2.0
04	37.8	8.2	27.5	2.5
05	38.0	9.5	28.7	2.4
06	38.4	10.0	28.0	2.7
07	39.2	10.3	29.2	3.0
08	40.3	10.6	30.1	3.2
09	42.9	9.4	31.3	3.5
10	41.8	9.8	32.4	4.0

Table - 2. Concentration of Chromium and Lead level	in urine of Cement industry workers
---	-------------------------------------



Fig.2.1

## **IV. CONCLUSION**

Health hazards associated with chronic exposures to urban air pollution are now well recognized. Although air pollution affects the population in general, children are particularly vulnerable to toxiceffects of air pollution because of more outdoor exposures, greater intake of air in relation to bodyweight, lower breathing zone where particulate concentration is higher and, most importantly, a child's underdeveloped immunity. Air pollution-related health effects ranged from low birth weight, reduced lung development and lung function to behavioral alterations. Compared with the developed nations, however, very little is known about the health effects of urban air pollution on children's health in developing countries including India. The result showed that Chromium and lead concentration is higher than standard. One of the main objectives of Air Act is to protect. Those who were having higher concentration of chromium and lead levels in blood and urine were having severe form of COPD and asthma and there were no neurological deficit as the leaf levels were below the toxic levels. This study recommends that the biomonitoring of air levels of chromium, PM and other organic pollutants levels at the work place and the nearby surroundings that can be used as a tool for mitigating health hazards and risk factors in the exposed community capacity. The result showed that Chromium and lead concentration is higher than standard.



#### REFERENCES

- [1] Shinggu, D.Y., Ogugbuaja, V., Toma, O. and Barminas, J.T. (2010) Determination of Heavy Metals in Street Dust in Yola, Nigeria. African Journal of Pure and Applied Chemistry, 4, 17-20.
- [2] Lopez, J.M., Callen, M.S., Murillo, R., Garcia, T., Navarro, M.V., de la Cruz, M.T. and Mastral, A.M. (2005) Levels of Selected Metals in Ambient Air PM10 in an Urban Site of Zaragoza, Spain. Environmental Research, 99, 58-67.
- [3] Melaku, S., Morris, V., Raghavan, D. and Hosten, C. (2008) Seasonal Variation of Heavy Metals in Ambient Air and Precipitation, at a Single Site in Washington, DC. Environmental Pollution, 155, 88-98.
- [4] Bhargava, A., Gupta, K., Bhargava, R. and Pardhi, S. (2003) The Effects Automobile Exhaust on Total N, P and HeavyMetals of Road Side Sugarcane at District, Saharanpur. Ad Plant Science, **16**, 557-560.
- [5] Awan, M.A. and Ahmed, S.H. (2011) Determination of Total Suspended Particulate Matter and Heavy Metals in Ambient Air of Four Cities of Pakistan. Ironical Journal of Energy & Environment, 2, 128-132.
- [6] Lazor, P., Tomas, J., Toth, T., Toth, J. and Ceryova, S. (2012) Monitoring of Air Pollution and Atmospheric Deposition fleavy Metals by Analysis of Honey. Journal of Microbiology, Biotechnology and Food Sciences, 1, 522-533.
- [7] Kim, K., Lee, J. and Jang, M. (2002) Metals in Airborne Particulate Matter from the First and Second Industrial Complex Area of Taejon City, Korea. Environmental Pollution, 118, 41-51.
- [8] Shah, J., Nagpa, J.T. and Brnadon, C. (2006) Urban air Quality Management Strategy in Asia-Guide Book. The WorldBank, USA, 17-21.
- [9] Ahmed, F. and Ishiga, H. (2006) Trace Metals Concentration in Street Dusts of Dhaka City Bangladesh. Atmospheric Environment, 40, 3835-3844. [10] Sezgin, N., Ozean, H.K., Demir, G., Nemlioglu, S. and Bayet, C. (2006) Determination of Heavy Metals in Street Dust in Istanbul Highway. Environmental International, 29, 797-985.\
- [10] Aybek, A. and Selcuk, A. (2007) Dust Exposures in Tractor and Combine Operations in Eastern Mediterranean, Turkey. Journal of Environmental Biology, 28, 717-722. <u>http://www.jeb.co.in/</u>
- [11] Pope, C.A., Bates, D.A. and Raizenne, M.E. (1995) Health Effects of Particulate Air Pollution: Time for Reassessment? Environmental Health Perspective, **103**, 472-479.
- [12] Gbadebo, A.M. and Bankole, O.D. (2007) Analysis of Potententially Toxic Metals in Airborne Cement Dust around Sagamu South Western Nigeria. Journal of Applied Sciences, 7, 35-40.
- [13] Rao, M.N. and Rao, H.V.N. (1989) Air Pollution. Data McGraw-Hill Publishing Company Ltd., New Delhi, 10-25.
- [14] Tsor, O.J. (2003) The Characterization of Atmospheric Aerosol during Harmattan in Three Nigerian Cities. M.Sc. Thesis, University of Jos, Jos
- [15] Stone, E.A., Yoon, S.C. and Schauer, J.J. (2011) Chemical Characterization of Fine and Coarse Particles in Gosan, Korea during Springtime Dust Events. Aerosol Air Quality Research, 11, 31-43.
- [16] Zhao, J., Zhang, F., Xu, Y., Chen, J., Yin, L., Shang, X. and Xu, L. (2011) Chemical Characterizations of ParticulateMatter during a Heavy Dust Episode in a Coastal City, Xiamen. Aerosol Air Quality Research, 11, 299-308
- [17] Sombo, T. (2006) Thesis Particulate Matter Sampling and Characterization. M.Sc. Thesis, University of Agriculture, Makurdi
- [18] [19] Peters, A., Dockery, D.W., Millerm J.F. and Mittelman, M.A. (2001) Increase Particulate Matter and the Triggering Myocardial Infarction. Circulation, 103, 2810-2815.
- [19] Curtis, L., Pea, W., Smith-Willis, P., Fenyves, E. and Pan, Y. (2006) Adverse Health Effects of Outdoor Air Pollutants. Environment International, 32, 815-830.
- [20] Environments of Northern India. Department of Civil Engineering, Indian Institute of Technology Roorkee, oorkee.
- [21] National Environmental Protection Council (NEPC) (2002) Ambient Air Quality Standards. Emmett, E. A. (1997). Occupational heal











45.98



IMPACT FACTOR: 7.129







# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24\*7 Support on Whatsapp)