



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 5

Issue: XI

Month of publication: November 2017

DOI:

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

A New Iterative Method for Riccati Differential Equation

Shivaji Tate

Department of Mathematics, Kisan Veer Mahavidyalaya, Wai, Maharashtra, India.

Abstract: Riccati differential equations are usually arises in many fields of applied science. In this paper, we suggest a new iterative method proposed by Daftardar-Gejji and Jafari to solve the Riccati differential equation. We use software Mathematic for computations. We discussed numerical examples to demonstrate the efficiency and the accuracy of the proposed Method.

Keywords: Quadratic Riccati Differential Equation, new iterative method

I. INTRODUCTION

The Riccati differential equation is named after the Italian nobleman Count Jacopo Francesco Riccati (1676-1754). The book of Reid [8] contains the fundamental theories of Riccati equation, with applications to random processes, optimal control, and diffusion problems. Beside important engineering and science applications that today are known as the classical proved, such as stochastic realization theory, optimal control, robust stabilization, and network synthesis, the newer applications include such areas as financial mathematics [2, 6].

In this paper, we discussed the solution of Riccati differential equation [7] using of New Iterative Method (NIM)

$$\frac{dy}{dt} = Q(t)y + R(t)y^2 + P(t), y(0) = G(t) \quad (1)$$

Where $Q(t), R(t), P(t), G(t)$ are known scalar functions.

The solution of Riccati differential equation can be find by classical numerical methods such as the forward Euler method and Runge-Kutta method. El-Tawil et al. [5] solved the nonlinear Riccati in an analytic form by Adomian decomposition method (ADM). Tan and Abbasbandy [9] applied Homotopy Analysis Method (HAM) to solve a quadratic Riccati equation. Abbasbandy [1] solved one example of the quadratic Riccati differential equation (with constant coefficient) by He's variation iteration method considering Adomian's polynomials

The paper is organized as follows: New iterative method described briefly in Section 2. Convergence results of NIM are stated in Section 3. Section 4 deals with illustrative examples and the conclusions are summarized in Section 5.

II. NEW ITERATIVE METHOD

The new iterative method (NIM) is described as bellow:

Consider the nonlinear equation [4]

$$u = f + N(u) \quad (2)$$

Where f is a given function, and N is nonlinear operator from a Banach space $B \rightarrow B$. It is assumed that the NIM solution for the Eq. (3) has the form:

$$u = \sum_{i=0}^{\infty} u_i \quad (3)$$

The convergence of series (3) is proved in [3] and described in Section 4.

The nonlinear operator N in Eq. (3) is decomposed by Daftardar-Gejji and Jafari as bellow:

$$N\left(\sum_{i=0}^{\infty} u_i\right) = N(u_0) + \sum_{i=1}^{\infty} \left\{ N\left(\sum_{j=0}^i u_j\right) - N\left(\sum_{j=0}^{i-1} u_j\right) \right\} \quad (4)$$

From Eq. (2) and (3), Eq. (4) is equivalent to

$$\sum_{i=0}^{\infty} u_i = f + N(u_0) + \sum_{i=1}^{\infty} \left\{ N\left(\sum_{j=0}^i u_j\right) - N\left(\sum_{j=0}^{i-1} u_j\right) \right\} \quad (5)$$

We define the following recurrence relation:

$$\begin{cases} G_0 = u_0 = f \\ G_1 = u_1 = N(u_0) \\ G_m = N(u_0 + u_1 + \dots + u_m) - N(u_0 + u_1 + \dots + u_{m-1}) \end{cases} \quad (6)$$

Then

$$\sum_{i=1}^{m+1} u_i = N\left(\sum_{i=1}^m u_i\right) \quad (7)$$

and

$$u(x) = f + \sum_{i=1}^{\infty} u_i \quad (8)$$

The m -term approximate solution of equation (2) is given by $u = \sum_{i=0}^{m-1} u_i$

III.CONVERGENCE OF NIM

The following convergence results for NIM are described by Daftardar-Gejji and Bhalekar[3].

Theorem 1. If N is C^∞ in a neighborhood of u_0 and $\|N^n(u_0)\| \leq L$, for any n and for some real $L > 0$ and $\|u_i\| \leq M < \frac{1}{e}$, $i = 1, 2, \dots$, then the series $\sum_{n=1}^{\infty} G_n$ is absolutely convergent and Moreover, $\|G_n\| \leq LM^n e^{n-1} (e - 1)$, $n=1, 2, 3, 4, \dots$

Theorem 2. If N is C^∞ and $\|N^n(u_0)\| \leq M \leq e^{-1}$, for all n , then $\sum_{n=0}^{\infty} G_n$ is absolutely convergent.

IV.ILLUSTRATIVE EXAMPLES

1) Example. 1. Consider the following equation [5]

$$\frac{dy}{dx} = -y^2(t) + 1 \quad (9)$$

subject to the initial condition

$$y(0) = 0 \quad (10)$$

Applying NIM, We get

$$y(t) = t - \int_0^t y^2(t) dt = f + N(y) \quad (11)$$

$$y_0 = t \quad (12)$$

$$y_1 = -\frac{t^3}{3} \quad (13)$$

$$y_2 = \frac{2}{15}t^5 - \frac{1}{63}t^7 \quad (14)$$

The solution obtained using NIM approximated to 3 components:

$$y(t) = t - \frac{1}{3}t^3 + \frac{2}{15}t^5 - \frac{1}{63}t^7 \quad (15)$$

The exact solution of Example 1 is $y(t) = \frac{e^{2t} - 1}{e^{2t} + 1}$




Fig. 1. The exact solution versus the NIM solution.

It is clear that the accuracy of the NIM can be increased by adding more terms in the solution. Fig. 1 shows the solution obtained using NIM is nearly identical to the exact solution.

2) *Example 2. Consider the following quadratic Riccati differential equation [5]*

$$\frac{dy}{dt} = 2y(t) - y^2(t) + 1 \quad (16)$$

subject to the initial condition

$$y(0) = 0 \quad (17)$$

Using NIM we get

$$y(t) = t + \int_0^t (2y(t) - y^2(t)) dt = f + N(y) \quad (18)$$

$$y_0 = t \quad (19)$$

$$y_1 = t^2 - \frac{1}{3}t^3 \quad (20)$$

$$y_2 = \frac{2}{3}t^3 - \frac{2}{3}t^4 - \frac{1}{15}t^5 + \frac{1}{9}t^6 - \frac{1}{63}t^7 \quad (21)$$

The solution obtained using NIM approximated to 3 components:

$$y(t) = t + t^2 + \frac{1}{3}t^3 - \frac{2}{3}t^4 - \frac{1}{15}t^5 + \frac{1}{9}t^6 - \frac{1}{63}t^7 \quad (22)$$

The exact solution of Example 2 is $y(t) = 1 + \sqrt{2} \tanh\left(\sqrt{2}t + \frac{1}{2} \log\left(\frac{\sqrt{2}-1}{\sqrt{2}+1}\right)\right)$




Fig. 2. The exact solution versus the NIM solution.

From the Fig. 2 it is clear that the solution obtained by NIM is in high agreement with the exact solution.

V. CONCLUSIONS

We have successfully applied new iterative method (NIM) to obtain approximate solutions of Quadratic Riccati Differential Equation. The result shows that NIM is very simple but powerful technique used to solve such type of equations and easy to implement on computer.

REFERENCES

- [1] S. Abbasbandy, "A new application of He's variational iteration method for quadratic Riccati differential equation by using Adomian's polynomials," J. Comput. Appl. Math., vol. 207, pp. 59–63, 2007.
- [2] B. D. O. Anderson and J. B. Moore, "Optimal Control: Linear Quadratic Methods," IEEE, p. 448, 2007.
- [3] S. Bhalekar and V. Daftardar-Gejji, "Convergence of the new iterative method," Int. J. Differ., 2011.
- [4] V. Daftardar-Gejji and H. Jafari, "An iterative method for solving nonlinear functional equations," J. Math. Anal., 753-763, 2006.
- [5] M. A. El-Tawil, A. A. Bahnasawi, and A. Abdel-Naby, "Solving Riccati differential equation using Adomian's decomposition method," Appl. Math. Comput., vol. 157, no. 2, pp. 503–514, 2004.
- [6] I. Lasiecka and R. Triggiani, "Differential and algebraic Riccati equations with application to boundary/point control problems: continuous theory and approximation theory," Lect. notes Control Inf. Sci., 1991.
- [7] Polyanin A. and V. Zaitsev, "Handbook of exact solutions for ordinary differential equations" ,2003.
- [8] W. Reid, "Riccati differential equations, volume 86 of Mathematics in Science and Engineering, New York and London," 1972.
- [9] Y. Tan and S. Abbasbandy, "Homotopy analysis method for quadratic Riccati differential equation," Comm Non-linear Sci Numer Simm, vol. 13, pp. 539–546, 2008.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)