



# IJRASET

International Journal For Research in  
Applied Science and Engineering Technology



---

# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume:** 2017 **Issue:** conference **Month of publication:** December 2017

**DOI:**

[www.ijraset.com](http://www.ijraset.com)

Call:  08813907089

E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)

# Adjacent Vertex Sum Polynomial of Path Related Graphs

A. M. Anto<sup>1</sup>

<sup>1</sup>Assistant Professor in Mathematics, Malankara Catholic College, Kaliyakavilai, Tamil Nadu, India.

**Abstract:** Let  $G = (V, E)$  be a graph. The adjacent vertex sum polynomial of  $G$  is defined as  $S(G, x) = \sum_{i=0}^{\Delta(G)} n_{\Delta(G)-i} x^{\alpha_{\Delta(G)-i}}$ , where  $n_{\Delta(G)-i}$  is the sum of the number of adjacent vertices of all the vertices of degree  $\Delta(G) - i$  and  $\alpha_{\Delta(G)-i}$  is the sum of the degree of adjacent vertices of all the vertices of degree  $\Delta(G) - i$ . In this paper I seek to find the Adjacent Vertex Sum Polynomial of some path related Graphs.

**Keywords:** Adjacent Vertex Sum Polynomial, Splitting graph, Degree splitting graph, Path.

## I. INTRODUCTION

Here I consider simple undirected graphs. The terms not defined here we can refer Frank Harary [3]. The vertex set is denoted by  $V$  and the edge set by  $E$ . For  $v \in V$ ,  $d(v)$  is the number of edges incident with  $v$ , the maximum degree of the graph  $G$  is defined as  $\Delta(G) = \max\{d(v)/v \in V\}$ . Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be two graphs, the union  $G_1 \cup G_2$  is defined to be  $G = (V, E)$  where  $V = V_1 \cup V_2$  and  $E = E_1 \cup E_2$ , the sum  $G_1 + G_2$  is defined as  $G_1 \cup G_2$  together with all the lines joining points of  $V_1$  to  $V_2$ . The Cartesian product of two graphs  $G_1$  and  $G_2$  denoted by  $G = G_1 \times G_2$  is the graph  $G$  such that  $V(G) = V(G_1) \times V(G_2)$ , that is every vertex of  $G_1 \times G_2$  is an ordered pair  $(u, v)$ , where  $u \in V(G_1)$  and  $v \in V(G_2)$  and two distinct vertices  $(u, v)$  and  $(x, y)$  are adjacent in  $G_1 \times G_2$  if either  $u = x$  and  $vy \in E(G_2)$  or  $v = y$  and  $ux \in E(G_1)$ . The graph  $G$  with  $V = S_1 \cup S_2 \cup \dots \cup S_i \cup T$ , where each  $S_i$  is a set of vertices having at least two vertices and having the same degree and  $T = V \setminus \cup S_i$ . The degree splitting graph of  $G$  denoted by  $DS(G)$  and is obtained from  $G$  by adding the vertices  $w_1, w_2, \dots, w_t$  and joining  $w_i$  to each vertex of  $S_i$ ,  $1 \leq i \leq t$  [5]. For each vertex  $v$  of a graph  $G$ , take a new vertex  $v'$ , join  $v'$  to all the vertices of  $G$  which are adjacent to  $v$ . The graph  $S(G)$  thus obtained is called splitting graph of  $G$  [1]. The Path consisting of length  $n$  is denoted by  $P_n$ . The graph  $G = (V, E)$  is simply denoted by  $G$ . Number of vertices in  $G$  is called order of  $G$ .

## II. MAIN RESULTS


### A. Theorem 2.1

Let  $P_m$  and  $P_n$  be paths with order  $m$  and  $n$  respectively. Then  $S(P_m \cup P_n, x) = [2(m+n) - 8]x^{4(m+n)-20} + 4x^8$ .

1) *Proof:* Let  $P_m$  and  $P_n$  be paths with order  $m$  and  $n$  respectively. In  $P_m$ ,  $m - 2$  vertices have degree 2 and 2 vertices have degree 1. In  $P_n$ ,  $n - 2$  vertices have degree 2 and 2 vertices have degree 1. Therefore in this graph  $P_m \cup P_n$ ,  $m - 2$  vertices have degree 2;  $n - 2$  vertices have degree 2 and 4 vertices have degree 1. Hence, sum of the number of adjacent vertices of all the vertices of degree 2 is  $2(m+n) - 8$ , sum of the degree of adjacent vertices of all the vertices of degree 2 is  $4(m+n) - 20$ , sum of the number of adjacent vertices of all the vertices of degree 1 is 4, sum of the degree of adjacent vertices of all the vertices of degree 1 is 8. This gives,  $S(P_m \cup P_n, x) = [2(m+n) - 8]x^{4(m+n)-20} + 4x^8$ .

### B. Example 2.2

Consider the graph  $P_3 \cup P_4$ , then  $S(P_3 \cup P_4, x) = 10x^8$ .



$$\begin{aligned} \text{Here, } S(P_3 \cup P_4, x) &= [2(3 + 4) - 8]x^{4(3+4)-20} + 4x^8. \\ &= 6x^8 + 4x^8. \\ &= 10x^8 \end{aligned}$$

C. Theorem 2.3

Let  $P_m$  and  $P_n$  be paths with order  $m$  and  $n$  respectively. Then  $S(P_m + P_n, x)$

$$\begin{aligned} &= (m - 2)(n + 2)x^{(m-4)[2(n+2)+(n-2)(m+2)+2(m+1)]+2[(2n+3)+(n-2)(m+2)+2(m+1)]} \\ &+ 2(n + 1)x^{2[(n+2)+(n-2)(m+2)+2(m+1)]} \\ &+ (n - 2)(m + 2)x^{(n-4)[2(m+2)+(m-2)(n+2)+2(n+1)]+2[(2m+3)+(m-2)(n+2)+2(n+1)]} \\ &+ 2(m + 1)x^{2[(m+2)+(m-2)(n+2)+2(n+1)]}, \text{ where } m, n \geq 4. \end{aligned}$$

1) *Proof:* Let  $P_m$  and  $P_n$  be paths with order  $m$  and  $n$  respectively. In  $P_m$ ,  $m - 2$  vertices have degree 2 and 2 vertices have degree 1. In  $P_n$ ,  $n - 2$  vertices have degree 2 and 2 vertices have degree 1. Therefore in this graph  $P_m + P_n$ ,  $m - 2$  vertices have degree  $n + 2$ ;  $n - 2$  vertices have degree  $m + 2$ , 2 vertices have degree  $n + 1$  and 2 vertices have degree  $m + 1$ . Hence, sum of the number of adjacent vertices of all the vertices of degree  $n + 2$  is  $(m - 2)(n + 2)$ , sum of the degree of adjacent vertices of all the vertices of degree  $n + 2$  is  $(m - 4)[2(n + 2) + (n - 2)(m + 2) + 2(m + 1)] + 2[(2n + 3) + (n - 2)(m + 2) + 2(m + 1)]$ , sum of the number of adjacent vertices of all the vertices of degree  $n + 1$  is  $2(n + 1)$ , sum of the degree of adjacent vertices of all the vertices of degree  $n + 1$  is  $2[(n + 2) + (n - 2)(m + 2) + 2(m + 1)]$ , sum of the number of adjacent vertices of all the vertices of degree  $m + 2$  is  $(n - 2)(m + 2)$ , sum of the degree of adjacent vertices of all the vertices of degree  $m + 2$  is  $(n - 4)[2(m + 2) + (m - 2)(n + 2) + 2(n + 1)] + 2[(2m + 3) + (m - 2)(n + 2) + 2(n + 1)]$ , sum of the number of adjacent vertices of all the vertices of degree  $m + 1$  is  $2(m + 1)$ , sum of the degree of adjacent vertices of all the vertices of degree  $m + 1$  is  $2[(m + 2) + (m - 2)(n + 2) + 2(n + 1)]$ .

This gives,  $S(P_m + P_n, x)$

$$\begin{aligned} &= (m - 2)(n + 2)x^{(m-4)[2(n+2)+(n-2)(m+2)+2(m+1)]+2[(2n+3)+(n-2)(m+2)+2(m+1)]} \\ &+ 2(n + 1)x^{2[(n+2)+(n-2)(m+2)+2(m+1)]} \\ &+ (n - 2)(m + 2)x^{(n-4)[2(m+2)+(m-2)(n+2)+2(n+1)]+2[(2m+3)+(m-2)(n+2)+2(n+1)]} \\ &+ 2(m + 1)x^{2[(m+2)+(m-2)(n+2)+2(n+1)]}. \end{aligned}$$

D. Example 2.4

Consider the graph  $P_4 \cup P_4$ , then  $S(P_4 + P_4, x) = 24x^{66} + 20x^{56}$ .




Figure: 2.2

Here,  $S(P_4 + P_4, x)$

$$\begin{aligned}
 &= (m - 2)(n + 2)x^{(m-4)[2(n+2)+(n-2)(m+2)+2(m+1)]+2[(2n+3)+(n-2)(m+2)+2(m+1)]} \\
 &+ 2(n + 1)x^{2[(n+2)+(n-2)(m+2)+2(m+1)]} \\
 &+ (n - 2)(m + 2)x^{(n-4)[2(m+2)+(m-2)(n+2)+2(n+1)]+2[(2m+3)+(m-2)(n+2)+2(n+1)]} \\
 &+ 2(m + 1)x^{2[(m+2)+(m-2)(n+2)+2(n+1)]}. \\
 &= 12x^{66} + 10x^{56} + 12x^{66} + 10x^{56}. \\
 &= 24x^{66} + 20x^{56}.
 \end{aligned}$$

*E. Theorem 2.5*

Let  $P_m$  and  $P_n$  be paths with order  $m$  and  $n$  respectively. Then  $S(s(P_m \cup P_n), x) = 4[(m - 2) + (n - 2)]x^{12[(m-4)+(n-4)]+36} + 2[(m - 2) + (n - 2) + 4]x^{8[(m-4)+(n-4)]+48} + 4x^{16}$ , where  $m, n \geq 4$ .

1) *Proof:* Let  $P_m$  and  $P_n$  be paths with order  $m$  and  $n$  respectively. In  $P_m$ ,  $m - 2$  vertices have degree 2 and 2 vertices have degree 1. In  $P_n$ ,  $n - 2$  vertices have degree 2 and 2 vertices have degree 1. Therefore in this graph  $s(P_m \cup P_n)$ ,  $m - 2$  vertices have degree 4;  $n - 2$  vertices have degree 4,  $m$  vertices have degree 2;  $n$  vertices have degree 2, and 4 vertices have degree 1. Hence, sum of the number of adjacent vertices of all the vertices of degree 4 is  $4[(m - 2) + (n - 2)]$ , sum of the degree of adjacent vertices of all the vertices of degree 4 is  $4(m + n) - 20$ , sum of the number of adjacent vertices of all the vertices of degree 2 is  $2[(m - 2) + (n - 2) + 4]$ , sum of the degree of adjacent vertices of all the vertices of degree 2 is  $8[(m - 4) + (n - 4)] + 48$ , sum of the number of adjacent vertices of all the vertices of degree 1 is 4, sum of the degree of adjacent vertices of all the vertices of degree 1 is 16. This gives,  $S(s(P_m \cup P_n), x) = 4[(m - 2) + (n - 2)]x^{12[(m-4)+(n-4)]+36}$

$$+ 2[(m - 2) + (n - 2) + 4]x^{8[(m-4)+(n-4)]+48} + 4x^{16}.$$

*F. Example 2.6*

Consider the graph  $s(P_3 \cup P_4)$ , then  $S(s(P_4 \cup P_4), x) = 16x^{36} + 16x^{48} + 4x^{16}$ .




Figure: 2.3

Here,  $S(s(P_m \cup P_n), x) = 4[(m - 2) + (n - 2)]x^{12[(m-4)+(n-4)]+36}$   
 $+ 2[(m - 2) + (n - 2) + 4]x^{8[(m-4)+(n-4)]+48} + 4x^{16}$ .  
 $= 16x^{36} + 16x^{48} + 4x^{16}$ .

G. Theorem 2.7

Let  $P_m$  and  $P_n$  be paths with order  $m$  and  $n$  respectively.

Then  $S(DS(P_m \cup P_n), x) = 3[(m - 2) + (n - 2)]x^{(n-4)(n+4)+(m-4)(m+4)+2(m+n-4)+20} + (m - 2)x^{3(m-2)} + (n - 2)x^{3(n-2)} + 12x^{28}$ , where  $m, n \geq 4$ .

1) Proof: Let  $P_m$  and  $P_n$  be paths with order  $m$  and  $n$  respectively. In  $P_m$ ,  $m - 2$  vertices have degree 2 and 2 vertices have degree 1. In  $P_n$ ,  $n - 2$  vertices have degree 2 and 2 vertices have degree 1. Therefore in this graph  $DS(P_m \cup P_n)$ ,  $m - 2$  and  $n - 2$  vertices have degree 3; 1 vertex has degree  $m - 2$ , 1 vertex has degree  $n - 2$ , 6 vertices have degree 2. Hence, sum of the number of adjacent vertices of all the vertices of degree 3 is  $3[(m - 2) + (n - 2)]$ , sum of the degree of adjacent vertices of all the vertices of degree 3 is  $(n - 4)(n + 4) + (m - 4)(m + 4) + 2(m + n - 4) + 20$ , sum of the number of adjacent vertices of all the vertices of degree  $m - 2$  is  $m - 2$ , sum of the degree of adjacent vertices of all the vertices of degree  $m - 2$  is  $3(m - 2)$ , sum of the number of adjacent vertices of all the vertices of degree  $n - 2$  is  $n - 2$ , sum of the degree of adjacent vertices of all the vertices of degree  $n - 2$  is  $3(n - 2)$ , sum of the number of adjacent vertices of all the vertices of degree 2 is 12, sum of the degree of adjacent vertices of all the vertices of degree 2 is 28. This gives,  $S(DS(P_m \cup P_n), x) = 3[(m - 2) + (n - 2)]x^{(n-4)(n+4)+(m-4)(m+4)+2(m+n-4)+20} + (m - 2)x^{3(m-2)} + (n - 2)x^{3(n-2)} + 12x^{28}$ , where  $m, n \geq 4$ .

H. Example 2.8

Consider the graph  $DS(P_4 \cup P_4)$ , then  $S(DS(P_4 \cup P_4), x) = 12x^{28} + 2x^6 + 2x^6 + 12x^{28}$ .




Figure: 2.4

Here,  $S(DS(P_m \cup P_n), x) = 3[(4 - 2) + (4 - 2)]x^{(4-4)(4+4)+(4-4)(4+4)+2(4+4-4)+20}$   
 $+ (4 - 2)x^{3(4-2)} + (4 - 2)x^{3(4-2)} + 12x^{28}$ .  
 $= 12x^{28} + 2x^6 + 2x^6 + 12x^{28}$ .

**REFERENCES**

- [1] E.Samathkumar and H.B.Waliker, On splitting graph of a graph, J. Karnatak Univ. Sci., (25-26) (1980-81), 13-16
- [2] E. Sukumaran, "Adjacent vertex sum polynomial" IJMA-6(6), 2015, 7-1
- [3] Frank Harary, 1872,"Graph Theory", Addition – Wesley Publishing Company
- [4] Gary Chartrand and Ping Znank, "Introduction to Graph Theory", Tata McGraw-Hill Edition
- [5] S. S. Sandhya, C. Jeyasekaran, C. D. Raj (2013), "Harmonic Mean Labelling Of Degree Splitting Graphs" Bulletin of Pure and Applied Sciences, 32E, 99-112.



10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)