iJRASET
International Journal For Research in Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 2017 **Issue:** Conference **Month of publication:** December 2017

DOI:

www.ijraset.com

Call: 08813907089 | E-mail ID: ijraset@gmail.com

Adjacent Vertex Sum Polynomial of Path Related Graphs

A. M. Anto[1]
[1]*Assistant Professor in Mathematics, MalanKara Catholic College, Kaliyakavilai, Tamil Nadu, India.*

Abstract: ***Let $G = (V, E)$ be a graph. The adjacent vertex sum polynomial of G is defined as $S(G, x) = \sum_{i=o}^{\Delta(G)} n_{\Delta(G)-i} x^{\alpha_{\Delta(G)-i}}$, where $n_{\Delta(G)-i}$ is the sum of the number of adjacent vertices of all the vertices of degree $\Delta(G) - i$ and $\alpha_{\Delta(G)-i}$ is the sum of the degree of adjacent vertices of all the vertices of degree $\Delta(G) - i$. In this paper I seek to find the Adjacent Vertex Sum Polynomial of some path related Graphs.***

Keywords: Adjacent Vertex Sum Polynomial, Splitting graph, Degree splitting graph, Path.

I. INTRODUCTION

Here I consider simple undirected graphs. The terms not defined here we can refer Frank Harary [3]. The vertex set is denoted by V and the edge set by E. For $v \epsilon V$, $d(v)$ is the number of edges incident with v, the maximum degree of the graph G is defined as $\Delta(G) = \max\{d(v)/v \epsilon V\}$. Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be two graphs, the union $G_1 \cup G_2$ is defined to be $G = (V, E)$ where $V = V_1 \cup V_2$ and $E = E_1 \cup E_2$, the sum $G_1 + G_2$ is defined as $G_1 \cup G_2$ together with all the lines joining points of V_1 to V_2. The Cartesian product of two graphs G_1 and G_2 denoted by $G = G_1 \times G_2$ is the graph G such that $V(G) = V(G_1) \times V(G_2)$, that is every vertex of $G_1 \times G_2$ is an ordered pair (u, v), where $u \in V(G_1)$ and $v \in V(G_2)$ and two distinct vertices (u, v) and (x, y) are adjacent in $G_1 \times G_2$ if either $u = x$ and $vy \in E(G_2)$ or $v = y$ and $ux \in E(G_1)$. The graph G with $V = S_1 \cup S_2 \cup \dots \cup S_i \cup T$, where each S_i is a set of vertices having at least two vertices and having the same degree and $T = V \backslash \cup S_i$. The degree splitting graph of G denoted by $DS(G)$ and is obtained from G by adding the vertices $w_1, w_2, \dots, w_t$ and joining w_i to each vertex of $S_i, 1 \leq i \leq t$ [5]. For each vertex v of a graph G, take a new vertex v′, join v′ to all the vertices of G which are adjacent to v. The graph $s(G)$ thus obtained is called splitting graph of G [1]. The Path consisting of length n is denoted by P_n. The graph $G = (V, E)$ is simply denoted by G. Number of vertices in G is called order of G.

II. MAIN RESULTS

A. Theorem 2.1

Let P_m and P_n be paths with order m and n respectively. Then $S(P_m \cup P_n, x) = [2(m + n) - 8]x^{4(m+n)-20} + 4x^8$.

1) Proof: Let P_m and P_n be paths with order m and n respectively. In P_m, $m - 2$ vertices have degree 2 and 2 vertices have degree 1. In P_n, $n - 2$ vertices have degree 2 and 2 vertices have degree 1. Therefore in this graph $P_m \cup P_n$, $m - 2$ vertices have degree 2; $n - 2$ vertices have degree 2 and 4 vertices have degree 1. Hence, sum of the number of adjacent vertices of all the vertices of degree 2 is $2(m + n) - 8$, sum of the degree of adjacent vertices of all the vertices of degree 2 is $4(m + n) - 20$, sum of the number of adjacent vertices of all the vertices of degree 1 is 4, sum of the degree of adjacent vertices of all the vertices of degree 1 is8. This gives, $S(P_m \cup P_n, x) = [2(m + n) - 8]x^{4(m+n)-20} + 4x^8$.

B. Example 2.2

Consider the graph $P_3 \cup P_4$, then $S(P_3 \cup P_4, x) = 10x^8$.




Figure: 2.1

©IJRASET (UGC Approved Journal): All Rights are Reserved

Here, $S(P_3 \cup P_4, x) = [2(3+4)-8]x^{4(3+4)-20} + 4x^8$.

$= 6x^8 + 4x^8$.

$= 10x^8$

C. Theorem 2.3

Let P_m and P_n be paths with order m and n respectively. Then $S(P_m + P_n, x)$

$= (m-2)(n+2)x^{(m-4)[2(n+2)+(n-2)(m+2)+2(m+1)]+2[(2n+3)+(n-2)(m+2)+2(m+1)]}$

$+2(n+1)x^{2[(n+2)+(n-2)(m+2)+2(m+1)]}$

$+(n-2)(m+2)x^{(n-4)[2(m+2)+(m-2)(n+2)+2(n+1)]+2[(2m+3)+(m-2)(n+2)+2(n+1)]}$

$+2(m+1)x^{2[(m+2)+(m-2)(n+2)+2(n+1)]}$, where $m, n \geq 4$.

1) Proof:Let P_m and P_n be paths with order m and n respectively. In P_m, $m-2$ vertices have degree 2 and 2 vertices have degree 1. In P_n, $n-2$ vertices have degree 2 and 2 vertices have degree 1. Therefore in this graph $P_m + P_n$, $m-2$ vertices have degree $n+2$; $n-2$ vertices have degree $m+2$, 2 vertices have degree $n+1$ and 2 vertices have degree $m+1$. Hence, sum of the number of adjacent vertices of all the vertices of degree $n+2$ is $(m-2)(n+2)$, sum of the degree of adjacent vertices of all the vertices of degree $n+2$ is $(m-4)[2(n+2)+(n-2)(m+2)+2(m+1)]+2[(2n+3)+(n-2)(m+2)+2(m+1)]$, sum of the number of adjacent vertices of all the vertices of degree $n+1$ is $2(n+1)$, sum of the degree of adjacent vertices of all the vertices of degree $n+1$ is $2[(n+2)+(n-2)(m+2)+2(m+1)]$, sum of the number of adjacent vertices of all the vertices of degree $m+2$ is $(n-2)(m+2)$, sum of the degree of adjacent vertices of all the vertices of degree $m+2$ is $(n-4)[2(m+2)+(m-2)(n+2)+2(n+1)]+2[(2m+3)+(m-2)(n+2)+2(n+1)]$, sum of the number of adjacent vertices of all the vertices of degree $m+1$ is $2(m+1)$, sum of the degree of adjacent vertices of all the vertices of degree $m+1$ is $2[(m+2)+(m-2)(n+2)+2(n+1)]$.

This gives, $S(P_m + P_n, x)$

$= (m-2)(n+2)x^{(m-4)[2(n+2)+(n-2)(m+2)+2(m+1)]+2[(2n+3)+(n-2)(m+2)+2(m+1)]}$

$+2(n+1)x^{2[(n+2)+(n-2)(m+2)+2(m+1)]}$

$+(n-2)(m+2)x^{(n-4)[2(m+2)+(m-2)(n+2)+2(n+1)]+2[(2m+3)+(m-2)(n+2)+2(n+1)]}$

$+2(m+1)x^{2[(m+2)+(m-2)(n+2)+2(n+1)]}$.

D. Example 2.4

Consider the graph $P_4 \cup P_4$, then $S(P_4 + P_4, x) == 24x^{66} + 20x^{56}$.

$P_4 + P_4$:




Figure: 2.2

©IJRASET (UGC Approved Journal): All Rights are Reserved

Here, $S(P_4 + P_4, x)$

$$= (m-2)(n+2)x^{(m-4)[2(n+2)+(n-2)(m+2)+2(m+1)]+2[(2n+3)+(n-2)(m+2)+2(m+1)]}$$

$$+2(n+1)x^{2[(n+2)+(n-2)(m+2)+2(m+1)]}$$

$$+(n-2)(m+2)x^{(n-4)[2(m+2)+(m-2)(n+2)+2(n+1)]+2[(2m+3)+(m-2)(n+2)+2(n+1)]}$$

$$+2(m+1)x^{2[(m+2)+(m-2)(n+2)+2(n+1)]}.$$

$$= 12x^{66} + 10x^{56} + 12x^{66} + 10x^{56}.$$

$$= 24x^{66} + 20x^{56}.$$

E. Theorem 2.5

Let P_m and P_n be paths with order m and n respectively. Then $S(s(P_m \cup P_n), x) = 4[(m-2) + (n-2)]x^{12[(m-4)+(n-4)]+36} + 2[(m-2) + (n-2) + 4]x^{8[(m-4)+(n-4)]+48} + 4x^{16}$, where $m, n \geq 4$.

1) Proof: Let P_m and P_n be paths with order m and n respectively. In P_m, $m-2$ vertices have degree 2 and 2 vertices have degree 1. In P_n, $n-2$ vertices have degree 2 and 2 vertices have degree 1. Therefore in this graph $s(P_m \cup P_n)$, $m-2$ vertices have degree 4; $n-2$ vertices have degree 4, m vertices have degree 2; n vertices have degree 2, and 4 vertices have degree 1. Hence, sum of the number of adjacent vertices of all the vertices of degree 4 is $4[(m-2) + (n-2)]$, sum of the degree of adjacent vertices of all the vertices of degree 4 is $4(m+n) - 20$, sum of the number of adjacent vertices of all the vertices of degree 2 is $2[(m-2) + (n-2) + 4]$, sum of the degree of adjacent vertices of all the vertices of degree 2 is $8[(m-4) + (n-4)] + 48$, sum of the number of adjacent vertices of all the vertices of degree 1 is 4, sum of the degree of adjacent vertices of all the vertices of degree 1 is16. This gives, $S(s(P_m \cup P_n), x) = 4[(m-2) + (n-2)]x^{12[(m-4)+(n-4)]+36}$

$$+2[(m-2) + (n-2) + 4]x^{8[(m-4)+(n-4)]+48} + 4x^{16}.$$

F. Example 2.6

Consider the graph $s(P_3 \cup P_4)$, then $S(s(P_4 \cup P_4), x) == 16x^{36} + 16x^{48} + 4x^{16}$.




Figure: 2.3

©IJRASET (UGC Approved Journal): All Rights are Reserved

Here, $S(s(P_m \cup P_n), x) = 4[(m-2)+(n-2)]x^{12[(m-4)+(n-4)]+36}$

$+2[(m-2)+(n-2)+4]x^{8[(m-4)+(n-4)]+48} + 4x^{16}$.

$= 16x^{36} + 16x^{48} + 4x^{16}$.

G. Theorem 2.7

Let P_m and P_n be paths with order m and n respectively.

Then $S(DS(P_m \cup P_n), x) = 3[(m-2)+(n-2)]x^{(n-4)(n+4)+(m-4)(m+4)+2(m+n-4)+20} + (m-2)x^{3(m-2)} + (n-2)x^{3(n-2)} + 12x^{28}$, where $m, n \geq 4$.

1) Proof: Let P_m and P_n be paths with order m and n respectively. In P_m, $m-2$ vertices have degree 2 and 2 vertices have degree 1. In P_n, $n-2$ vertices have degree 2 and 2 vertices have degree 1. Therefore in this graph $DS(P_m \cup P_n)$, $m-2$ and $n-2$ vertices have degree 3; 1 vertex has degree $m-2$, 1 vertex has degree $n-2$, 6 vertices have degree 2. Hence, sum of the number of adjacent vertices of all the vertices of degree 3 is $3[(m-2)+(n-2)]$, sum of the degree of adjacent vertices of all the vertices of degree 3 is $(n-4)(n+4)+(m-4)(m+4)+2(m+n-4)+20$, sum of the number of adjacent vertices of all the vertices of degree $m-2$ is $m-2$, sum of the degree of adjacent vertices of all the vertices of degree $m-2$ is $3(m-2)$, sum of the number of adjacent vertices of all the vertices of degree $n-2$ is $n-2$, sum of the degree of adjacent vertices of all the vertices of degree $n-2$ is $3(n-2)$,sum of the number of adjacent vertices of all the vertices of degree 2 is 12, sum of the degree of adjacent vertices of all the vertices of degree 2 is 28. This gives, $S(DS(P_m \cup P_n), x) = 3[(m-2)+(n-2)]x^{(n-4)(n+4)+(m-4)(m+4)+2(m+n-4)+20}$

$+(m-2)x^{3(m-2)} + (n-2)x^{3(n-2)} + 12x^{28}$, where $m, n \geq 4$.

H. Example 2.8

Consider the graph $DS(P_4 \cup P_4)$, then $S(DS(P_4 \cup P_4), x) == 12x^{28} + 2x^6 + 2x^6 + 12x^{28}$.

$DS(P_4 \cup P_4)$:




Figure: 2.4

Here, $S(DS(P_m \cup P_n), x) = 3[(4-2)+(4-2)]x^{(4-4)(4+4)+(4-4)(4+4)+2(4+4-4)+20}$

$+(4-2)x^{3(4-2)} + (4-2)x^{3(4-2)} + 12x^{28}$.


$= 12x^{28} + 2x^6 + 2x^6 + 12x^{28}$.

©IJRASET (UGC Approved Journal): All Rights are Reserved

REFERENCES

[1] E.Sampathkumar and H.B.Walikar, On splitting graph of a graph, J. Karnatak Univ. Sci., (25-26) (1980-81), 13-16
[2] E. Sukumaran, "Adjacent vertex sum polynomial" IJMA-6(6), 2015, 7-1
[3] Frank Harary, 1872,"Graph Theory", Addition – Wesley Publishing Company
[4] Gary Chartrant and Ping Znank, "Introduction to Graph Theory", Tata McGraw-Hill Edition
[5] S. S. Sandhya, C. Jeyasekaran, C. D. Raj (2013), "Harmonic Mean Labelling Of Degree Splitting Graphs" Bulletin of Pure and Applied Sciences, 32E, 99-112.

©IJRASET (UGC Approved Journal): All Rights are Reserved



10.22214/IJRASET

45.98

IMPACT FACTOR: 7.129

IMPACT FACTOR: 7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)