iJRASET
International Journal For Research in Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 2017 **Issue:** Conference **Month of publication:** December 2017

DOI:

www.ijraset.com

Call: 08813907089 | E-mail ID: ijraset@gmail.com

Intuitionistic Fuzzy Hx Right and Left Ideal of a Hx Ring

R. Muthuraj[1] , M.S.Muthuraman[2]
[1] PG & Research Department of Mathematics, H.H.The Rajah's College, Pudukkottai – 622 001, Tamilnadu, India. E-mail:
[2]Department of Mathematics, PSNA College of Engineering and Technology, Dindigul -624 622, Tamilnadu, India. . E-

Abstract : In this paper, we define the notion of intuitionistic fuzzy HX right and eft ideal of a HX ring and some of their related properties are investigated. We define the necessity and possibility operators of an intuitionistic fuzzy subset of an HX ring and discuss some of its properties. We introduce the concept of an image, pre-image of an intuitionistic fuzzy HX ring and discuss in detail a series of homomorphic and anti homomorphic properties of an intuitionistic fuzzy HX ring.
Keywords: intuitionistic fuzzy set, fuzzy HX ring, intuitionistic fuzzy HX right and left ideal, homomorphism and anti homomorphism of an intuitionistic fuzzy HX right and left ideal, image and pre-image of an intuitionistic fuzzy set.

I. INTRODUCTION

In 1965, Zadeh [14] introduced the concept of fuzzy subset μ of a set X as a function from X into the closed unit interval [0, 1] and studied their properties. Fuzzy set theory is a useful tool to describe situations in which the data or imprecise or vague and it is applied to logic, set theory, group theory, ring theory, real analysis, measure theory etc. In 1967, Rosenfeld [13] defined the idea of fuzzy subgroup and gave some of its properties. Li Hong Xing [5] introduced the concept of HX group. In 1982 Wang-jin Liu[7] introduced the concept of fuzzy ring and fuzzy ideal. With the successful upgrade of algebraic structure of group many researchers considered the algebraic structure of some other algebraic systems in which ring was considered as first. In 1988, Professor Li Hong Xing [6] proposed the concept of HX ring and derived some of its properties, then Professor Zhong [2,3] gave the structures of HX ring on a class of ring. R.Muthuraj et.al[12]., introduced the concept of fuzzy HX ring. In this paper we define a new algebraic structure of an intuitionistic fuzzy HX right and left ideal of a HX ring and investigate some related properties. We define the necessity and possibility operators of an intuitionistic fuzzy subset of an intuitionistic fuzzy HX ring and discuss some of its properties. Also we introduce the image and pre-image of an intuitionistic fuzzy set in an intuitionistic fuzzy HX right and left ideal of a HX ring and discuss some of its properties.

II. PRELIMINARIES

In this section, we site the fundamental definitions that will be used in the sequel. Throughout this paper, R = (R ,+, ·) is a Ring, e is the additive identity element of R and xy, we mean x.y

A. Definition

Let R be a ring. In $2^R - \{\phi\}$, a non-empty set $\vartheta \subset 2^R - \{\phi\}$ with two binary operation ' + ' and '.' is said to be a HX ring on R if ϑ is a ring with respect to the algebraic operation defined by

1) $A + B = \{a + b / a \in A \text{ and } b \in B\}$, which its null element is denoted by Q , and the negative element of A is denoted by – A.

2) $AB = \{ab / a \in A \text{ and } b \in B\}$,

3) $A (B + C) = AB + AC$ and $(B + C) A = BA + CA$.

B. Definition

Let R be a ring. Let μ be a fuzzy ring defined on R. Let $\vartheta \subset 2^R - \{\phi\}$ be a HX ring. A fuzzy subset λ^μ of ϑ is called a fuzzy HX ring on ϑ or a fuzzy ring induced by μ if the following conditions are satisfied. For all $A,B \in \vartheta$,

1) $\lambda^\mu (A - B) \geq \min \{ \lambda^\mu (A), \lambda^\mu (B) \}$,

2) $\lambda^\mu (AB) \geq \min \{ \lambda^\mu (A), \lambda^\mu (B) \}$

where $\lambda^\mu (A) = \max\{ \mu(x) / \text{ for all } x \in A \subseteq R \}$.

III. PROPERTIES OF AN INTUITIONISTIC FUZZY HX RIGHT AND LEFT IDEAL

A. Definition

©IJRASET (UGC Approved Journal): All Rights are Reserved

Let R be a ring. Let H = { ⟨x, μ(x), η(x)⟩ / x∈R } be an intuitionistic fuzzy set defined on a ring R, where μ : R→[0,1], η : R→[0,1] such that $0 \le \mu(x) + \eta(x) \le 1$. Let $\Re \subset 2^R - \{\phi\}$ be a HX ring. An intuitionistic fuzzy subset $\lambda^H = \{ \langle A, \lambda^\mu(A), \lambda^\eta(A)\rangle / A \in \Re$ and $0 \le \lambda^\mu(A) + \lambda^\eta(A) \le 1\}$ of $\Re$ is called an intuitionistic fuzzy HX right ideal on $\Re$ or an intuitionistic fuzzy right ideal induced by H if the following conditions are satisfied. For all A,B ∈$\Re$,

$\lambda^\mu(A - B) \ge \min\{\lambda^\mu(A), \lambda^\mu(B)\}$,

$\lambda^\mu(AB) \ge \lambda^\mu(A)$

$\lambda^\eta(A - B) \le \max\{\lambda^\eta(A), \lambda^\eta(B)\}$

$\lambda^\eta(AB) \le \lambda^\eta(A)$.

where $\lambda^\mu(A) = \max\{\mu(x) /$ for all $x \in A \subseteq R\}$ and $\lambda^\eta(A) = \min\{\eta(x) /$ for all $x \in A \subseteq R\}$.

B. Definition

Let R be a ring. Let H = { ⟨x, μ(x), η(x)⟩ / x∈R } be an intuitionistic fuzzy set defined on a ring R, where μ : R→[0,1], η : R→[0,1] such that $0 \le \mu(x) + \eta(x) \le 1$. Let $\Re \subset 2^R - \{\phi\}$ be a HX ring. An intuitionistic fuzzy subset $\lambda^H = \{ \langle A, \lambda^\mu(A), \lambda^\eta(A)\rangle / A \in \Re$ and $0 \le \lambda^\mu(A) + \lambda^\eta(A) \le 1\}$ of $\Re$ is called an intuitionistic fuzzy HX left ideal on $\Re$ or an intuitionistic fuzzy left ideal induced by H if the following conditions are satisfied. For all A,B ∈$\Re$,

1) $\lambda^\mu(A - B) \ge \min\{\lambda^\mu(A), \lambda^\mu(B)\}$,

2) $\lambda^\mu(AB) \ge \lambda^\mu(B)$

3) $\lambda^\eta(A - B) \le \max\{\lambda^\eta(A), \lambda^\eta(B)\}$

4) $\lambda^\eta(AB) \le \lambda^\eta(B)$.

where $\lambda^\mu(A) = \max\{\mu(x) /$ for all $x \in A \subseteq R\}$ and $\lambda^\eta(A) = \min\{\eta(x) /$ for all $x \in A \subseteq R\}$

C. Theorem

If H is an intuitionistic fuzzy right ideal of a ring R then the intuitionistic fuzzy subset λ^H is an intuitionistic fuzzy HX right ideal of a HX ring $\Re$.

1) Proof: Let H be an intuitionistic fuzzy right ideal of R.

$\min\{\lambda^\mu(A), \lambda^\mu(B)\} = \min\{\max\{\mu(x)/$for all $x \in A \subseteq R\}$,

$\max\{\mu(y)/$for all $y \in B \subseteq R\}\}$

$= \min\{\mu(x_0), \mu(y_0)\}$

$\le \mu(x_0 - y_0)$

$\le \max\{\mu(x-y)/$for all $x-y \in A-B \subseteq R\}$

$\le \lambda^\mu(A - B)$

$\lambda^\mu(A-B) \ge \min\{\lambda^\mu(A), \lambda^\mu(B)\}$.

$\lambda^\mu(AB) = \max\{\mu(xy) /$ for all $x \in A \subseteq R$ and $y \in B \subseteq R\}$

$= \mu(x_0 y_0)$, for $x_0 \in A$ and $y_0 \in B$.

$\ge \mu(x_0)$

$= \max\{\mu(x) /$ for all $x \in A \subseteq R\}$

$\ge \lambda^\mu(A)$.

$\lambda^\mu(AB) \ge \lambda^\mu(A)$.

$\max\{\lambda^\eta(A), \lambda^\eta(B)\} = \max\{\min\{\eta(x)/$for all $x \in A \subseteq R\}$,

$\min\{\eta(y)/$for all $y \in B \subseteq R\}\}$

$= \max\{\eta(x_0), \eta(y_0)\}$

$\ge \eta(x_0 - y_0)$

$\ge \min\{\eta(x-y)/$for all $x-y \in A-B \subseteq R\}$

$\ge \lambda^\eta(A - B)$

$\lambda^\eta(A-B) \le \max\{\lambda^\eta(A), \lambda^\eta(B)\}$.

$\lambda^\eta(AB) = \min\{\eta(xy) /$ for all $x \in A \subseteq R$ and $y \in B \subseteq R\}$

©IJRASET (UGC Approved Journal): All Rights are Reserved

$= \quad \eta(x_0 y_0)$, for $x_0 \in A$ and $y_0 \in B$.

$\leq \quad \eta(x_0)$

$= \quad \min \{ \eta(x) / \text{for all } x \in A \subseteq R\}$

$\leq \quad \lambda^{\eta}(A)$.

$\lambda^{\eta}(AB) \quad \leq \quad \lambda^{\eta}(A)$.

Hence, λ^H is an intuitionistic fuzzy HX right ideal of a HX ring $\Re$.

D. Theorem

If H is an intuitionistic fuzzy left ideal of a ring R then the intuitionistic fuzzy subset λ^H is an intuitionistic fuzzy HX left ideal of a HX ring $\Re$.

1) Proof: It is clear.

E. Theorem

Let G and H be any two intuitionistic fuzzy sets on R. Let γ^G and λ^H be any two intuitionistic fuzzy HX right ideals of a HX ring $\Re$ then their intersection, $\gamma^G \cap \lambda^H$ is also an intuitionistic fuzzy HX right ideal of a HX ring $\Re$.

1) Proof: Let $G = \{ \langle x, \alpha(x), \beta(x)\rangle / x \in R \}$ and $H = \{ \langle x, \mu(x), \eta(x)\rangle / x \in R \}$ be any two intuitionistic fuzzy sets defined on a ring R.

Then, $\gamma^G = \{ \langle A, \gamma^{\alpha}(A), \gamma^{\beta}(A)\rangle / A \in \Re \}$ and $\lambda^H = \{ \langle A, \lambda^{\mu}(A), \lambda^{\eta}(A)\rangle / A \in \Re \}$ be any two intuitionistic fuzzy HX right ideals of a HX ring $\Re$.

$\gamma^G \cap \lambda^H = \{ \langle A, (\gamma^{\alpha} \cap \lambda^{\mu})(A), (\gamma^{\beta} \cup \lambda^{\eta})(A)\rangle / A \in \Re \}$

Let $A,B \in \Re$

a) $(\gamma^{\alpha} \cap \lambda^{\mu})(A-B) = \min \{ \gamma^{\alpha}(A-B) , \lambda^{\mu}(A-B)\}$

$\geq \quad \min \{ \min \{\gamma^{\alpha}(A), \gamma^{\alpha}(B)\}, \min\{ \lambda^{\mu}(A), \lambda^{\mu}(B)\}\}$

$= \quad \min \{ \min \{\gamma^{\alpha}(A), \lambda^{\mu}(A)\}, \min\{\gamma^{\alpha}(B), \lambda^{\mu}(B)\}\}$

$= \quad \min \{ (\gamma^{\alpha} \cap \lambda^{\mu})(A) , (\gamma^{\alpha} \cap \lambda^{\mu})(B)\}$

$(\gamma^{\alpha} \cap \lambda^{\mu})(A-B) \geq \min \{ (\gamma^{\alpha} \cap \lambda^{\mu})(A) , (\gamma^{\alpha} \cap \lambda^{\mu})(B)\}$.

b) $(\gamma^{\alpha} \cap \lambda^{\mu})(AB) = \min \{ \gamma^{\alpha}(AB) , \lambda^{\mu}(AB)\}$

$\geq \quad \min\{ \gamma^{\alpha}(A), \lambda^{\mu}(A)\}$

$= \quad (\gamma^{\alpha} \cap \lambda^{\mu})(A)$.

$(\gamma^{\alpha} \cap \lambda^{\mu})(AB) \geq (\gamma^{\alpha} \cap \lambda^{\mu})(A)$.

c) $(\gamma^{\beta} \cup \lambda^{\eta})(A-B) = \max \{ \gamma^{\beta}(A-B) , \lambda^{\eta}(A-B)\}$

$\leq \quad \max \{\max\{\gamma^{\beta}(A), \gamma^{\beta}(B)\}, \max\{ \lambda^{\eta}(A), \lambda^{\eta}(B)\}\}$

$= \quad \max \{\max \{\gamma^{\beta}(A), \lambda^{\eta}(A)\}, \max\{\gamma^{\beta}(B), \lambda^{\eta}(B)\}\}$

$= \quad \max \{ (\gamma^{\beta} \cup \lambda^{\eta})(A) , (\gamma^{\beta} \cup \lambda^{\eta})(B)\}$

$(\gamma^{\beta} \cup \lambda^{\mu})(A-B) \leq \max \{ (\gamma^{\beta} \cup \gamma^{\eta})(A) , (\gamma^{\beta} \cup \lambda^{\eta})(B)\}$.

d) $(\gamma^{\beta} \cup \lambda^{\eta})(AB) = \max \{ \gamma^{\beta}(AB) , \lambda^{\eta}(AB)\}$

$\leq \quad \max\{ \gamma^{\beta}(A), \lambda^{\eta}(A)\}$

$= \quad (\gamma^{\beta} \cup \lambda^{\eta})(A)$.

$(\gamma^{\beta} \cup \lambda^{\eta})(AB) \leq (\gamma^{\beta} \cup \lambda^{\eta})(A)$.

Hence, $\gamma^G \cap \lambda^H$ is an intuitionistic fuzzy HX right ideal of a HX ring $\Re$.

F. Theorem

Let G and H be any two intuitionistic fuzzy sets on R. Let γ^G and λ^H be any two intuitionistic fuzzy HX left ideals of a HX ring $\Re$ then their intersection, $\gamma^G \cap \lambda^H$ is also an intuitionistic fuzzy HX left ideal of a HX ring $\Re$.

1) Proof: Let $G = \{ \langle x, \alpha(x), \beta(x)\rangle / x \in R \}$ and $H = \{ \langle x, \mu(x), \eta(x)\rangle / x \in R \}$ be any two intuitionistic fuzzy sets defined on a ring R.

Then, $\gamma^G = \{ \langle A, \gamma^{\alpha}(A), \gamma^{\beta}(A)\rangle / A \in \Re \}$ and $\lambda^H = \{ \langle A, \lambda^{\mu}(A), \lambda^{\eta}(A)\rangle / A \in \Re \}$ be any two intuitionistic fuzzy HX left ideals of a HX ring $\Re$.

$\gamma^G \cap \lambda^H = \{ \langle A, (\gamma^{\alpha} \cap \lambda^{\mu})(A), (\gamma^{\beta} \cup \lambda^{\eta})(A)\rangle / A \in \Re \}$

Let $A,B \in \Re$

©IJRASET (UGC Approved Journal): All Rights are Reserved

a) $(\gamma^{\alpha} \cap \lambda^{\mu})(A-B) = \min\{\gamma^{\alpha}(A-B), \lambda^{\mu}(A-B)\}$

$\geq \min\{\min\{\gamma^{\alpha}(A), \gamma^{\alpha}(B)\}, \min\{\lambda^{\mu}(A), \lambda^{\mu}(B)\}\}$

$= \min\{\min\{\gamma^{\alpha}(A), \lambda^{\mu}(A)\}, \min\{\gamma^{\alpha}(B), \lambda^{\mu}(B)\}\}$

$= \min\{(\gamma^{\alpha} \cap \lambda^{\mu})(A), (\gamma^{\alpha} \cap \lambda^{\mu})(B)\}$

$(\gamma^{\alpha} \cap \lambda^{\mu})(A-B) \geq \min\{(\gamma^{\alpha} \cap \lambda^{\mu})(A), (\gamma^{\alpha} \cap \lambda^{\mu})(B)\}$.

b) $(\gamma^{\alpha} \cap \lambda^{\mu})(AB) = \min\{\gamma^{\alpha}(AB), \lambda^{\mu}(AB)\}$

$\geq \min\{\gamma^{\alpha}(B), \lambda^{\mu}(B)\}$

$= (\gamma^{\alpha} \cap \lambda^{\mu})(B)$.

$(\gamma^{\alpha} \cap \lambda^{\mu})(AB) \geq (\gamma^{\alpha} \cap \lambda^{\mu})(B)$.

c) $(\gamma^{\beta} \cup \lambda^{\eta})(A-B) = \max\{\gamma^{\beta}(A-B), \lambda^{\eta}(A-B)\}$

$\leq \max\{\max\{\gamma^{\beta}(A), \gamma^{\beta}(B)\}, \max\{\lambda^{\eta}(A), \lambda^{\eta}(B)\}\}$

$= \max\{\max\{\gamma^{\beta}(A), \lambda^{\eta}(A)\}, \max\{\gamma^{\beta}(B), \lambda^{\eta}(B)\}\}$

$= \max\{(\gamma^{\beta} \cup \lambda^{\eta})(A), (\gamma^{\beta} \cup \lambda^{\eta})(B)\}$

$(\gamma^{\beta} \cup \lambda^{\mu})(A-B) \leq \max\{(\gamma^{\beta} \cup \gamma^{\eta})(A), (\gamma^{\beta} \cup \lambda^{\eta})(B)\}$.

d) $(\gamma^{\beta} \cup \lambda^{\eta})(AB) = \max\{\gamma^{\beta}(AB), \lambda^{\eta}(AB)\}$

$\leq \max\{\gamma^{\beta}(B), \lambda^{\eta}(B)\}$

$= (\gamma^{\beta} \cup \lambda^{\eta})(B)$.

$(\gamma^{\beta} \cup \lambda^{\eta})(AB) \leq (\gamma^{\beta} \cup \lambda^{\eta})(B)$.

Hence, $\gamma^{G} \cap \lambda^{H}$ is an intuitionistic fuzzy HX left ideal of a HX ring $\Re$.

G. Theorem

Let G and H be any two intuitionistic fuzzy sets on R. Let γ^{G} be an intuitionistic fuzzy HX ring and λ^{H} be an intuitionistic fuzzy HX right (left) ideals of a HX ring $\Re$ then their intersection, $\gamma^{G} \cap \lambda^{H}$ is also an intuitionistic fuzzy HX right (left) ideal of a HX ring $\Re$.

1) Proof: It is clear.

H. Theorem

Let G and H be any two intuitionistic fuzzy sets on R. Let γ^{G} be an intuitionistic fuzzy HX ring and λ^{H} be an intuitionistic fuzzy HX ideal of a HX ring $\Re$ then their intersection, $\gamma^{G} \cap \lambda^{H}$ is also an intuitionistic fuzzy HX ideal of a HX ring $\Re$.

I. Theorem

Let G and H be any two intuitionistic fuzzy sets on R. Let γ^{G} and λ^{H} be any two intuitionistic fuzzy HX right ideals of a HX ring $\Re$ then their union, $\gamma^{G} \cup \lambda^{H}$ is also an intuitionistic fuzzy HX right ideal of a HX ring $\Re$.

1) Proof: Let $G = \{\langle x, \alpha(x), \beta(x)\rangle / x \in R\}$ and $H = \{\langle x, \mu(x), \eta(x)\rangle / x \in R\}$ be any two intuitionistic fuzzy sets defined on a ring R.

Then, $\gamma^{G} = \{\langle A, \gamma^{\alpha}(A), \gamma^{\beta}(A)\rangle / A \in \Re\}$ and $\lambda^{H} = \{\langle A, \lambda^{\mu}(A), \lambda^{\eta}(A)\rangle / A \in \Re\}$ be any two intuitionistic fuzzy HX right ideals of a HX ring $\Re$.Then,

$\gamma^{G} \cup \lambda^{H} = \{\langle A, (\gamma^{\alpha} \cup \lambda^{\mu})(A), (\gamma^{\beta} \cap \lambda^{\eta})(A)\rangle / A \in \Re\}$

Let $A,B \in \Re$

a) $(\gamma^{\alpha} \cup \lambda^{\mu})(A-B) = \max\{\gamma^{\alpha}(A-B), \lambda^{\mu}(A-B)\}$

$\geq \max\{\min\{\gamma^{\alpha}(A), \gamma^{\alpha}(B)\}, \min\{\lambda^{\mu}(A), \lambda^{\mu}(B)\}\}$

$= \min\{\max\{\gamma^{\alpha}(A), \lambda^{\mu}(A)\}, \max\{\gamma^{\alpha}(B), \lambda^{\mu}(B)\}\}$

$= \min\{(\gamma^{\alpha} \cup \lambda^{\mu})(A), (\gamma^{\alpha} \cup \lambda^{\mu})(B)\}$

$(\gamma^{\alpha} \cup \lambda^{\mu})(A-B) \geq \min\{(\gamma^{\alpha} \cup \lambda^{\mu})(A), (\gamma^{\alpha} \cup \lambda^{\mu})(B)\}$.

b) $(\gamma^{\alpha} \cup \gamma^{\mu})(AB) = \max\{\gamma^{\alpha}(AB), \lambda^{\mu}(AB)\}$

$\geq \max\{\gamma^{\alpha}(A), \lambda^{\mu}(A)\}$

$= (\gamma^{\alpha} \cup \lambda^{\mu})(A)$.

$(\gamma^{\alpha} \cup \lambda^{\mu})(AB) \geq (\gamma^{\alpha} \cup \lambda^{\mu})(A)$.

©IJRASET (UGC Approved Journal): All Rights are Reserved

c) $(\gamma^{\beta} \cap \lambda^{\eta})(A-B) = \min\{\gamma^{\beta}(A-B), \lambda^{\eta}(A-B)\}$

$\leq \min\{\max\{\gamma^{\beta}(A), \gamma^{\beta}(B)\}, \max\{\lambda^{\eta}(A), \lambda^{\eta}(B)\}\}$

$= \max\{\min\{\gamma^{\beta}(A), \lambda^{\eta}(A)\}, \min\{\gamma^{\beta}(B), \lambda^{\eta}(B)\}\}$

$= \max\{(\gamma^{\beta} \cap \gamma^{\eta})(A), (\gamma^{\beta} \cup \lambda^{\eta})(B)\}$

$(\gamma^{\beta} \cap \lambda^{\mu})(A-B) \leq \max\{(\gamma^{\beta} \cap \gamma^{\eta})(A), (\gamma^{\beta} \cap \lambda^{\eta})(B)\}.$

d) $(\gamma^{\beta} \cap \lambda^{\eta})(AB) = \min\{\gamma^{\beta}(AB), \lambda^{\eta}(AB)\}$

$\leq \min\{\gamma^{\beta}(A), \lambda^{\eta}(A)\}$

$= (\gamma^{\beta} \cap \lambda^{\eta})(A).$

$(\gamma^{\beta} \cap \lambda^{\eta})(AB) \leq (\gamma^{\beta} \cap \lambda^{\eta})(A).$

Hence, $\gamma^{G} \cup \lambda^{H}$ is an intuitionistic fuzzy HX right ideal of a HX ring $\Re$.

J. Theorem

Let G and H be any two intuitionistic fuzzy sets on R. Let γ^{G} and λ^{H} be any two intuitionistic fuzzy HX left ideals of a HX ring $\Re$ then their union, $\gamma^{G} \cup \lambda^{H}$ is also an intuitionistic fuzzy HX left ideal of a HX ring $\Re$.

1) Proof: It is clear.

K Theorem

Let G and H be any two intuitionistic fuzzy sets of R_1 and R_2 respectively. Let $\Re_1 \subset 2^{R_1} - \{\phi\}$ and $\Re_2 \subset 2^{R_2} - \{\phi\}$ be any two HX rings. If γ^{G} and λ^{H} are any two intuitionistic fuzzy HX right ideals of $\Re_1$ and $\Re_2$ respectively then, $\gamma^{G} \times \lambda^{H}$ is also an intuitionistic fuzzy HX right ideal of a HX ring $\Re_1 \times \Re_2$.

1) Proof: Let $G = \{\langle x, \alpha(x), \beta(x)\rangle / x \in R_1\}$ and $H = \{\langle x, \mu(x), \eta(x)\rangle / x \in R_2\}$ be any two intuitionistic fuzzy sets defined on a ring R_1 and R_2 respectively.

Then, $\gamma^{G} = \{\langle A, \gamma^{\alpha}(A), \gamma^{\beta}(A)\rangle / A \in \Re_1\}$ and $\lambda^{H} = \{\langle A, \lambda^{\mu}(A), \lambda^{\eta}(A)\rangle / A \in \Re_2\}$ be any two intuitionistic fuzzy HX right ideal of a HX ring $\Re_1$ and $\Re_2$ respectively. Then,

$(\gamma^{G} \times \lambda^{H}) = \{\langle (A,B), (\gamma^{\alpha} \cap \lambda^{\mu})(A, B), (\gamma^{\beta} \cup \lambda^{\eta})(A, B)\rangle / (A, B) \in \Re_1 \times \Re_2\},$

where $(\gamma^{\alpha} \cap \lambda^{\mu})(A, B) = \min\{\gamma^{\alpha}(A), \lambda^{\mu}(B)\}$, for all $(A, B) \in \Re_1 \times \Re_2$,

$(\gamma^{\beta} \cup \lambda^{\eta})(A, B) = \max\{\gamma^{\beta}(A), \lambda^{\eta}(B)\}$, for all $(A, B) \in \Re_1 \times \Re_2$.

Let $A, B \in \Re_1 \times \Re_2$, where $A = (C,D)$, $B = (E,F)$,

a) $(\gamma^{\alpha} \cap \lambda^{\mu})(A - B) = \min\{\gamma^{\alpha}(C - E), \lambda^{\mu}(D - F)\}$

$\geq \min\{\min\{\gamma^{\alpha}(C), \gamma^{\alpha}(E)\}, \min\{\lambda^{\mu}(D), \lambda^{\mu}(F)\}\}$

$= \min\{\min\{\gamma^{\alpha}(C), \lambda^{\mu}(D)\}, \min\{\gamma^{\alpha}(D), \lambda^{\mu}(E)\}\}$

$= \min\{(\gamma^{\alpha} \cap \lambda^{\mu})(C, D), (\gamma^{\alpha} \cap \lambda^{\mu})(E, F)\}$

$(\gamma^{\alpha} \cap \lambda^{\mu})(A - B) \geq \min\{(\gamma^{\alpha} \cap \lambda^{\mu})(A), (\gamma^{\alpha} \cap \lambda\lambda^{\mu})(B)\}.$

b) $(\gamma^{\alpha} \cap \lambda^{\mu})(AB) = (\gamma^{\alpha} \cap \lambda^{\mu})((C, D) \cdot (E, F))$

$= (\gamma^{\alpha} \cap \lambda^{\mu})(CE, DF)$

$= \min\{\gamma^{\alpha}(CE), \lambda^{\mu}(DF)\}$

$\geq \min\{\gamma^{\alpha}(C), \lambda^{\mu}(D)\}$

$= (\gamma^{\alpha} \cap \lambda^{\mu})(C, D)$

$(\gamma^{\alpha} \cap \lambda^{\mu})(AB) \geq (\gamma^{\alpha} \cap \lambda^{\mu})(A).$

c) $(\gamma^{\beta} \cup \lambda^{\eta})(A - B) = \max\{\gamma^{\beta}(C - E), \lambda^{\eta}(D - F)\}$

$\leq \max\{\max\{\gamma^{\beta}(C), \gamma^{\beta}(E)\}, \max\{\lambda^{\eta}(D), \lambda^{\eta}(F)\}\}$

$= \max\{\max\{\gamma^{\beta}(C), \lambda^{\eta}(D)\}, \max\{\gamma^{\beta}(E), \lambda^{\eta}(F)\}\}$

$= \max\{(\gamma^{\beta} \cup \lambda^{\eta})(C, D), (\gamma^{\beta} \cup \lambda^{\eta})(E, F)\}$

$(\gamma^{\beta} \cup \lambda^{\mu})(A - B) \leq \max\{(\gamma^{\beta} \cup \gamma^{\eta})(A), (\gamma^{\beta} \cup \lambda^{\eta})(B)\}.$

d) $(\gamma^{\beta} \cup \lambda^{\eta})(AB) = (\gamma^{\beta} \cup \lambda^{\eta})((C, D) \cdot (E, F))$

$= (\gamma^{\beta} \cup \lambda^{\eta})(CE, DF)$

$= \max\{\gamma^{\beta}(CE), \lambda^{\eta}(DF)\}$

$\leq \max\{\gamma^{\beta}(C), \lambda^{\eta}(D)\}$

©IJRASET (UGC Approved Journal): All Rights are Reserved

$= \max \{\gamma^{\beta}(C), \lambda^{\eta}(D)\}$

$= (\gamma^{\beta} \cup \lambda^{\eta})(C, D)$

$(\gamma^{\beta} \cup \lambda^{\mu})(AB) \leq (\gamma^{\beta} \cup \gamma^{\eta})(C, D).$

Hence, $\gamma^{G} \times \lambda^{H}$ is an intuitionistic fuzzy HX right ideal of a HX ring $\mathfrak{R}$.

K. Theorem

Let G and H be any two intuitionistic fuzzy sets of R_1 and R_2 respectively. Let $\mathfrak{R}_1 \subset 2^{R}{}_1 - \{\phi\}$ and $\mathfrak{R}_2 \subset 2^{R}{}_2 - \{\phi\}$ be any two HX rings. If γ^{G} and λ^{H} are any two intuitionistic fuzzy HX left ideals of $\mathfrak{R}_1$ and $\mathfrak{R}_2$ respectively then, $\gamma^{G} \times \lambda^{H}$ is also an intuitionistic fuzzy HX left ideal of a HX ring $\mathfrak{R}_1 \times \mathfrak{R}_2$.

1) Proof: It is clear.

L. Definition

Let $\lambda^{H} = \{\langle A, \lambda^{\mu}(A), \lambda^{\gamma}(A) \rangle / \text{ for all } A \in \mathfrak{I}\}$ be an intuitionistic fuzzy subset of a HX ring $\mathfrak{I}$. We define the following "necessity" and "possibility" operations :

$\Box \lambda^{H} = \{ \langle A, \lambda^{\mu}(A), 1 - \lambda^{\mu}(A) \rangle / A \in \mathfrak{I}\}.$

$\Diamond \lambda^{H} = \{ \langle A, 1 - \lambda^{\gamma}(A), \lambda^{\gamma}(A) \rangle / A \in \mathfrak{I}\}$

M. Theorem

Let H be an intuitionistic fuzzy set on R. Let λ^{H} be an intuitionistic fuzzy HX right ideal of a HX ring $\mathfrak{R}$ then $\Box\lambda^{H}$ is an intuitionistic fuzzy HX right ideal of a HX ring $\mathfrak{R}$.

1) Proof: Let λ^{H} be an intuitionistic fuzzy HX right ideal of a HX ring $\mathfrak{R}$. Then,

a) $\lambda^{\mu}(A - B) \geq \min \{ \lambda^{\mu}(A), \lambda^{\mu}(B) \}$

b) $\lambda^{\mu}(AB) \geq \lambda^{\mu}(A$

c) $\lambda^{\eta}(A - B) \leq \max \{ \lambda^{\eta}(A), \lambda^{\eta}(B) \}$

d) $\lambda^{\eta}(AB) \leq \lambda^{\eta}(A).$

Now, $\lambda^{\mu}(A - B) \geq \min \{ \lambda^{\mu}(A), \lambda^{\mu}(B) \}$

$1 - \lambda^{\mu}(A - B) \leq 1 - \min \{ \lambda^{\mu}(A), \lambda^{\mu}(B) \}$

$\leq \max \{ 1 - \lambda^{\mu}(A), 1 - \lambda^{\mu}(B) \}.$

That is, $1 - \lambda^{\mu}(A - B) \leq \max \{ 1 - \lambda^{\mu}(A), 1 - \lambda^{\mu}(B) \}.$

We have, $\lambda^{\mu}(AB) \geq \lambda^{\mu}(A)$

$1 - \lambda^{\mu}(AB) \leq 1 - \lambda^{\mu}(A).$

Hence, $\Box\lambda^{H}$ is an intuitionistic fuzzy HX right ideal of a HX ring $\mathfrak{R}$.

N. Theorem

Let H be an intuitionistic fuzzy set on R. Let λ^{H} be an intuitionistic fuzzy HX left ideal of a HX ring $\mathfrak{R}$ then $\Box\lambda^{H}$ is an intuitionistic fuzzy HX left ideal of a HX ring $\mathfrak{R}$

O. Theorem

Let H be an intuitionistic fuzzy set on R. Let λ^{H} be an intuitionistic fuzzy HX right ideal of a HX ring $\mathfrak{R}$ then $\Diamond\lambda^{H}$ is an intuitionistic fuzzy HX right ideal of a HX ring $\mathfrak{R}$.

1) Proof: Let λ^{H} be an intuitionistic fuzzy HX right ideal of a HX ring $\mathfrak{R}$. Then,

a) $\lambda^{\mu}(A - B) \geq \min \{ \lambda^{\mu}(A), \lambda^{\mu}(B) \}$,

b) $\lambda^{\mu}(AB) \geq \lambda^{\mu}(A)$

c) $\lambda^{\eta}(A - B) \leq \max \{ \lambda^{\eta}(A), \lambda^{\eta}(B) \}$

d) $\lambda^{\eta}(AB) \leq \lambda^{\eta}(A).$

Now, $\lambda^{\eta}(A - B) \leq \max \{ \lambda^{\eta}(A), \lambda^{\eta}(B) \}$

$1 - \lambda^{\eta}(A - B) \geq 1 - \max \{ \lambda^{\eta}(A), \lambda^{\eta}(B) \}$

$\geq \min \{ 1 - \lambda^{\eta}(A), 1 - \lambda^{\eta}(B) \}.$

That is, $1 - \lambda^{\eta}(A - B) \geq \min \{ 1 - \lambda^{\eta}(A), 1 - \lambda^{\eta}(B) \}.$

We have, $\lambda^{\eta}(AB) \leq \lambda^{\eta}(A)$

$1 - \lambda^{\eta}(AB) \geq 1 - \lambda^{\eta}(A).$

©IJRASET (UGC Approved Journal): All Rights are Reserved

Hence, $\Diamond\lambda^H$ is an intuitionistic fuzzy HX right ideal of a HX ring $\Re$.

P. Theorem

H be an intuitionistic fuzzy set on R. Let λ^H be an intuitionistic fuzzy HX left ideal of a HX ring $\Re$ then $\Diamond\lambda^H$ is an intuitionistic fuzzy HX left ideal of a HX ring $\Re$.

1) Proof: It is clear.

III. HOMOMORPHISM AND ANTI HOMOMORPHISM OF AN INTUITIONISTIC FUZZY HX RIGHT AND LEFT IDEAL OF A HX RING

In this section, we introduce the concept of an image, pre-image of an intuitionistic fuzzy subset of a HX ring and discuss the properties of homomorphic and anti homomorphic images and pre-images of an intuitionistic fuzzy HX right and left ideal of a HX ring $\Re$.

A. Theorem

Let $\Re_1$ and $\Re_2$ be any two HX rings on the rings R_1 and R_2 respectively. Let $f : \Re_1 \to \Re_2$ be a homomorphism onto HX rings. Let H be an intuitionistic fuzzy subset of R_1. Let λ^H be an intuitionistic fuzzy HX right ideal of $\Re_1$ then $f(\lambda^H)$ is an intuitionistic fuzzy HX right ideal of $\Re_2$, if λ^H has a supremum property and λ^H is f-invariant.

1) Proof: Let $H = \{ \langle x, \mu(x), \eta(x)\rangle / x \in R_1 \}$ be an intuitionistic fuzzy sets defined on a ring R_1.

Then, $\lambda^H = \{ \langle X, \lambda^\mu(X), \lambda^\eta(X)\rangle / X \in \Re_1 \}$ be an intuitionistic fuzzy HX right ideal of a HX ring $\Re_1$. Then, $f(\lambda^H) = \{ \langle f(X), f(\lambda^\mu)(f(X)), f(\lambda^\eta)(f(X))\rangle / X \in \Re_1 \}$.

There exist $X, Y \in \Re_1$ such that $f(X), f(Y) \in \Re_2$,

$$
\begin{aligned}
(f(\lambda^\mu))(f(X) - f(Y)) &= (f(\lambda^\mu))(f(X-Y)), \\
&= \lambda^\mu(X-Y) \\
&\geq \min\{\lambda^\mu(X), \lambda^\mu(Y)\} \\
&= \min\{(f(\lambda^\mu))(f(X)), (f(\lambda^\mu))(f(Y))\} \\
(f(\lambda^\mu))(f(X) - f(Y)) &\geq \min\{(f(\lambda^\mu))(f(X)), (f(\lambda^\mu))(f(Y))\}
\end{aligned}
$$

$$
\begin{aligned}
(f(\lambda^\mu))(f(X)\,f(Y)) &= (f(\lambda^\mu))(f(XY)) \\
&= \lambda^\mu(XY) \\
&\geq \lambda^\mu(X) \\
&= (f(\lambda^\mu))(f(X)) \\
(f(\lambda^\mu))(f(X)f(Y)) &\geq f(\lambda^\mu))(f(X)).
\end{aligned}
$$

$$
\begin{aligned}
(f(\lambda^\eta))(f(X) - f(Y)) &= (f(\lambda^\eta))(f(X-Y)), \\
&= \lambda^\eta(X-Y) \\
&\leq \max\{\lambda^\eta(X), \lambda^\eta(Y)\} \\
&= \max\{(f(\lambda^\eta))(f(X)), (f(\lambda^\eta))(f(Y))\} \\
(f(\lambda^\eta))(f(X) - f(Y)) &\leq \max\{(f(\lambda^\eta))(f(X)), (f(\lambda^\eta))(f(Y))\}
\end{aligned}
$$

$$
\begin{aligned}
(f(\lambda^\eta))(f(X)\,f(Y)) &= (f(\lambda^\eta))(f(XY)) \\
&= \lambda^\eta(XY) \\
&\leq \lambda^\eta(X) \\
&= (f(\lambda^\eta))(f(X)) \\
(f(\lambda^\eta))(f(X)f(Y)) &\leq f(\lambda^\eta))(f(X)).
\end{aligned}
$$

Hence, $f(\lambda^H)$ is an intuitionistic fuzzy HX right ideal of $\Re_2$.

B. Theorem

Let $\Re_1$ and $\Re_2$ be any two HX rings on the rings R_1 and R_2 respectively. Let $f : \Re_1 \to \Re_2$ be a homomorphism onto HX rings. Let H be an intuitionistic fuzzy subset of R_1. Let λ^H be an intuitionistic fuzzy HX left ideal of $\Re_1$ then $f(\lambda^H)$ is an intuitionistic fuzzy HX left ideal of $\Re_2$, if λ^H has a supremum property and λ^H is f-invariant.

1) Proof: It is clear.

©IJRASET (UGC Approved Journal): All Rights are Reserved

C. Theorem

Let $\Re_1$ and $\Re_2$ be any two HX rings on R_1 and R_2 respectively. Let $f : \Re_1 \to \Re_2$ be a homomorphism on HX rings. Let G be an intuitionistic fuzzy subset of R_2. Let γ^G be an intuitionistic fuzzy HX right ideal of $\Re_2$, then $f^{-1}(\gamma^G)$ is an intuitionistic fuzzy HX right ideal of $\Re_1$.

1) Proof: Let $G = \{ \langle y, \alpha(y), \beta(y)\rangle / y \in R_2 \}$ be an intuitionistic fuzzy sets defined on a ring R_2.

Then, $\gamma^G = \{ \langle Y, \gamma^\alpha(Y), \gamma^\beta(Y)\rangle / Y \in \Re_2 \}$ be an intuitionistic fuzzy HX right ideal of a HX ring $\Re_2$. Then, $f^{-1}(\gamma^G) = \{ \langle X, f^{-1}(\gamma^\alpha)(X), f^{-1}(\gamma^\beta)(X)\rangle / X \in \Re_1 \}$.

For any $X,Y \in \Re_1$, $f(X), f(Y) \in \Re_2$,

$$(f^{-1}(\gamma^\alpha))(X-Y) = \gamma^\alpha(f(X-Y))$$
$$= \gamma^\alpha(f(X) - f(Y))$$
$$\geq \min\{\gamma^\alpha(f(X)), \gamma^\alpha(f(Y))\}$$
$$= \min\{(f^{-1}(\gamma^\alpha))(X), (f^{-1}(\gamma^\alpha))(Y)\}$$
$$(f^{-1}(\gamma^\alpha))(X-Y) \geq \min\{(f^{-1}(\gamma^\alpha))(X), (f^{-1}(\gamma^\alpha))(Y)\}.$$

$$(f^{-1}(\gamma^\alpha))(XY) = \gamma^\alpha(f(XY))$$
$$= \gamma^\alpha(f(X)\, f(Y))$$
$$\geq \gamma^\alpha(f(X))$$
$$= (f^{-1}(\gamma^\alpha))(X)$$
$$(f^{-1}(\gamma^\alpha))(XY) \geq (f^{-1}(\gamma^\alpha))(X).$$

$$(f^{-1}(\gamma^\beta))(X-Y) = \gamma^\beta(f(X-Y))$$
$$= \gamma^\beta(f(X) - f(Y))$$
$$\leq \max\{\gamma^\beta(f(X)), \gamma^\beta(f(Y))\}$$
$$= \max\{(f^{-1}(\gamma^\beta))(X), (f^{-1}(\gamma^\beta))(Y)\}$$
$$(f^{-1}(\gamma^\beta))(X-Y) \leq \max\{(f^{-1}(\gamma^\beta))(X), (f^{-1}(\gamma^\beta))(Y)\}.$$

$$(f^{-1}(\gamma^\beta))(XY) = \gamma^\beta(f(XY))$$
$$= \gamma^\beta(f(X)\, f(Y))$$
$$\leq \gamma^\beta(f(X))$$
$$= (f^{-1}(\gamma^\beta))(X)$$
$$(f^{-1}(\gamma^\beta))(XY) \leq (f^{-1}(\gamma^\beta))(X).$$

Hence, $f^{-1}(\gamma^G)$ is an intuitionistic fuzzy HX right ideal of $\Re_1$.

D. Theorem

Let $\Re_1$ and $\Re_2$ be any two HX rings on R_1 and R_2 respectively. Let $f : \Re_1 \to \Re_2$ be a homomorphism on HX rings. Let G be an intuitionistic fuzzy subset of R_2. Let γ^G be an intuitionistic fuzzy HX left ideal of $\Re_2$, then $f^{-1}(\gamma^G)$ is an intuitionistic fuzzy HX left ideal of $\Re_1$.

1) Proof: It is clear.

E. Theorem

Let $\Re_1$ and $\Re_2$ be any two HX rings on the rings R_1 and R_2 respectively. Let $f : \Re_1 \to \Re_2$ be an anti homomorphism onto HX rings. Let H be an intuitionistic fuzzy subset of R_1. Let λ^H be an intuitionistic fuzzy HX right ideal of $\Re_1$ then $f(\lambda^H)$ is an intuitionistic fuzzy HX left ideal of $\Re_2$, if λ^H has a supremum property and λ^H is f-invariant.

1) Proof: Let $H = \{ \langle x, \mu(x), \eta(x)\rangle / x \in R \}$ be an intuitionistic fuzzy sets defined on a ring R_1. Then, $\lambda^H = \{ \langle X, \lambda^\mu(X), \lambda^\eta(X)\rangle / X \in \Re \}$ be an intuitionistic fuzzy HX right ideal of a HX ring $\Re_1$. Then, $f(\lambda^H) = \{ \langle f(X), f(\lambda^\mu)(f(X)), f(\lambda^\eta)(f(X))\rangle / X \in \Re \}$.

There exist $X,Y \in \Re_1$ such that $f(X), f(Y) \in \Re_2$,

©IJRASET (UGC Approved Journal): All Rights are Reserved

a) $(f(\lambda^{\mu}))(f(X) - f(Y)) = (f(\lambda^{\mu}))(f(X-Y))$,

$= \lambda^{\mu}(X-Y)$

$= \min\{\lambda^{\mu}(X), \lambda^{\mu}(Y)\}$

$= \min\{(f(\lambda^{\mu}))(f(X)), (f(\lambda^{\mu}))(f(Y))\}$

$(f(\lambda^{\mu}))(f(X) - f(Y)) \geq \min\{(f(\lambda^{\mu}))(f(X)), (f(\lambda^{\mu}))(f(Y))\}$

b) $(f(\lambda^{\mu}))(f(X)\, f(Y)) = (f(\lambda^{\mu}))(f(YX))$

$= \lambda^{\mu}(YX)$

$\geq \lambda^{\mu}(Y)$

$= (f(\lambda^{\mu}))(f(Y))$

$(f(\lambda^{\mu}))(f(X)f(Y)) \geq (f(\lambda^{\mu}))(f(Y))$.

c) $(f(\lambda^{\eta}))(f(X) - f(Y)) = (f(\lambda^{\eta}))(f(X-Y))$,

$= \lambda^{\eta}(X-Y)$

$\leq \max\{\lambda^{\eta}(X), \lambda^{\eta}(Y)\}$

$= \max\{(f(\lambda^{\eta}))(f(X)), (f(\lambda^{\eta}))(f(Y))\}$

$(f(\lambda^{\eta}))(f(X) - f(Y)) \leq \max\{(f(\lambda^{\eta}))(f(X)), (f(\lambda^{\eta}))(f(Y))\}$

d) $(f(\lambda^{\eta}))(f(X)\, f(Y)) = (f(\lambda^{\eta}))(f(YX))$

$= \lambda^{\eta}(YX)$

$\leq \lambda^{\eta}(Y)$

$= (f(\lambda^{\eta}))(f(Y))$

$(f(\lambda^{\eta}))(f(X)f(Y)) \leq (f(\lambda^{\eta}))(f(Y))$.

Hence, $f(\lambda^{H})$ is an intuitionistic fuzzy HX left ideal of $\Re_2$.

F. Theorem

Let $\Re_1$ and $\Re_2$ be any two HX rings on the rings R_1 and R_2 respectively. Let $f : \Re_1 \to \Re_2$ be an anti homomorphism onto HX rings. Let H be an intuitionistic fuzzy subset of R_1. Let λ^{H} be an intuitionistic fuzzy HX left ideal of $\Re_1$ then $f(\lambda^{H})$ is an intuitionistic fuzzy HX right ideal of $\Re_2$, if λ^{H} has a supremum property and λ^{H} is f-invariant.

1) Proof: It is clear.

G. Theorem

Let $\Re_1$ and $\Re_2$ be any two HX rings on R_1 and R_2 respectively. Let $f : \Re_1 \to \Re_2$ be an anti homomorphism on HX rings. Let G be an intuitionistic fuzzy subset of R_2. Let γ^{G} be an intuitionistic fuzzy HX right ideal of $\Re_2$, then $f^{-1}(\gamma^{G})$ is an intuitionistic fuzzy HX left ideal of $\Re_1$.

1) Proof: Let $G = \{ \langle y, \alpha(y), \beta(y) \rangle / y \in R_2 \}$ be an intuitionistic fuzzy sets defined on a ring R_2.

Then, $\gamma^{G} = \{ \langle Y, \gamma^{\alpha}(Y), \gamma^{\beta}(Y) \rangle / Y \in \Re_2 \}$ be an intuitionistic fuzzy HX right ideal of a HX ring $\Re_2$. Then, $f^{-1}(\gamma^{G}) = \{ \langle X, f^{-1}(\gamma^{\alpha})(X), f^{-1}(\gamma^{\beta})(X) \rangle / X \in \Re_1 \}$.

For any $X, Y \in \Re_1$, $f(X), f(Y) \in \Re_2$,

$(f^{-1}(\gamma^{\alpha}))(X-Y) = \gamma^{\alpha}(f(X-Y))$

$= \gamma^{\alpha}(f(X) - f(Y))$

$\geq \min\{\gamma^{\alpha}(f(X)), \gamma^{\alpha}(f(Y))\}$

$= \min\{(f^{-1}(\gamma^{\alpha}))(X), (f^{-1}(\gamma^{\alpha}))(Y)\}$

$(f^{-1}(\gamma^{\alpha}))(X-Y) \geq \min\{(f^{-1}(\gamma^{\alpha}))(X), (f^{-1}(\gamma^{\alpha}))(Y)\}$.

$(f^{-1}(\gamma^{\alpha}))(XY) = \gamma^{\alpha}(f(XY))$

$= \gamma^{\alpha}(f(Y)\, f(X))$

$\geq \gamma^{\alpha}(f(Y))$

$= (f^{-1}(\gamma^{\alpha}))(Y)$

©IJRASET (UGC Approved Journal): All Rights are Reserved

$(f^{-1}(\gamma^{\alpha}))(XY) \geq (f^{-1}(\gamma^{\alpha}))(Y).$

$(f^{-1}(\gamma^{\beta}))(X-Y) = \gamma^{\beta}(f(X-Y))$

$= \gamma^{\beta}(f(X) - f(Y))$

$\leq \max\{\gamma^{\beta}(f(X)), \gamma^{\beta}(f(Y))\}$

$= \max\{(f^{-1}(\gamma^{\beta}))(X), (f^{-1}(\gamma^{\beta}))(Y)\}$

$(f^{-1}(\gamma^{\beta}))(X-Y) \leq \max\{(f^{-1}(\gamma^{\beta}))(X), (f^{-1}(\gamma^{\beta}))(Y)\}.$

$(f^{-1}(\gamma^{\beta}))(XY) = \gamma^{\beta}(f(XY))$

$= \gamma^{\beta}(f(Y)\, f(X))$

$\leq \gamma^{\beta}(f(Y))$

$= (f^{-1}(\gamma^{\beta}))(Y)$

$(f^{-1}(\gamma^{\beta}))(XY) \leq (f^{-1}(\gamma^{\beta}))(Y).$

Hence, $f^{-1}(\gamma^{G})$ is an intuitionistic fuzzy HX left ideal of $\Re_1$.

H. Theorem

Let $\Re_1$ and $\Re_2$ be any two HX rings on R_1 and R_2 respectively. Let $f : \Re_1 \to \Re_2$ be a homomorphism on HX rings. Let G be an intuitionistic fuzzy subset of R_2. Let γ^{G} be an intuitionistic fuzzy HX left ideal of $\Re_2$, then $f^{-1}(\gamma^{G})$ is an intuitionistic fuzzy HX right ideal of $\Re_1$.

1) Proof: It is clear.


IV. ACKNOWLEDGEMENTS

The authors would like to express their sincere thanks to all our friends for their help to make this paper a successful one.

REFERENCES

[1] Atanassov.K.T, intuitionistic fuzzy sets,Fuzzy Sets and System 20 (1986) ,pp 87-96.
[2] Bing-xueYao and Yubin-Zhong, Upgrade of algebraic structure of ring, Fuzzy information and Engineering (2009)2:219-228.
[3] Bing-xueYao and Yubin-Zhong, The construction of power ring, Fuzzy information and Engineering (ICFIE),ASC 40,pp.181-187,2007
[4] International Journal of computational cognition, Volume 9, No.2, 45-50, June 2011.
[5] Li Hong Xing, HX group, BUSEFAL, 33(4), 31-37, October 1987.
[6] Li Hong Xing, HX ring, BUSEFAL, 34(1) 3-8, January 1988.
[7] Liu.W.J., Fuzzy invariant subgroups and fuzzy ideals, Fuzzy sets and systems,8,133-139
[8] Muthuraj.R, Manikandan K.H., Sithar Selvam.P.M., Intuitionistic Q-fuzzy normal HX group ,International Journal of Mathematical Archive , Vol.2,no10.,pp 1828-1832.
[9] Muthuraj.R, Manikandan K.H., Intuitionistic Q-fuzzy HX subgroup and $< \alpha, \beta >$ level sub HX groups, CIIT International DOI – FS 122011010.
[10] Muthuraj.R and Muthuraman.M.S, Intuitionistic anti fuzzy semiprimary HX ideal in HX ring, International journal of applied engineering research, 2015, vol.10, no.39, pp. 30050 – 30056. (Annexure II - SCI)
[11] Muthuraj.R and Muthuraman.M.S, Intuitionistic fuzzy HX k – ideal of a semi HX ring, SS International journal of pure and applied mathematics, 2015, pp. 31-41.
[12] Muthuraj.R, Ramila Gandhi.N., Induced fuzzy k-ideal and its level k-ideals on a HX ring ,IOSR Journal of Mathematics (IOSR-JM) Volume 9, Issue 6 (Jan 2014) , pp v138-143.
[13] Rosenfeld. A., Fuzzy groups,J.Math.Anal.,35(1971),512-517.
[14] Zadeh.L.A., Fuzzy sets, Information and control,8,338-353.

©IJRASET (UGC Approved Journal): All Rights are Reserved



10.22214/IJRASET

45.98

IMPACT FACTOR: 7.129

IMPACT FACTOR: 7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)