



IJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 2017 **Issue:** conference **Month of publication:** December 2017

DOI:

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

ON (k,d) – HERONIAN MEAN GRAPHS

Akilandeswari. K¹, Tamilselvi. M²

¹Research Scholar, ²Associate Professor, PG and Research Department of Mathematics, Seethalakshmi Ramaswami College, Tiruchirappalli – 620002

Abstract: Mean labeling was first introduced by S. Somasundram and R. Ponraj. Heronian mean labeling was introduced by S. S. Sandhya, E. Ebin Raja Merly and S. D. Deepa. We have extended this notion to a labeling called k-Heronian mean labeling. In this paper, we introduce (k,d)-Heronian mean labeling of some graphs. Here k and d are denoted as any positive integer greater than or equal to 1.

Keywords: (k,d)-Heronian mean labeling, (k,d)-Heronian mean graph, Path, $L_n \odot K_1$, $T_n \odot K_1$, $Q_n \odot K_1$, $TL_n \odot K_1$, H_n , Peterson graph.

2010 Mathematics Subject Classification: 05C78

I. INTRODUCTION

We begin with simple, finite, connected and undirected graph $G(V, E)$ with p vertices and q edges. For a detailed survey of graph labeling we refer to Gallian [2]. Terms are not defined here are used in the sense of Harary [3]. S. Somasundram and R. Ponraj were introduced mean labeling of graphs in [6] [7]. The concept of Heronian mean labeling was introduced by S.S. Sandhya et.al [4] [5].

We introduced the concept of k-Heronian mean labeling in [1]. In this paper, we introduce and investigate (k, d)-Heronian mean labeling of some graphs. For brevity, we use (k,d)-HML for (k,d)-Heronian mean labeling and k and d are any positive integer greater than or equal to 1.

A. Definition 1.1

A graph $G=(V,E)$ with p vertices and q edges is said to be a (k,d)-Heronian Mean graph if it is possible to label the vertices $x \in V$ with distinct labels $f(x)$ from $k, k+d, k+2d, \dots, k+qd$ in such a way that when each edge $e=uv$ is labeled with, $f^*(e) = \left\lfloor \frac{f(u)+f(v)+\sqrt{f(u)f(v)}}{3} \right\rfloor$ or $\left\lceil \frac{f(u)+f(v)+\sqrt{f(u)f(v)}}{3} \right\rceil$, then the resulting edge labels are distinct. In this case f is called a (k,d)-Heronian Mean labeling of G.

B. Definition 1.2

If G has order n, the corona of G with H, $G \odot H$ is the graph obtained by taking one copy of G and n copies of H and joining the i th vertex of G with an edge to every vertex in the i th copy of H

C. Definition 1.3

A triangular snake (T_n) is obtained from a path by identifying each edge of the path with an edge of the cycle C_3 .

D. Definition 1.4

The H – graph of a path P_n denoted by H_n is the graph obtained from two copies of P_n with vertices v_1, v_2, \dots, v_n and u_1, u_2, \dots, u_n by joining the vertices $v_{\frac{n+1}{2}}$ and $u_{\frac{n+1}{2}}$; if n is odd and the vertices $v_{\frac{n}{2}+1}$ and $u_{\frac{n}{2}}$; if n is even.

E. Definition 1.5

A Triangular Ladder $T(L_n)$ is a graph obtained from L_n by adding the edges $u_i v_{i+1}$, $1 \leq i \leq n-1$, where $1 \leq i \leq n$ are the vertices of L_n such that $u_1 u_2 u_3 \dots u_n$ and $v_1 v_2 v_3 \dots v_n$ are two paths of length n in the graph L_n .

F. Definition 1.6

The ladder graph L_n , is obtained from the cartesian product of two path graphs.

II. MAIN RESULTS

A. Theorem 2.1

Any path P_n is a (k, d) -Heronian mean graph, for all $n \geq 2$.

Proof:

Let $V(P_n) = \{u_i; 1 \leq i \leq n\}$ and $E(P_n) = \{e_i = (u_i, u_{i+1}); 1 \leq i \leq n - 1\}$ be the vertices and edges of P_n respectively.

Define $f: V(P_n) \rightarrow \{k, k + d, k + 2d, \dots, k + (n - 1)d\}$ by

$$f(u_i) = k + d(i - 1); 1 \leq i \leq n.$$

Now the induced edge labels are

$$f^*(e_i) = k + d(i - 1); 1 \leq i \leq n - 1.$$

Here $p = n$ and $q = n - 1$.

Clearly, f is (k, d) -Heronian mean labeling of P_n .

Hence P_n is a (k, d) -Heronian mean graph, for all $n \geq 2$.

1) Example 2.2: $(50, 2)$ -Heronian mean labeling of P_5 is given in the figure 2.1:



Figure 2.1: $(50, 2)$ -HML of P_5

B. Theorem 2.3

The graph $L_n \odot K_1$ is a (k, d) -Heronian mean labeling, for all $n \geq 2$.

Proof;

Let $V(L_n \odot K_1) = \{u_i, u'_i, v_i, v'_i; 1 \leq i \leq n\}$ and

$$E(L_n \odot K_1) = \{e_i = (u_i, u'_i), e'_i = (u'_i, v'_i), e''_i = (v_i, v'_i); 1 \leq i \leq n\} \cup$$

$\{e_i''' = (u'_i, u'_{i+1}), e_i^{iv} = (v'_i, v'_{i+1}); 1 \leq i \leq n - 1\}$ be the vertices and edges of $L_n \odot K_1$ respectively.

Define $f: V(L_n \odot K_1) \rightarrow \{k, k + d, \dots, k + (5n - 2)d\}$ by

$$\begin{aligned} f(u_1) &= k \\ f(u_i) &= k + d(5i - 6); 2 \leq i \leq n \\ f(u'_1) &= k + d \\ f(u'_i) &= k + 5d(i - 1); 2 \leq i \leq n \\ f(v'_1) &= k + 2d \\ f(v'_i) &= k + d(5i - 4); 2 \leq i \leq n \\ f(v_i) &= k + d(5i - 2); 1 \leq i \leq n \end{aligned}$$

Now the induced edge labels are

$$\begin{aligned} f^*(e_i) &= k + 5d(i - 1); 1 \leq i \leq n \\ f^*(e'_i) &= k + d(5i - 4); 1 \leq i \leq n \\ f^*(e''_i) &= k + d(5i - 3); 1 \leq i \leq n \\ f^*(e_i''') &= k + d(5i - 2); 1 \leq i \leq n - 1 \\ f^*(e_i^{iv}) &= k + d(5i - 1); 1 \leq i \leq n - 1 \end{aligned}$$

Here $p = 4n$ and $q = 5n - 2$.

Clearly f is a (k, d) -Heronian mean labeling.

Hence $L_n \odot K_1$ is a (k, d) -Heronian mean graph for all $n \geq 2$.

1) Example 2.4

$(75, 2)$ -Heronian mean labeling of $\square_7 \odot \square_7$ is given in figure 2.2:




Figure 2.2: (75, 2)-HML of $L_7 \odot K_1$

C. Theorem 2.5

The graph $T_n \odot K_1$ is a (k,d)- Heronian mean labeling, for all $n \geq 2$.

Proof;

Let $V(T_n \odot K_1) = \{u_i, v_i; 1 \leq i \leq n\} \cup \{w_i, x_i; 1 \leq i \leq n-1\}$ and

$$E(T_n \odot K_1) = \{e_i = (u_i, v_i), \quad 1 \leq i \leq n\} \cup \{e'_i = (v_i, w_i), \quad e''_i = (w_i, x_i), e'''_i = (v_i, v_{i+1}), e_i^{iv} = (w_i, x_i); 1 \leq i \leq n-1\}$$

be the vertices and edges of $T_n \odot K_1$ respectively.

Define $f: V(T_n \odot K_1) \rightarrow \{k, k + d, \dots, k + (5n - 4)d\}$ by

$$\begin{aligned} f(u_i) &= k + 5d(i - 1); \quad 1 \leq i \leq n \\ f(v_i) &= k + d(5i - 4); \quad 1 \leq i \leq n \\ f(w_i) &= k + d(5i - 3); \quad 1 \leq i \leq n - 1 \\ f(x_i) &= k + d(5i - 2); \quad 1 \leq i \leq n - 1 \end{aligned}$$

Now the induced edge labels are

$$\begin{aligned} f^*(e_i) &= k + 5d(i - 1); \quad 1 \leq i \leq n \\ f^*(e'_i) &= k + d(5i - 4); \quad 1 \leq i \leq n - 1 \\ f^*(e''_i) &= k + d(5i - 3); \quad 1 \leq i \leq n - 1 \\ f^*(e'''_i) &= k + d(5i - 2); \quad 1 \leq i \leq n - 1 \\ f^*(e_i^{iv}) &= k + d(5i - 1); \quad 1 \leq i \leq n - 1 \end{aligned}$$

Here $p = 4n-2$ and $q = 5n-4$.

Clearly f is a (k,d)- Heronian mean labeling.

Hence $T_n \odot K_1$ is a (k,d)-Heronian mean graph for all $n \geq 2$.

1) Example 2.6: (150,3)-Heronian mean labeling of $T_6 \odot K_1$ is given in figure 2.3:




Figure 2.3: (150,3)-HML of $T_6 \odot K_1$

D. Theorem 2.7

The graph $Q_n \odot K_1$ is a (k,d) - Heronian mean labeling, for all $n \geq 2$.

Proof;

Let $V(Q_n \odot K_1) = \{u_i, v_i; 1 \leq i \leq n\} \cup \{w_i, x_i, w'_i, x'_i; 1 \leq i \leq n-1\}$ and $E(Q_n \odot K_1) = \{e_i = (u_i, v_i), 1 \leq i \leq n\} \cup \{e'_i = (v_i, w_i), e''_i = (w_i, x_i), e'''_i = (w_i, w'_i), e^{iv}_i = (w'_i, x'_i), e^v_i = (v_i, v_{i+1}), e^{vi}_i = (w'_i, x_i); 1 \leq i \leq n-1\}$ be the vertices and edges of $Q_n \odot K_1$ respectively.

Define $f: V(Q_n \odot K_1) \rightarrow \{k, k + d, \dots, k + (7n - 6)d\}$ by

$$\begin{aligned} f(u_i) &= k + 7d(i - 1); 1 \leq i \leq n \\ f(v_i) &= k + d(7i - 6); 1 \leq i \leq n \\ f(w_i) &= k + d(7i - 5); 1 \leq i \leq n - 1 \\ f(x_i) &= k + d(7i - 4); 1 \leq i \leq n - 1 \\ f(w'_i) &= k + d(7i - 3); 1 \leq i \leq n - 1 \\ f(x'_i) &= k + d(7i - 2); 1 \leq i \leq n - 1 \end{aligned}$$

Now the induced edge labels are

$$\begin{aligned} f^*(e_i) &= k + 7d(i - 1); 1 \leq i \leq n \\ f^*(e'_i) &= k + d(7i - 6); 1 \leq i \leq n - 1 \\ f^*(e''_i) &= k + d(7i - 5); 1 \leq i \leq n - 1 \\ f^*(e'''_i) &= k + d(7i - 4); 1 \leq i \leq n - 1 \\ f^*(e^{iv}_i) &= k + d(7i - 3); 1 \leq i \leq n - 1 \\ f^*(e^v_i) &= k + d(7i - 2); 1 \leq i \leq n - 1 \\ f^*(e^{vi}_i) &= k + d(7i - 1); 1 \leq i \leq n - 1 \end{aligned}$$

Here $p = 6n-4$ and $q = 7n-6$.

Clearly f is a (k,d) - Heronian mean labeling.

Hence $Q_n \odot K_1$ is a (k,d) -Heronian mean graph for all $n \geq 2$.

1) Example 2.8: (200,6)-Heronian mean labeling of $Q_4 \odot K_1$ is given in figure 2.4:




Figure 2.4: (200, 6)-HML of $Q_4 \odot K_1$

E. Theorem 2.9

The Total graph $TL_n \odot K_1$ is a (k,d) - Heronian mean labeling, for all $n \geq 2$.

Proof:

Let $V(TL_n \odot K_1) = \{u_i, v_i, w_i, x_i; 1 \leq i \leq n\}$ and

$$E(TL_n \odot K_1) = \{e_i = (u_i, v_i), e'_i = (v_i, w_i), e''_i = (w_i, x_i); 1 \leq i \leq n\} \cup \{e'''_i = (v_i, v_{i+1}), e^{iv}_i = (v_i, w_{i+1}), e^v_i = (w_i, w_{i+1}); 1 \leq i \leq n-1\}$$

be the vertices and edges of $TL_n \odot K_1$ respectively.

Define $f: V(TL_n \odot K_1) \rightarrow \{k, k + d, \dots, k + (6n - 3)d\}$ by

$$\begin{aligned} f(u_i) &= k + 6d(i - 1); 1 \leq i \leq n \\ f(v_i) &= k + d(6i - 5); 1 \leq i \leq n \\ f(w_i) &= k + d(6i - 4); 1 \leq i \leq n \\ f(x_i) &= k + d(6i - 3); 1 \leq i \leq n \end{aligned}$$

Now the induced edge labels are

$$\begin{aligned} f^*(e_i) &= k + 6d(i - 1); 1 \leq i \leq n \\ f^*(e'_i) &= k + d(6i - 5); 1 \leq i \leq n \\ f^*(e''_i) &= k + d(6i - 4); 1 \leq i \leq n \\ f^*(e'''_i) &= k + d(6i - 3); 1 \leq i \leq n - 1 \\ f^*(e^{iv}_i) &= k + d(6i - 2); 1 \leq i \leq n - 1 \\ f^*(e^v_i) &= k + d(6i - 1); 1 \leq i \leq n - 1 \end{aligned}$$

Here $p = 4n$ and $q = 6n-3$.

Clearly f is a (k,d) - Heronian mean labeling.

Hence $TL_n \odot K_1$ is a (k,d) -Heronian mean graph for all $n \geq 2$.

1) Example 2.10

$(185,7)$ -Heronian mean labeling of $TL_7 \odot K_1$ is given in figure 2.5:




Figure 2.5: $(185,7)$ -HML of $TL_7 \odot K_1$

F. Theorem 2.11

The graph H_n is a (k,d) -Heronian mean graph for all $n \geq 2$.

Proof:

Let $V(H_n) = \{u_i, v_i; 1 \leq i \leq n\}$ and

$$E(H_n) = \{e_i = (u_i, u_{i+1}), e'_i = (v_i, v_{i+1}); 1 \leq i \leq n-1\} \cup \{e\}$$

be the vertices and edges of H_n respectively.

Define $f: V(H_n) \rightarrow \{k, k + d, \dots, k + (2n - 1)d\}$ by

$$\begin{aligned} f(u_i) &= k + d(i - 1); 1 \leq i \leq n \\ f(v_i) &= k + d(n + i - 1); 1 \leq i \leq n \end{aligned}$$

Now the induced edge labels are

$$\begin{aligned} f^*(e_i) &= k + d(i - 1); 1 \leq i \leq n - 1 \\ f^*(e'_i) &= k + d(n + i - 1); 1 \leq i \leq n - 1 \\ f^*(e) &= k + d(n - 1) \end{aligned}$$

Here $p = 2n$ and $q = 2n-1$.

Clearly f is a (k,d) -Heronian mean labeling.

Hence H_n is a (k,d) -Heronian mean graph for all $n \geq 2$.

1) *Example 2.12:* $(125,8)$ -Heronian mean labeling of H_6 is given in figure 2.6:




Figure 2.6: $(125, 8)$ -HML of H_6

G. Theorem 2.13

The Peterson graph is a (k,d) -Heronian mean graph.

Proof:

Let G be Peterson Graph.

Let $V(G) = \{u_i, v_i; 1 \leq i \leq 5\}$ and $E(G) = \{e_1 = (u_1, u_3), e_2 = (u_3, u_5), e_3 = (u_5, u_2), e_4 = (u_2, u_4)\} \cup \{e_5 = (u_4, u_1)\} \cup \{e'_i = (v_i, v_{i+1}); 1 \leq i \leq 4\} \cup \{e''_5 = (v_1, v_5)\} \cup \{e''_i = (v_i, u_i); 1 \leq i \leq 5\}$ be the vertices and edges of G .

Define $f: V(G) \rightarrow \{k, k + d, \dots, k + 15d\}$

$$f(u_i) = k + d(i - 1); 1 \leq i \leq 4$$

$$f(u_5) = k + 5d$$

$$f(v_i) = k + d(i + 9); 1 \leq i \leq 5$$

Now the induced edge labels are

$$f^*(e_i) = k + d(i - 1); 1 \leq i \leq 5$$

$$f^*(e'_i) = k + d(i + 9); 1 \leq i \leq 2$$

$$f^*(e'_i) = k + d(i + 10); 3 \leq i \leq 4$$

$$f^*(e'_5) = k + 12d$$

$$f^*(e''_i) = k + d(i + 4); 1 \leq i \leq 5$$

Here $p = 2n$ and $q = 2n-1$.

Clearly f is a (k,d) -Heronian mean labeling.

Hence H_n is a (k,d) -Heronian mean graph for all $n \geq 2$.

1) *Example 2.14*

$(10,9)$ -Heronian mean labeling of Peterson Graph.




Figure 2.7: (k,d)-HML of Peterson Graph

REFERENCES

- [1] Akilandeswari. K and Tamilselvi. M, "On k-Heronian Mean Labeling, International Journal of Innovative Research in Science, Engineering and Technology, Vol. 6, Issue 9, September 2017, 18019-18023.
- [2] J.A. Gallian, "A dynamic survey of graph labeling", Electronic Journal of Combinatorics, 18 (2016) # DS6
- [3] F. Harary, Graph Theory, Addison Wesley, Massachusetts (1972).
- [4] S.S.Sandhya, E.Ebin Raja Merly and S.D.Deepa, "Heronian Mean Labeling of Graphs", Communicated to International Journal of Mathematical Form.
- [5] S.S.Sandhya, E.Ebin Raja Merly and S.D.Deepa, "Some Results on Heronian Mean Labeling of Graphs", communicated to Journal of Discrete Mathematical Sciences and Cryptography.
- [6] S. Somasundaram and R. Ponraj, Mean labeling of graphs, National Academy Science Letter, 26 (2003), 210-213.
- [7] S.Somasundaram, R.Ponraj and S.S.Sandhya, "Harmonic Mean Labeling of Graphs", communicated to Journal of Combinatorial Mathematics and Combinatorial Computing.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)