



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 5 Issue: XII Month of publication: December 2017 DOI:

www.ijraset.com

Call: 🛇 08813907089 🕴 E-mail ID: ijraset@gmail.com



Domain and Time Domain Approaches in Forecasting Techniques

R.V.S.S. Nagabhushana Rao¹, B. Ramana Murthy², C.Narayana³, K.Vasu⁴, G. Mokesh Rayalu⁵

¹Assistant professor, Department of Statistics, Vikrama Simhapuri University, Nellore

²Assistant Professor, Department of Mathematics and Statistics, Acharya N.G. Ranga Agricultural University College, Tirupati.

³Assistant professor, Department of Mathematics, Sriharsha Institute of P.G. Studies, Nellore

⁴Assistant Professor, Vidya Educational Institutions, Yadamari, Chittoor

⁵Assistant Professor, Department of Mathematics, School of Advanced sciences, VIT, Vellore

Abstract: Time series analysis has been created by four types of experts namely Engineers and Physical Scientists; Economists; Applied Statisticians and Econometricians; and Mathematical Statisticians and Probability. The richness of the subject springs from this diversity of its origins. The history of time series analysis was not as smooth as one might think. Time series analysis was divided into two approaches namely Frequency Domain and Time Domain approaches. For Engineers and Physical Scientists, it is natural to regard a time series as made up of oscillations at different frequencies. This gives frequency domain analysis. For Statisticians and Applied research workers in other fields, it is natural to treat the data from the standpoint of correlation and regression relations between successive observations. This gives time domain analysis. There is duality between the frequency domain and time domain approaches to the time series analysis.

I. INTRODUCTION

In univariate time series regression models, the regressor variable is time. A linear trend relationship can be written as

$$Y = \alpha + \beta T + \epsilon \tag{1.1}$$

Where T indicates time. The T variable may be specified in many ways. when T has zero mean, the normal equations for fitting (1.1) will become

$$a = \overline{Y}and b = \frac{\sum Y}{\sum T^2}$$

One way of modeling such behavior is by means of auto regression. The simplest autoregressive model is

$$Y_{t} = \alpha + \beta Y_{t-1} + \epsilon_{t} \tag{1.2}$$

This is called a first order autoregressive model and is denoted by AR(1). The order indicates the maximum lag in the equation. From the equation (1.2), we make the following assumptions about \in variable

$$E\left(\epsilon_{i}^{2}\right) = \sigma^{2} \text{ for all I}$$

$$E\left(\epsilon_{i} \in j\right) = 0 \text{ for all } i \neq j$$

$$(1.3)$$

These assumptions define a white noise series. Here the crucial question is, how does the Y series behave over time. Assuming that process stated a very long time ago, we take

$$Y_{t} = \alpha \left(1 + \beta + \beta^{2} + \dots \right) + \left(\in_{t} + \beta \in_{t-1} + \beta^{2}_{t} \in_{t-2} + \dots \right)$$

$$E(Y_{t}) = \alpha \left(1 + \beta + \beta^{2} + \dots \right)$$

$$(1.4)$$

This expectation exists only when the infinite geometric series on the right hand side has limit. The necessary and sufficient conditions is $|\beta| < 1$.

The expectation then

then
$$E(Y_t) = \mu = \frac{\alpha}{1-\beta}$$
 (1.5)

f 11 :



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor :6.887

Volume 5 Issue XII December 2017- Available at www.ijraset.com

(1.7)

(2.1)

Here variance Y will be
$$\operatorname{Var}(Y) = \sigma_y^2 - \frac{\sigma^2}{1 - \beta^2}$$
 (1.6)

The Y series has a constant unconditional variance, independent of time. The covariance of Y with a lagged value itself is known as autocovariance. The lag auto covariance is defined as

 $\gamma_s = \beta^s \sigma_v^2$ s = 0,1,2,

So that first lag autocovariance is

 $\gamma_1 = \beta \sigma_y^z$

The autocovariances depend only on the lag length and are independent of t Dividing the covariances by the variance gives the autocorrelation coefficients, also known as series correlation coefficients. These will be defined as

$$\rho_{\rm s} = \gamma_{\rm s} / \gamma_{\rm o} , \qquad {\rm s} = 0, 1, 2..... \tag{1.8}$$

Where $\gamma_{o} = \sigma_{u}^{z}$

Plotting the autocorrelation coefficients against the lag lengths gives the correlogram of series.

When $|\beta| < 1$ the mean, variance, and covariances of Y series are constants independent of time. The Y series is then said to be weekly or covariance stationary.

When $\beta = 1$, the AR(1) process is said to have unit root. The equation becomes.

(1.9)

Which is called a random walk with drift. The conditional expectation and conditional variance are

 $E(Y_t/Y_0) = at + Y_o$

Which increases or decreases without limit as t increases $Var(Y_t/Y_0) = t\sigma^2$

Which increases without limit. In this case the unconditional mean and variance do not exist. The Y series is then said to be nonstationary.

A. Autoregressive Model With Order p:AR(p)

A common approach for modeling univariate time series is the autoregressive (AR) model. The AR(p) model is defined as

$$X_{t} = \delta + \phi X_{t-1} + \phi_{2} X_{t-2} + \dots + \phi_{p} X_{t-p} + \epsilon_{t}$$

where X_{t} is the time series, $\delta = \left(1 - \sum_{i=1}^{p} \phi_{i}\right) \mu$, where μ is the process mean

 \in_{t} is the white noise

 $\varphi_1,\varphi_2.....\varphi_p$ are the parameters of the model

p is the order of the AR model

Some constraints are necessary on the values of the parameters of this model so that model remains wide-sense stationary. For an

AR(P) model to be wide-sense stationary, the roots of the polynomial $Z^p - \sum_{i=1}^{p} \phi_i Z^{p-i}$ must lie within the unit circle, i.e., each root

 $Z_{i} \text{ must satisfy } \mid Z_{i} \mid > \mid .$

1) Estimation of AR Parameters: The AR(p) model is given by the equation

$$X_{t} = \sum_{i=1}^{p} \phi_{i} X_{t-i} + \epsilon_{t}$$

$$(2.2)$$

It is based on the parameters ϕ_i where i=1,, p.



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor :6.887 Volume 5 Issue XII December 2017- Available at www.ijraset.com

There is a direct correspondence between these parameters and the covariance function of the process. This correspondence can be inverted to determine the parameters from the autocorrelation function by using yule-walker equations.

$$\gamma_{m} = \sum_{k=l}^{p} \phi_{k} \gamma_{m-k} + \sigma_{t}^{2} \delta_{m,0}$$
(2.3)

where

 $m=0,1, \dots p.$ yielding p+1 equations.

 $\gamma_{\rm m}$ is the autocorrelation function of x

 $\boldsymbol{\sigma}_{_{\!\boldsymbol{\mathcal{C}}}}$ is the standard deviation of the input noise process

$$\delta_{m,0}$$
 is the Kronecker delta function

The last part of the equation is non-zero only if m=0. Hence the equation is solved by representing it as a matrix for m>0. Thus we get the equation.

$$\begin{bmatrix} \gamma_1 \\ \gamma_2 \\ \gamma_3 \\ \vdots \\ \vdots \end{bmatrix} = \begin{bmatrix} \gamma_0 & \gamma_{-1} & \gamma_{-2} & \dots \\ \gamma_1 & \gamma_0 & \gamma_{-1} & \dots \\ \gamma_2 & \gamma_1 & \gamma_0 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix} \begin{bmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \\ \vdots \end{bmatrix}$$

Solving all ϕ can be otained. For m =0, $\gamma_0 = \sum_{k=1}^p \phi_k \gamma_{-k} + \sigma_t^2$ Which allows us to solve σ_{\in}^2

1) Derivation: The equation defining the AR process is

$$X_t = \sum_{i=1}^p \phi_i X_{t-i} + \in_t$$

Multiplying both sides by X_{t-m} and taking expectation,

$$E\left[X_{t}X_{t-m}\right] = E\left[\sum_{i=1}^{p}\phi_{i}X_{t-i}X_{t-m}\right] + E\left[\epsilon_{t}X_{t-m}\right]$$

By the definition of autocorrelation function,

$$E\!\left[X_{t},X_{t-m}\right]\!=\!\gamma_{m}$$

The values of noise function are independent of each other and X_{t-m} is independent of \in_t when 'm' is greater than zero.

For m>0,
$$E[\in_{t} X_{t-m}] = 0$$

For m=0, $E[\in_{t} X_{t-m}], E\left[\in_{t} \left(\sum_{i=1}^{p} \phi_{i} X_{t-i} + \epsilon_{t}\right)\right]$
 $= \sum_{i=1}^{p} \phi_{i} E[\in_{t} X_{t-i}] + E\left[\epsilon_{t}^{2}\right]$
 $= 0 + \sigma_{t}^{2}$
 $= \sigma_{t}^{2}$
(2.4)
Now, we have for $m \ge 0$, $\gamma_{m} = E\left[\sum_{i=1}^{p} \phi_{i} X_{t-i} X_{t-m}\right] + \sigma_{t}^{2} \gamma_{m}$

©IJRASET (UGC Approved Journal): All Rights are Reserved



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor :6.887 Volume 5 Issue XII December 2017- Available at www.ijraset.com

$$\text{Further, } E \Bigg[\sum_{i=1}^{p} \phi_i X_{t-i} X_{t-m} \Bigg] = \sum_{i=1}^{p} \phi_i E \Big[X_t \ X_{t-m+i} \Big] \quad = \ \sum_{i=1}^{p} \phi_i \ \gamma_{m-i}$$

Which yields the Yule-Walkes equations.

$$\gamma_{m} = \sum_{i=1}^{p} \phi_{i} \gamma_{m-i} + \sigma_{t}^{2} \gamma_{m} \text{ for } m \ge 0$$
(2.5)

For m<0,
$$\gamma_{\rm m} = \gamma - {\rm m} = \sum_{i=1}^{P} \phi_i \gamma_{|{\rm m}|-i} + \sigma_{\epsilon}^2 \delta_{\rm m}$$
 (2.6)

2) Moving Average Model With Order Q: Ma (Q)

In time series analysis, the moving average (MA) model is common approach for modeling univariate time series models. The moving average model with order qMA(q) is

$$X_{t} = \mu + \epsilon_{t} + \theta_{1} \epsilon_{t-1} + \dots + \theta_{q} \epsilon_{t-q}$$

$$(3.1)$$

Where, μ is the mean of the series,

 $\theta_{1}, \theta_{2}, \dots, \theta_{q}$ are the parameters of the model,

 $\in_{t}, \in_{t-1}, \dots$ are white noise error terms.

q is the order of the moving average model.

B. Autoregressive Moving Average (Arma) Model

It is convenient to use the notation ARMA (p, q), where p is the order of the autoregressive part and q is the order of the moving average part.

The general AR (p) model was represented as

$$X_{t} = \phi_{1}X_{t-1} + \phi_{2}X_{t-2} + \dots + \phi_{p}X_{t-p} + \epsilon_{t}$$
(4.1)

Multiplying both sides by X_{t-k} yields

$$X_{t-k}X_{t} = \phi_{1}X_{t-k}X_{t-1} + \phi_{2}X_{t-k}X_{t-2} + \dots + \phi_{p}X_{t-k}X_{t-p} + X_{t-k} \in_{t}$$

Taking expected value both sides and assuming stationarity gives

$$\gamma_{k} = \phi_{1}\gamma_{k-1} + \phi_{2}\gamma_{k-2} + \dots + \phi_{p}\gamma_{k-p}$$

Where γ_k is the covariance between $\,X_t\,$ and $\,X_{t-k}\,$

The MA (q) model is written as

$$\mathbf{X}_{\mathbf{k}} = \mathbf{e}_{\mathbf{t}} - \mathbf{\theta}_{\mathbf{1}} \mathbf{e}_{\mathbf{t}-\mathbf{1}} - \mathbf{\theta}_{\mathbf{2}} \mathbf{e}_{\mathbf{t}-\mathbf{2}} \dots \dots \mathbf{\theta}_{\mathbf{q}} \mathbf{e}_{\mathbf{t}-\mathbf{q}}$$

Multiplying both sides by X_{t-k} yields

$$X_{t-k}X_{t} = \left(e_{t} - \theta_{1} e_{t-1} - \theta_{2} e_{t-2} \dots \theta_{q} e_{t-q}\right) X \left(e_{t-k} - \theta_{1} e_{t-k-1} - \theta_{2} e_{t-k-2} \dots \theta_{q} e_{t-k-q}\right) (4.2)$$

The expected value of the above equation will depend upon the value of k.

If k=0, and all other terms of the equation on (4.2) dropout because

$$E(e_t e_{t+i}) = 0 \text{ for } i \neq 0 \text{ and } E(e_t e_{t+i}) = \sigma_e^2 \text{ for } i = 0.$$

Thus (4.1) becomes, $\gamma_0 = \sigma_e^2 + \theta_1^2 \sigma_e^2 + \theta_2^2 \sigma_e^2 + \dots + \theta_q^2 \sigma_e^2$

To obtain the initial estimates for ARMA models, combine AR and MA models:

$$\gamma_{k} = \phi_{l} E(x_{t} x_{t-k}) + \dots + \phi_{p} E(X_{t-p} X_{t-k}) + E(e_{t} X_{t-k}) - \theta_{l} E(e_{t-1} X_{t-k}) - \dots - \theta_{q} E(e_{t-q} X_{t-k})$$
(4.3)

If k>q, the terms $E(e_t X_{t-k}) = 0$ which leaves



(4.5)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor :6.887 Volume 5 Issue XII December 2017- Available at www.ijraset.com

 $\gamma_k = \phi_1 \gamma_{k-1} + \phi_2 \gamma_{k-2} + \dots + \phi_p \gamma_{k-p}$

When k < q, the past errors and the X_{t-k} will be correlated and the autocovariances will be affected by the moving average part of the process, requiring that it will be included. The variance and autocovariances of an ARMA (1,1) process are therefore obtained as follows:

 $\boldsymbol{X}_t = \boldsymbol{\varphi}_1 \boldsymbol{X}_{t-1} + \boldsymbol{e}_t - \boldsymbol{\theta}_1 \, \boldsymbol{e}_{t-1}$

Multiplying both sides by X_{t-k} gives

$$X_{t-k}X_{t} = \phi_{l}X_{t-k}X_{t-l} + X_{t-k}e_{t} - \theta_{l}X_{t-k}e_{t-l}$$

Taking expected values both sides results in

$$E[X_{t-k}X_{t}] = \phi E[X_{t-k}X_{t-1}] + E[X_{t-k}e_{t}] - \theta_{1}E[X_{t-k}e_{t-1}]$$

If k=0, $\gamma_{0} = \theta_{1}\gamma_{1} + E[(\phi_{1}X_{t-1} + e_{t} - \theta_{1}e_{t-1})e_{t}] - \theta_{1}E[(\phi_{1}X_{t-1} + e_{t} - \theta_{1}e_{t-1})e_{t-1}]$ (4.4)
Since $x_{t} = \phi_{1}k_{t-1} + e_{t} - \theta_{1}e_{t-1}$

 $\gamma_0 = \phi_1 \gamma_1 + \sigma_e^2 - \theta_1 (\phi_1 - \theta_1) \sigma_e^2$ nilerly if k-1 $\gamma_e = \phi \gamma_e - \theta \sigma_e^2$

Similarly if k=1, $\gamma_1 = \phi_1 \gamma_0 - \theta_1 \sigma_e^2$

Solving the equations (4.2) ad (4.3) for γ_0 and γ_1 we get

$$\gamma_0 = \frac{1 + \theta_1^2 - 2\phi_1 \theta_1}{1 - \theta_1^2}$$
(4.6)

$$\gamma_1 = \frac{\left(1 - \phi_1 \theta_1\right) \left(\phi_1 - \theta_1\right)}{1 - \theta_1^2} \tag{4.7}$$

Dividing (4.5) by (4.4) gives

$$\rho_{1} = \frac{(1 - \phi_{1}\theta_{1})(\phi_{1} - \theta_{1})}{1 + \theta_{1}^{2} - 2\phi_{1}\theta_{1}}$$
(4.8)

C. Autoregressive Integrated Moving Average (Aima) Model

To identify the appropriate ARIMA model for a time series, one has to begin by identifying the order (r) of differencing needing to stationarize the series and remove the gross features of seasonality, in conjunction with a variance–stabilizing transformation such as lagging or deflating.

The equation for the simplest case ARIMA (1, 1, 1) is as follows

$$(1-B).(1-\phi_1B)X_t = \mu^1 + (1-\phi_1B)e_t$$

The terms can be multiplied out and rearranged as follows.

$$\begin{bmatrix} 1 - B(1 + \phi_1) + \phi_1 B^2 \end{bmatrix} x_t = \mu^1 + e_t - \theta_1 e_{t-1}$$

$$X_t = (1 + \phi_1) X_{t-1} - \phi_2 X_{t-2} + \mu^1 + e_t - \theta_1 e_{t-1}$$
(5.1)

In this form, the ARIMA model looks more like a conventional regression equation, except that there is more than one error term on the right hand side. ARIMA models are quite flexible and can represent a wide range of characteristics of time series that occur in practice.

1) ARIMA Model : pth order autoregressive model

Y.

$$= \phi_0 + \phi_1 Y_{t-1} + \phi_2 Y_{t-2} + \dots + \phi_p Y_{t-p} + \epsilon_t$$
(5.2)

2) ARIMA Model: qth order moving average model

$$Y_{t} = \mu + \in_{t} - W_{1} \in_{t-1} - W_{2} \in_{t-2} \dots W_{q} t_{t-q}$$
(5.3)



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor :6.887 Volume 5 Issue XII December 2017- Available at www.ijraset.com

(6.3)

3) ARIMA Model: ARMA (p, q) model

$$Y_{t} = \phi_{0} + \phi_{1}Y_{t-1} + \phi_{2}Y_{t-2} + \dots + \phi_{p}Y_{t-p} + \epsilon_{t}$$

-W₁ \epsilon_{t-1} - W_2 \epsilon_{t-2} \dots + W_q \epsilon_{t-q} (5.4)

$$\mathbf{Y}_{t} - \mathbf{Y}_{t-1} = \boldsymbol{\varepsilon}_{t} - \mathbf{W}_{1} \boldsymbol{\varepsilon}_{t-1} \tag{5.6}$$

D. Autoregresive Distributed Lag Models (Ardl)

In case where the variables in the long-run relation of interest are trend stationary, the general practice has been to de-trend the series and to model the detrended series as stationary distributed lag or autoregressive distributed lag (ARDL) models. Estimation and inference concerning the long-run properties of the model are then carried out using standard asymptotic normal theory. The analysis becomes more complicated when the variables are difference–stationary, or integrated of order 1. The auto-regressive distributed lag model with p lags of dependent variable Y_t and q lags of additional regressor X_t , ADL (p, q) is defined as:

$$ADL(p,q) = Y_{t} = \beta_{0} + \beta_{1}Y_{t-1} + \dots + \beta_{p}Y_{t-p} + \delta_{1}X_{t-1} + \dots + \delta_{q}X_{t-q} + \mu_{t}$$
(6.1)

 $\beta(L)Y_t = \beta_0 + \delta\bigl(L\bigr)X_{t-1} \quad \text{with lag-polynomials defined by}$

$$\beta(\mathbf{L})\mathbf{Y}_{t} = 1 - \beta_{1}\mathbf{L} - \dots - \beta_{p}\mathbf{L}_{p}$$

$$\delta(\mathbf{L}) = \delta_{1} + \delta_{2}\mathbf{L} + \dots + \delta_{q}\mathbf{L}_{q-1}$$
(6.2)

k additional predictors = ADL (p, q_1, \dots, q_k)

$$\beta(L)Y_{t} = \beta_{0} + \delta_{1}(L)X_{1,t-1} + \delta_{2}(L)X_{2,t-1} + \dots + \delta_{k}(L)X_{k,t-1}$$

Model Assumptions

$$E\left[u_{t}/Y_{t-1,t-2},\ldots,X_{1t-1},X_{1t-2},\ldots,X_{kt-1},X_{kt-2},\ldots\right] = 0$$
(6.4)

- (a). $(\mathbf{Y}_{t}, \mathbf{X}_{1t}, \dots, \mathbf{X}_{kt})$ are (strictly) stationary
 - (b). $\left(\mathbf{Y}_{t}, \mathbf{X}_{1t}, \dots, \mathbf{X}_{kt}\right)$ are ergodic (6.5)

 $(Y_{t,}X_{lt,}...,X_{kt})$ and $(Y_{t-j,}X_{lt-j},...,X_{kt-j})$ becomes independent for $j \rightarrow \infty$

 Y_t and X_{1t_1}, X_{kt} have nonzero, finite fourth moments

no perfect multicollinearity

E. Non Stationary Time Series Regression Models

A time series X_t is said to be stationary if its expected value and population variance are independent of time and if the population covariance between its values at time t and t+s depends on s but not on time.

An example of a stationary time series is an AR (1) process

$$X_{t} = \beta_2 X_{t-1} + \epsilon_{t} \tag{7.1}$$

Here $-1 < \beta_2 < 1$ and \in_t is white noise

If equation (7.1) is valid for time period t, it is also valid for time period t-1 i.e. the series is stationary.

$$X_{t-1} = \beta_2 X_{t-2} + \in_{t-1}$$

Substituting for X_{t-1} in equation (7.1), we get

$$\mathbf{X}_{t} = \beta_{2}^{2} \mathbf{X}_{t-2} + \beta_{2} \in_{t-1} + \in \mathbf{Y}_{t-1}$$

Continuing this process of lagging and substituting, we get



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor :6.887 Volume 5 Issue XII December 2017- Available at www.ijraset.com

$$X_{t} = \beta_{2}^{t} X_{0} + \beta_{2}^{t-1} \in_{1} + \dots + \beta_{2} \in_{t-1} + \in_{t}$$
(7.2)

In the previous examples, if $\beta_2\,$ is equal to 1, the original series becomes

$$\mathbf{X}_{t} = \mathbf{X}_{t-1} + \boldsymbol{\epsilon}_{t} \tag{7.3}$$

This is an example of a nonstationary process which is known as a random walk If it starts at X_0 at time 0, its value at time t is given by

$$\mathbf{X}_{t} = \mathbf{X}_{0} + \boldsymbol{\epsilon}_{t} + \dots \boldsymbol{\epsilon}_{t} \tag{7.4}$$

The key difference between this process and the corresponding A.R(1) process is that the contribution of each innovation is permanently built into the time series. In the more general version of the autogressive process with the constant β_1 , the process becomes a random walk with drift, if β_2 equals 1.

$$X_{t} = \beta_{1} + X_{t-1} + \epsilon_{t}$$
(7.5)
If the series starts at X₀ at time 0, X_t is given by

 $X_t = X_0 + \beta_{1t} + \epsilon_1 + \dots + \epsilon_t$

The expectation of X_t at time '0', $(X_0 + \beta_{1t})$ is a function of 't'. Another common example of a nonstationary time series is one possessing a time trend:

$$\mathbf{X}_{t} = \boldsymbol{\beta}_{1} + \boldsymbol{\beta}_{2} \mathbf{t} + \boldsymbol{\epsilon}_{t} \tag{7.6}$$

This type of trend is sometimes described as a deterministic trend. The expected value of X_t at time 't' $(\beta_1 + \beta_2 t)$ is not independent of t and so X_t is nonstationary.

F. Difference-Stationarity And Trend–Stationarity

If a nonstationary process can be transformed into a stationary one by differencing, it is said to be difference – stationary. A random walk, with or without drift, is an example. If X_t is a random walk with drift,

$$\Delta X_{t} = (X_{t} - X_{t-1}) = \beta_{1} + t_{1}$$
(7.7)

This is a stationary process with population mean β , and variance σ_t^2 , both independent of time. If a nonstationary of time series

can be transformed into a stationary process by differencing once, it is described as integrated of order 1, or I(1). If a time series can be made stationary by differencing twice, it is known as I(2), and so an. To complete the picture, a stationary process, which by definition need no differencing is described as I(0). In practice most series as I(0), I(1) or occasionally, I(2). The stochastic component in (7.1) is white noise. More generally, the stationary process reached after differencing may be ARMA (p, q) in which case the original series is characterized as an ARIMA (p, d, q) time series, where d is the number of times it has to be differenced to render it stationary.

A nonstationary time series is described as being trend-stationary if it can be transformed into stationary process by extracting a time trend. For example, the very simple model given by equation (7.1) can be attended by fitting the equation.

$$\mathbf{X}_{t} = \mathbf{b}_{1} + \mathbf{b}_{2}\mathbf{t}$$

An defining a new variable

$$\widehat{\mathbf{X}}_{t} = \mathbf{X}_{t} - \widehat{\mathbf{X}}_{t} = \mathbf{X}_{t} - \mathbf{b}_{1} - \mathbf{b}_{2}\mathbf{t}$$

The new, detrended, variable is of course just the residuals from the regression of x on t.

II. CONCLUSIONS

Time series analysis has advance from univariate modeling based on a single variable to multivariate models that employ the interrelationships between several such variables. Constructing such models requires the performing of tests to determine and to discover the interactions that exist between a given time series variables and one or more other variables. The given variable can be



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor :6.887 Volume 5 Issue XII December 2017- Available at www.ijraset.com

influenced not only by certain exogenous events occurring at particular points in time but also by contemporaneous, lagged and leading values of another variable or additional variables.

REFERENCES

- [1] Amemiya, T. (1980), "Selection of Regressors", Internatinal Economic Review, 21, 331-354.
- [2] Antoniadis, A., Gijbels, I., and Gregoire, G. (1997), "Model Selection Using Wavelet Decomposition and Applications", Biometrika, 84, 4, 751-763.
- [3] Aoki, M., and Havenner, A. (1991), "State-Space Modelling of Multiple Time Series", Econometric Reviews, 10, 1-59.
- [4] Atikinson, A.C. (1970), "A Method for Discriminating Between Models", Journal of the Royal Statistical Society B, 32, 323-353.
- [5] Box, G.E.P., and Jenkins, G.M. (1976), "Time Series Analysis, Forecasting and Control", Rev. ed. San Francisco: Holden-Day.
- [6] Davidson, R., and MacKinnon, J.G. (1993), "Estimation and Inference in Econometrics", New York : Oxford University Press.
- [7] Eubank, R.L., and Hart, J.D. (1992), "Testing Goodness-of-Fit in Regression Via Order Selection Criteria", Ann. Statist, 20, 1412-1425.
- [8] Fan, J. (1993), "Local Linear Regression smoothers and Their Minimax Efficiencies", Annals of Statistics", 21, 196-216.
- [9] Guttman, I., Pereyra, V., and Scolnik, H.D. (1973), "Least Squares Estimation for a Class of Non-Linear Model", Technometrics, 15, 2.
- [10] Hall, A. (1987), "The Information Matrix Test for the Linear Model", Review of Economic Studies, 54, 257-263.
- [11] Loh, W. (1985), "A New Method for Testing Separate Families of Hypotheses", Journal of the American Statistical Association, 80, 362-368.
- [12] MacKinnon, J.G. (1983), "Model Specification Tests Against Non-Nested Alternatives", Econometric Reviews, 2, 85-110.
- [13] Rao, C. R. (1973), "Linear Statistical Inference and its Applications", 2nd ed. (Wiley, New York).
- [14] White, H., and Olson, L. (1979), "Determinants of Wage Change on the Job: A Symmetric Test of Non-Nested Hypotheses", Mimeo. (University of Rochester, Rochester, NY).
- [15] Wu, P. (1993), "Variable Addition Test", Econometric Theory, Problem, 93.1.2, 9: 145-146.
- [16] Zellner, A. (1971), "An Introduction to Bayesian Influence in Econometrics", Wiley, New York.











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24*7 Support on Whatsapp)