



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 6 Issue: I Month of publication: January 2018

DOI: <http://doi.org/10.22214/ijraset.2018.1232>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

Isolation, Characterization and Activity of Amidase Producing *Paenibacillus Polymexa* from Semi Arid Soils

H.s.surti¹, V.C. Solank², A.T. Thakkar³, S.A.bhatt⁴

^{1, 2, 3, 4} Dept. of Life Sciences, HNGU, Patan

Abstract: Soil samples from different locations were collected and the amidase producing organisms were isolated by enrichment culture method in soil extract medium using acetamide as sole source of C and N and incubating at 37°C. A total of 29 bacterial isolates were obtained. Eight strains showed amidase production. The highest amidase producing strain (HSS22) was revealed as *Paenibacillus polymexa*. Acetamide containing mineral salt media when supplemented with Sodium citrate for *P. polymexa* resulted in higher growth of bacteria but addition of NaCl did not show remarkable effect on bacterial growth, this indicated that amides could be efficiently utilized as N source. Isolate *P. polymexa* was found to show highest amidase activity (19.34 U/ml) among the nine amidase producing isolates. The pH tolerance, temperature and amidase production capabilities of *P. Polymexa* were compared. It was found that *P. polymexa* was able to grow at temperature of 37°C, the pH 8.0, whereas another six isolates showed amidase activity in the pH range of (6.0-9.0) and temperature (30°C) thus *P. polymexa* was selected for subsequent studies. The results for enzyme activity under the performance of different physiochemical conditions and purification as well as molecular weights are recorded and discussed.

Keywords: Amidase, Soil Samples, Bacteria, Physiochemical condition, Purification.

I. INTRODUCTION

An amidase (EC 3.5.1.4), acylamidases, acylase (misleading), amidohydrolase (ambiguous), deaminase (ambiguous), fatty acylamidases, N-acetylaminohydrolase (ambiguous) are enzymes that catalyze the hydrolysis of an amide.

Amidase (Richards & Rolinson, 1961) act on the peptide linkage of the prosthetic group. Amidase belongs to the Family of hydrolases, those acting on carbon-nitrogen bonds other than peptide bonds, specifically in linear amides. These enzymes are involved in nitrogen metabolism in cells and are widely distributed in nature. They have been found in prokaryotic and eukaryotic cells. (Kotlova *et al.*, 1998) This enzyme utilizes two substrates, which are monocarboxylic acid amide and H₂O and result in the formation of monocarboxylate and NH₃.

In enzymology, an amidase (EC 3.5.1.4), acylamidase, acylase (misleading), amidohydrolase (ambiguous), deaminase (ambiguous), fatty acylamidase, N-acetylaminohydrolase (ambiguous) are enzyme that catalyze the hydrolysis of an amide.

A. Potential Applications Of Amidase

- 1) Food industry
- 2) Waste treatment
- 3) Production of acrylic acid
- 4) Biosynthesis of hydroxamic acids
- 5) Production of nicotinic acid
- 6) Peptide synthesis
- 7) Production of amino acids and their derivatives

II. MATERIALS AND METHOD

A. Collection Of Soil Sample

The soil samples for the isolation of amidase producing bacteria were collected from different fields viz., cultivated soil – Farm of Unjha district, Non- Cultivated soil – garden of Gandhinagar district, Arid soil – Santalpur and High saline soil – white desert, Kutch district. The pH of all sites was between 6.0 and 7.5. Samples were brought to the laboratory and were kept at 4°C in refrigerator till further processing.

B. Physico-Chemical Test Of Soil

- 1) Enrichment And Isolation
- 2) General media –nutrient broth/agar
- 3) Selective media- soil extract broth/agar

C. Characterization Of Isolates

- 1) **Cultural Characters:** - since isolate hss-22 exited highest amidase activity among all the isolates, therefore, this bacterium was selected for further characterization.
- 2) **Colony Morphology:** - bacterial strains were identified by gram's staining and various morphological characteristics like margine, pigment etc. all the results were observed and recorded according to the methods given by (goss, 1972) and (case and johanson, 1984).
- 3) **Biochemical Characters:-** various biochemical tests like tsi test, urea hydrolysis test, indole test, emb test, m.r. test, v.p. test, catalase test, starch utilization test, citrate test, h₂s test, gelatine liquidification test, lipid hydrolysis test, nitrate reduction test, dehydrogenase test, carbohydrate fermentation, were performed to identify the bacterial isolate hss-22.
- 4) **16s Rdna Sequence Determination:-**the isolates were identified by morphological and biochemical tests and 16s rna sequence analysis. 16s rna gene sequencing of the isolated microorganisms were carried out at chromous biotech pvt. ltd., bangalore, india

D. Screening of Enzyme

- 1) **Qualitative Assay:** A qualitative estimation of amidase producing organism was done using a modified method. (Rahim *et al.*, 2003)
- 2) **Quantitative assay:**
 - a) **Effect Of Different Growth Factors On Amidase Enzyme:-**
 - i) EFFECT OF TEMPERATURE
 - ii) EFFECT OF PH
 - iii) EFFECT OF CARBON SOURCE
 - iv) EFFECT OF NITROGEN SOURCES
 - v) EFFECT OF INCUBATION TIME
 - b) **Determination Of Pgp Characters:-**
 - i) INDOLE ACETIC ACID ANALYSIS
 - ii) PHOSPHATE SOLUBILISATION
 - iii) PRODUCTION OF AMMONIA:-
 - iv) SIDEROPHORE PRODUCTION:
 - v) HYDROGEN CYANIDE PRODUCTION
 - c) **Purification Of Amidase:** Purification of amidase enzyme performed by SDS-PAGE.

III. RESULTS AND DISCUSSION

Table:- 1 - Physico-chemical analysis of soil samples

Chemical analysis							
Soil sample	pH	EC	C (%)	P (mg/L)	K (mg/L)	Ca (mg/L)	Mg (mg/L)
Arid soil	6.8	0.5	0.12	0.10	70	4.5	4.4
Non- Cultivated soil	7.9	0.3	0.18	0.02	110	5.4	4.6
Cultivated soil	6.9	0.4	0.16	0.03	68	4.6	2.4
High saline soil	7.3	0.6	0.25	0.04	78	6.0	1.2
Physical analysis							
Soil texture (%)							WHC (%)
Soil sample	Gravel	Coarse sand	Fine sand	Silt	Clay		
Arid soil	1.0	1.0	1.10	3.30	93.59		28.35

Non- Cultivated soil	0.09	0.06	0.65	7.54	91.66	24.54
Cultivated soil	0.74	1.08	9.46	58.0	30.68	31.68
High saline soil	22.18	4.77	7.27	35.25	28.53	26.71

A. Screening

- 1) A qualitative analysis of amidase producing organism was done using a modified method by Rahim *et. al.*, 2003.
- 2) The pH of the medium was adjusted to 6.6 ± 0.2 .
- 3) The broth and plates were regularly examined for colour change from yellow to pink.
- 4) Among twenty nine isolates, eight isolates were give positive result in qualitative screening of Amidase.

B. Qualitative Screening




Fig: 1 Qualitative Screening of HSS22

Table:- 2 – Quantitative assay

Source of Soil	Bacterial isolates	Enzyme activity (U/ml)	Source of Soil	Bacterial isolates	Enzyme activity (U/ml)	Bacterial isolates	Enzyme activity (U/ml)	Bacterial isolates	Enzyme activity (U/ml)	Bacterial isolates	Enzyme activity (U/ml)
Arid Soil	HSS 1	---	Non Cultivated Soil	HSS 8	11.64	HSS 11	---	HSS 14	11.04	HSS 14	---
	HSS 2	12.21		HSS 9	---	HSS 12	---	HSS 15	---	HSS 15	---
	HSS 3	13.08		HSS 10	---	HSS 13	---	HSS 16	---	HSS 16	---
Saline Soil	HSS 4	---	Cultivated Soil	HSS 17	---	HSS 21	---	HSS 25	---	HSS 29	---
	HSS 5	---		HSS 18	16.92	HSS 22	19.32	HSS 26	---		
	HSS 6	---		HSS 19	---	HSS 23	---	HSS 27	---		
	HSS 7	---		HSS 20	12.6	HSS 24	12.9	HSS 28	---		

C. Characterization & Identification:

Table:- 3 – Morphological Characteristics

Organism	HSS 22
Size	Small
Shape	Irregular
Margin	Entire
Elevation	Mucoid
Texture	Smooth
Consistency	dry
Opacity	Translucent
Pigment	White shiny
Gram reaction	Gram positive


Table:- 4 - Biochemical Characterization

No.	Test		HSS 22
1	TSI Test		-
2	Urea Hydrolysis Test		-
3	Indole Test		-
4	EMB Test		Blue colour
5	M.R. Test		+
6	V.P. Test		-
7	Catalase Test		+
8	Starch Utilization Test		-
9	Citrate Test		+
10	H ₂ S Test		-
11	Gelatine liquidification Test		+
12	Lipid hydrolysis Test		+
13	Nitrate Reduction Test		-
14	Dehydrogenase Test		+
15	Carbohydrate Fermentation	Glucose	-
16		Lactose	-
17		Xylose	-
18		Mannose	-
19		Maltose	-

Table:- 5 - Identification of isolate by 16S rRNA:


Sample Id	Sequence	Organism	Similarity	Accession No.
HSS 22	TTTGCCACTCCTGGAT.....CTATTGCCCTTGTAGGACT G	<i>Paenibacillus polymexa</i>	100%	LN774600.1

D. Physio-chemical Characteristics




The results are shown in fig.2 *P. polymexa* showed highest activity at 35°C beyond these temperatures isolate recorded decrease in activity. Isolate recorded highest unit activity of 20.76 U/ml 35 °C

E. Effect of pH




The results are shown in fig. 3. The highest activity of amidase from *P. polymexa* was at pH 8 and the activity was 20.52U/ml.

F. Effect Of Carbon Source




The results are shown in fig. 4. It is clear from that starch is better carbon source for *P. polymexa* and the activity was 20.16.

G. Effect of nitrogen source



The results are shown in fig. 5. The activity of amidase with different nitrogen source used for growth of organisms. The highest activity of amidase by *P. polymexa* showed high enzyme activity in peptone was used as nitrogen source and the activity was 23.24 U/ml.

H. Effect Of Incubation Time



The results are shown in fig. 6. The optimize activity of amidase on different incubation time. The highest activity of amidase producing organism *P. polymexa* was at 48h and the activity was 14.72 U/ml

I. Purification of enzyme




Fig. 7 result of purification

Amidase enzyme from *Paenibacillus spp.* was purified by SDS PAGE method.

The enzyme was corresponding with std. (range 10 to 110 kDa)

The molecular weight of amidase from *Paenibacillus polymexa* was found 95 kDa.

IV. CONCLUSION

From this study was focused on amidase enzyme producing bacteria and its activity. The objectives set forth for the study were achieved and out some of the work is summarized below:

- A. In this work total 29 bacterial strains were isolated from various types of soil collected from different locations of North Gujarat region. On the basis of qualitative as well as quantitative screening, one bacterial isolated was selected as potent amidase producer.
- B. HSS 22 showed maximum enzyme activity 19.32 U/ml compared to other isolates.
- C. HSS 22 was identified as *Paenibacillus polymexa* on the basis of morphological characteristics, biochemical tests and 16 S rRNA sequencing.
- D. Effect of various physico chemical parameters was checked on the amidase activity of *P. polymexa* which showed maximum amidase activity at temperature 35°C, pH - 8, starch as a sole source of carbon, peptone as a nitrogen source and 48 hrs incubation time.
- E. The enzyme from *P. polymexa* by SDS PAGE method and the molecular weight were recorded as 95 kDa for *P. polymexa*.

REFERENCES

- [1] Asano, Y., Tachibana, M., Tani, Y., and Yamada, H. (1982) aeruginosa FEBS Lett Biochem Pseudomonas Purification and characterization of amidase which participates in nitrile degradation. *Agr* 46: 1175–1181.
- [2] Bork, P., and Koonin, E. V. (1994) A new family of carbon– nitrogen hydrolases. *Protein Sci* 3: 1344–1346.
- [3] Botrè F, Botrè C, Lorenti G, Mazzei F, Porcelli F, et al. (1993) Determination of L-glutamate and L-glutamine in pharmaceutical formulations by amperometric L-glutamate oxidase based enzyme sensors. See comment in PubMed Commons below *J Pharm Biomed Anal* 11: 679–686.
- [4] *Environ Microbiol* 61:998–1003 Google
- [5] CLARKE, P. H. (1970). The aliphatic amidases of *Pseudomonas aeruginosa*. *Advances in Microbial Physiology* 4, 179–222.
- [6] DuChateau, P., Thelwell, R., & Butters, G. (2004). Analysis of an adjoint problem approach to the identification of an unknown diffusion coefficient. *Inverse Problems*, 20(2), 601.
- [7] Dunstan, D. W., Zimmet, P. Z., Welborn, T. A., De Courten, M. P., Cameron, A. J., Sicree, R. A., ... & Atkins, R. (2002). The rising prevalence of diabetes and impaired glucose tolerance. *Diabetes care*, 25(5), 829–834.
- [8] Fyfe, P. K., Rao, V. A., Zemla, A., Cameron, S., & Hunter, W. N. (2009). Specificity and Mechanism of *Acinetobacter baumannii* Nicotinamidase: Implications for Activation of the Front-Line Tuberculosis Drug Pyrazinamide. *Angewandte Chemie International Edition*, 48(48), 9176–9179.
- [9] Gabor EM, Janssen DB (2004) Increasing the synthetic performance of penicillin acylase PAS2 by structure-inspired semi-random mutagenesis. *Protein Eng Des Sel* 17:571–579
- [10] Kato, Y., Ooi, R., & Asano, Y. (1998). Isolation and characterization of a bacterium possessing a novel aldoxime-dehydration activity and nitrile-degrading enzymes. *Archives of microbiology*, 170(2), 85–90.
- [11] Kelly, M., & Kornberg, H. L. (1964). Purification and properties of acyltransferases from *Pseudomonas aeruginosa*. *Biochemical Journal*, 93(3), 557.
- [12] Mehta, P. K., Bhatia, S. K., Bhatia, R. K., & Bhalla, T. C. (2015). Thermostable amidase catalyzed production of isonicotinic acid from isonicotinamide. *Process Biochemistry*, 50(9), 1400–1404.
- [13] Mehta, P. K., Bhatia, S. K., Bhatia, R. K., & Bhalla, T. C. (2015). Thermostable amidase catalyzed production of isonicotinic acid from isonicotinamide. *Process Biochemistry*, 50(9), 1400–1404.
- [14] O'reilly, C., & Turner, P. D. (2003). The nitrilase family of CN hydrolysing enzymes—a comparative study. *Journal of applied microbiology*, 95(6), 1161–1174.
- [15] Rawlings, N. D., & Barrett, A. J. (2000). MEROPS: the peptidase database. *Nucleic acids research*, 28(1), 323–325.
- [16] Rolf, M. J., & Lim, H. C. (1985). Experimental adaptive on-line optimization of cellular productivity of a continuous bakers' yeast culture. *Biotechnology and bioengineering*, 27(8), 1236–1245.
- [17] Rutherford ST, Bassler BL (2012) Bacterial quorum sensing: its role in virulence and possibilities for its control. *Cold Spring Harb Perspect Med*. doi:10.1101/cshperspect.a012427
- [18] Sharma M, Sharma N, Bhalla TC (2009) Amidases: versatile enzymes in nature. *Rev Env Sci Biotech* 8:343–366.
- [19] Takashima, F., & Hibiya, T. (1995). An atlas of fish histology: normal and pathological features.
- [20] Torii, K. U., Mitsukawa, N., Oosumi, T., Matsuura, Y., Yokoyama, R., Whittier, R. F., & Komeda, Y. (1996). The Arabidopsis ERECTA gene encodes a putative receptor protein kinase with extracellular leucine-rich repeats. *The Plant Cell*, 8(4), 735–746.
- [21] Yasuhisa Asano, Mineo Tachibana, Yoshiki Tani & Hideaki Yamada (1982) Purification and Characterization of Amidase which Participates in Nitrile Degradation, *Agricultural and Biological Chemistry*, 46:5, 1175–1181, DOI: 10.1080/00021369.1982.10865239



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)