iJRASET
International Journal For Research in Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 6 **Issue:** I **Month of publication:** January 2018

DOI: http://doi.org/10.22214/ijraset.2018.1262

www.ijraset.com

Call: 08813907089 | E-mail ID: ijraset@gmail.com

International Journal for Research in Applied Science & Engineering Technology (IJRASET)
ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor :6.887
Volume 6 Issue I, January 2018- Available at www.ijraset.com

Fuzzy Ideals and Anti Fuzzy Ideals of Near-Ring

T. L. Dewangan[1], Prof. M. M. Singh[2]
[1]*Department of Mathematics, Govt. Polytechnic Ramanujganj, Chhattisgarh, India*
[2]*Department of Mathematics, Shri Shankaracharya Technical Campus, Bhilai, India*

Abstract: The aim of this paper is to extend the notion of a fuzzy subnear-ring, fuzzy ideals of a near ring, anti fuzzy ideals of near-ring and to give some properties of fuzzy ideals and anti fuzzy ideals of a near-ring.
Keywords: Near-ring, Near-subring, Ideals of near-ring, Fuzzy set, Fuzzy subring, Fuzzy ideals of near-ring,Anti fuzzy ideals

I. INTRODUCTION

The theory of fuzzy set was introduced by Zadeh[3], applying which Rosenfeld[6] in 1971 defined fuzzy subgroups. Salah Abou-Zaid[4]introduced the theory of a fuzzy subnear-ring and fuzzy ideals of a near-ring. Fuzzy ideals of a ring and a characterization of a regular ring studied by Lui[7]. The notion of fuzzy ideals of near rings with interval valued membership functions introduced by B. Davvaz[10] in 2001. In 2001, Kyung Ho Kim and Young BaeJun[11] in their paper entitled "Normal fuzzy R-subgroups in near-rings" introduced the concept of a normal fuzzy R-subgroup in near-rings and explored some related properties. In 2005, Syam Prasad Kuncham and Satyanarayana Bhavanari in their paper entitled "Fuzzy Prime ideal of a Gamma-near-ring" introduced fuzzy prime ideal in Γ-near-rings.The anti-fuzzy ideals of near-ring defined by F. A. Azam, A. A. Mamun and F. Nasrin. In this paper we study the concept of fuzzy ideals of a near ring and some difference properties of fuzzy ideals and anti-fuzzy ideals of a near-ring.

II. PRELIMINARIES

For the sake of continuity we recall some basic definitions.

A. *Definition 2.1*

A set *N* together with two binary operations + (called *addition*) and · (called *multiplication*) is called a (right) *near-ring* if:

1) *N* is a group (not necessarily abelian) under addition;
2) multiplication is associative (so *N* is a semi group under multiplication); and
3) multiplication distributes over addition on the *right*: for any $x, y, z \in N$ it holds that $(x + y).z = (x.z) + (y.z)$.This near-ring will be termed as right near-ring. If $z.(x + y) = z.x + z.y$ instead of condition (3), the set N satisfies, then we call N a left near-ring. Near-rings are generalised of a rings, addition needs not be commutative and (more important) only one distributive law is postulated.

B. *Examples 2.2*

(1)Z be the Set of positive and negative integers with 0. $(Z, +)$ is a group . Define '.' on Z by $a.b = a$ for all $a, b \in Z$. Clearly $(Z, +, .)$ is a near ring. (2) Let $Z_{12} = \{0,1,2,3, \dots ,11\}$. $(Z_{12},+)$ is a group under '+' modulo 12. Define '.' on Z_{12} by $a.b = a$ for all $a, b \in Z$. Clearly $(Z_{12}, +, .)$ is a near ring. (3) Let $M_{2\times2} = \{(a_{ij})/ Z : Z$ is treated as a near ring$\}$. $M_{2\times2}$ under the operation of '+' and matrix multiplication '.' is defind by the following:

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} . \begin{bmatrix} e & f \\ g & h \end{bmatrix} = \begin{bmatrix} a+b & a+b \\ c+d & c+d \end{bmatrix}$$

Because we use the multiplication in Z i.e. $a.b = a$. So

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} . \begin{bmatrix} e & f \\ g & h \end{bmatrix} = \begin{bmatrix} ae+bg & af+bh \\ ce+dg & cf+dh \end{bmatrix} = \begin{bmatrix} a+b & a+b \\ c+d & c+d \end{bmatrix}.$$

It is easily verified $M_{2\times2}$ is a near ring. We denote xy instead of $.y$. Note that $x.0 = 0$and $x(-y) = -xy$ but in general $0x \neq 0$ for some $x \in R$. An ideal I of a near-ring R is a subset of R such that
(1) $(I, +)$is a normal subgroup of $(R, +)$, (2) $RI \subseteq I$ (3) $(r + i)s - rs \in I$for any $i \in I$ and any $r, s \in R$.

©IJRASET (UGC Approved Journal): All Rights are Reserved

International Journal for Research in Applied Science & Engineering Technology (IJRASET)
ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor :6.887
Volume 6 Issue I, January 2018- Available at www.ijraset.com

III. FUZZY IDEALS OF NEAR-RINGS

A. *Definition 3.1*

Let R be a near-ring and μ be a fuzzy subset of R. We say a fuzzy subnear-ring of R if (1) $\mu(x-y) \geq \min\{\mu(x), \mu(y)\}$, (2) $\mu(xy) \geq \min\{\mu(x), \mu(y)\}$, for all $x, y \in R$.

B. *Definition3.2*

Let R be a near-ring and μ be a fuzzy subset of R. μ is called a fuzzy left ideal of R if μ is a fuzzy subnear-ring of R and satisfies: for all $x, y \in R$.

1) $\mu(x-y) \geq \min\{\mu(x), \mu(y)\}$,
2) $\mu(y+x-y) \geq \mu(x)$,
3) $\mu(xy) \geq \mu(y)$ or $\mu(xy) \geq \mu(x)$

C. *Definition3.3*

Let R be a near-ring and μ be a fuzzy subset of R. μ is called a fuzzy right ideal of R if μ is a fuzzy subnear-ring of R and satisfies: for all $x, y \in R$.(1) $\mu(x-y) \geq \min\{\mu(x), \mu(y)\}$, (2) $\mu(xy) \geq \min\{\mu(x), \mu(y)\}$, (3) $\mu(y+x-y) \geq \mu(x)$, (4) $\mu((x+i)y - xy) \geq \mu(i)$.

D. *Example 3.4*

Let $R := \{a, b, c, d\}$ be a set with two binary operations as follows:

+	a	b	c	d
a	a	b	c	d
b	b	a	d	c
c	c	d	b	a
d	d	c	a	b

.	a	b	c	d
a	a	a	a	a
b	a	a	a	a
c	a	a	a	a
d	a	a	b	b

The we can easily see that $(R, +)$ is a group and $(R, .)$ is an semigroup and satisfies left distributive law. Hance $(R, +, .)$ is a left near-ring. Define a fuzzy subset $\mu : R \to [0,1]$ by $\mu(c) = \mu(d) < \mu(b) < \mu(a)$. Then μ is a left fuzzy ideal of R.

E. *Example 3.5*

Let $R := \{a, b, c, d\}$ be a set with two binary operations as follows.

+	a	b	c	d
a	a	b	c	d
b	b	a	d	c
c	c	d	b	a
d	d	c	a	b

.	a	b	c	d
a	a	a	a	a
b	a	a	a	a
c	a	a	a	a
d	a	b	c	b

Then we can easily see that $(R, +, .)$ is a left near-ring. Define a fuzzy subset $\mu : R \to [0,1]$ by $\mu(c) = \mu(d) < \mu(b) < \mu(a)$. Then μ is a fuzzy left ideal of R, but not fuzzy right ideal of R, Since $\mu((c+d)d - cd) = \mu(d) < \mu(b)$.

F. *Proposition 3.6*

If a fuzzy subset μ of R satisfies the properties $\mu(x-y) \geq \min\{\mu(x), \mu(y)\}$ then

©IJRASET (UGC Approved Journal): All Rights are Reserved

1) $\mu(0_R) \geq \mu(x)$
2) $\mu(-x) = \mu(x)$, for all $x, y \in R$

Proof.(1) We have that for any $x \in R$

$$\mu(0_R) = \mu(x - x)$$
$$\geq \min\{\mu(x), \mu(x)\}$$
$$= \mu(x)$$

Hence $\mu(0_R) \geq \mu(x)$.

(2) By (1), we have that

$$\mu(-x) = \mu(0_R - x)$$
$$\geq \min\{\mu(0_R), \mu(x)\}$$
$$= \mu(x)$$

Hence $\mu(-x) = \mu(x)$. ■

G. Proposition 3.7

Let μ be a fuzzy ideal of R. If $\mu(x - y) = \mu(0_R)$ then $\mu(x) = \mu(y)$.

Proof. Assume that $\mu(x - y) = \mu(0_R)$ for all $x, y \in R$. Then

$$\mu(x) = \mu(x - y + y)$$
$$\geq \min\{\mu(x - y), \mu(y)\}$$
$$= \min\{\mu(0_R), \mu(x)\}$$
$$= \mu(y)$$

So, $\mu(x) \geq \mu(y)$ (1)

Also, $\mu(y) = \mu(y - x + x)$
$$\geq \min\{\mu(y - x), \mu(x)\}$$
$$= \min\{\mu(0_R), \mu(x)\}$$
$$= \mu(x)$$

So, $\mu(y) \geq \mu(x)$ (2)

From equation (1) and (2), Hence $\mu(x) = \mu(y)$. □

H. Proposition 3.8

If $\mu: R \to [0,1]$ is a fuzzy ideals of near-ring R with multiplicative identity 1_R. Then $\mu(0_R) \geq \mu(x) \geq \mu(1_R)\ \forall\ x \in R$.

Proof: We know that, $\mu(x) = \mu(-x)$

And now,

$$\mu(0_R) = \mu(x - x)$$

$= \mu(x + (-x))$

$\geq \min\{\mu(x), \mu(-x)\}$

$= \mu(x)$ (1)

Also $\mu(x) = \mu(x, 1_R)$

$\geq \min\{\mu(x), \mu(1_R)\}$

$\geq \mu(1_R)$ (2)

From equation (1) and (2),

$\mu(0_R) \geq \mu(x) \geq \mu(1_R) \; \forall \; x \in R.$ ■

IV. ANTI-FUZZY IDEALS OF NEAR-RING

A. Definition 4.1

Let *R* be a near-ring and μ be a fuzzy subset of *R*. μ is called an anti-fuzzy left ideal of R if μ is a fuzzy subnear-ring of R and satisfies: for all $x, y \in R$.

1) $\mu(x - y) \leq \max\{\mu(x), \mu(y)\}$,

2) $\mu(y + x - y) \leq \mu(x)$,

3) $\mu(xy) \leq \mu(y)$ or $\mu(xy) \leq \mu(x)$

B. Definition4.2

Let *R* be a near-ring and μ be a fuzzy subset of *R*. μ is called a anti fuzzy right ideal of R if μ is a fuzzy subnear-ring of R and satisfies: for all $x, y \in R$,

1) $\mu(x - y) \leq \max\{\mu(x), \mu(y)\}$,

2) $\mu(xy) \leq \max\{\mu(x), \mu(y)\}$

3) $\mu(y + x - y) \leq \mu(x)$

4) $\mu((x + i)y - xy) \leq \mu(i)$.

C. Proposition 4.3 For every anti fuzzy ideals μ of *R*,

1) $\mu(0_R) \leq \mu(x), \forall \; x \in R.$

2) $\mu(x) = \mu(-x), \forall \; x \in R.$

3) $\mu(x - y) = \mu(0_R) \Rightarrow \mu(x) = \mu(y), \forall \; x, y \in R.$

Proof.(1) $\mu(0_R) = \mu(x - x)$

$\leq \max\{\mu(x), \mu(x)\}$

$= \mu(x)$.

(2) $\mu(-x) = \mu(0_R - x)$

$\leq \max\{\mu(0_R), \mu(x)\}$

$= \mu(x).$

For all $x \in R$. Since x is arbitrary, we conclude that $\mu(-x) = \mu(x)$.

(3) Assume that $\mu(x-y) = \mu(0_R)$ for all $x, y \in R$. Then

$$\mu(x) = \mu(x - y + y)$$
$$\leq max\{\mu(x-y), \mu(y)\}$$
$$= \max\{\mu(0_R), \mu(x)\}$$
$$= \mu(y)$$

So, $\mu(x) \leq \mu(y)$ (1)

Also,

$$\mu(y) = \mu(y - x + x)$$
$$\leq max\{\mu(y-x), \mu(x)\}$$
$$= \max\{\mu(0_R), \mu(x)\}$$
$$= \mu(x)$$

So, $\mu(y) \leq \mu(x)$ (2)


From equation (1) and (2)

Hence $\mu(x) = \mu(y)$. ■

REFERENCES

[1] M. Akram, Anti fuzzy Lie ideals of Lie algebras, Quasigroups Related Systems 14 (2006)123-132
[2] Y. Bingxue, Fuzzy semi-ideal and generalized fuzzy quotient ring, Iran. J. Fuzzy Syst. 5 (2008) 87-92.
[3] L. A. Zadeh, Fuzzy sets, Information and Control 8 (1965) 338-353
[4] S. A. Zaid, On fuzzy ideals and fuzzy quotient rings of a ring, Fuzzy Sets and Systems 59(1993) 205-210
[5] M. Zhou, D. Xiang and J. Zhan, Onanti fuzzy ideals of ¡-rings, Ann. Fuzzy Math. Inform.
[6] A. Rosenfeld, Fuzzy groups, J. Math. Anal. Appl. 35 (1971) 512-517.
[7] W. Liu, Fuzzy invariant subgroups and fuzzy ideals, Fuzzy Sets and Systems 8 (1982) 133-139.
[8] T. K. Dutta and B. K. Biswas, Fuzzy ideals of near-rings, Bull. Calcutta Math. Soc.,89(1997), 447–456.
[9] S.M. Hong, Y.B. Jun, H.S. Kim, Fuzzy ideals in near-rings, Bulletin of KoreanMathematical Society, 35(3), (1998), 455–464
[10] B.Davvaz, Fuzzy ideals of near-rings with interval valued membership functions,Journal of Sciences, Islamic republic of Iran, 12(2001), no. 2, 171–175.
[11] K. H. Kim andY.B. Jun, Anti fuzzy ideals in near rings, Iranian Journal of FuzzySystems, Vol. 2 (2005), 71–80.

©IJRASET (UGC Approved Journal): All Rights are Reserved



10.22214/IJRASET

45.98

IMPACT FACTOR: 7.129

IMPACT FACTOR: 7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)