iJRASET
International Journal For Research in Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 6 **Issue:** I **Month of publication:** January 2018

DOI: http://doi.org/10.22214/ijraset.2018.1364

www.ijraset.com

Call: 08813907089 | E-mail ID: ijraset@gmail.com

International Journal for Research in Applied Science & Engineering Technology (IJRASET)
ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor :6.887
Volume 6 Issue I, January 2018- Available at www.ijraset.com

On Ternary Quadratic Diophantine Equation

$$15x^2+15y^2+24xy=438z^2$$

G.Janaki[1] R. Radha[2]
[1, 2,] *Department of Mathematics, Cauvery college for women, Trichy*

Abstract: ***The ternary quadratic Diophantine equation*** $15x^2+15y^2+24xy=438z^2$ ***is analyzed for its non-trivial distinct integral solutions. Six different patterns of integral solutions are obtained. A few interesting relations among the solutions and special polygonal numbers are presented.***

Keywords: ***Ternary quadratic equation, Integral solutions.***

Notations:

$T_{3,n}=\frac{n(n+1)}{2}$ = Triangular number of rank n.

$T_{7,n}=\frac{n(5n-3)}{2}$ = Heptagonal number of rank n.

$T_{10,n}=n(4n-3)$ = Decagonal number of rank n.

$T_{12,n}=n(5n-4)$ = Dodecagonal number of rank n.

$T_{13,n}=\frac{n(11n-9)}{2}$ = Tridecagonal number of rank n.

$T_{17,n}=\frac{n(15n-13)}{2}$ = Heptadecagonal number of rank n.

$T_{18,n}=n(8n-7)$ = Octadecagonal number of rank n.

$Gno_n=(2n-1)$ = Gnomonic number of rank n.

Mathematical Classification: 11D09.

I. INTRODUCTION

Ternary quadratic equations are rich in variety. For more detailed understanding one can see [1-7]. For the non-trivial integral solutions of ternary quadratic Diophantine equations [8-9] has been studied. [10-13] has been referred for various ternary quadratic Diophantine equations. In this communication, we consider yet another interesting ternary quadratic equation $15x^2+15y^2+24xy=438z^2$ and obtain different patterns of non-trivial integral solutions. Also, a few interesting relations among the solutions and special polygonal, Centered, Gnomonic, Star numbers are presented.

II. METHOD OF ANALYSIS

The Quadratic Diophantine equation with four unknowns under consideration is

$$15x^2+15y^2+24xy=438z^2 \quad (1)$$

The substitution of the linear transformations

$$x=u+v \quad and \quad y=u-v \quad (2)$$

in (1) leads to

$$9u^2+v^2=73z^2 \quad (3)$$

©IJRASET (UGC Approved Journal): All Rights are Reserved

Four different choices of solutions to (3) are presented below. Once the values of u and v are known, using (2), the corresponding values of X and Y are obtained.

A. Pattern 1

In (3), $9u^2 + v^2 = 73z^2$

Assume that

$$z = 9a^2 + b^2, \quad a, b \neq 0 \qquad (4)$$

We can write $73 = (3+8i)(3-8i)$ (5)

Substituting (4) and (5) in (3), we get

$$(3u + iv)(3u - iv) = (3+8i)(3-8i)(3a + ib)^2 (3a - ib)^2$$

Equating the positive and negative factors, we get

$$(3u + iv) = (3+8i)(3a + ib)^2 \qquad (6)$$

$$(3u - iv) = (3-8i)(3a - ib)^2 \qquad (7)$$

Equating the real and imaginary parts in either (6) or (7), we get

$$u = 9a^2 - b^2 - 16ab \qquad (8)$$

$$v = 72a^2 - 8b^2 + 18ab \qquad (9)$$

Substituting (8) and (9) in (2), we get

$$\left.\begin{aligned} x = x(a,b) &= 81a^2 - 9b^2 + 2ab \\ y = y(a,b) &= -63a^2 + 7b^2 - 34ab \\ z = z(a,b) &= 9a^2 + b^2 \end{aligned}\right\} \qquad (10)$$

Thus (10) represent non-zero distinct integer solution to (1) in two parameters.

B. Observations

1) $x(a,a) - 74T_{4,a} = 0$

2) $y(a,a) + 90T_{4,a} = 0$

3) $x(2a,a) + y(2a,a) = 6a^2$, a nasty number.

4) $z(a(a+1), b(b+1)) - 9P_a^2 - P_b^2 = 0$

5) $x(a,a+1) + y(a,a+1) + z(1,5a) + 29 = (3a-6)^2$, a perfect square.

6) $x(a,a) + y(a,a) + z(a,a) = -6a^2$, a nasty number.

C. Pattern 2

In (3), $9u^2 + v^2 = 73z^2$

Consider the linear transformations

$$\left.\begin{aligned} z &= X + 9T \\ u &= X + 73T \end{aligned}\right\} \qquad (11)$$

or

$$\left.\begin{aligned} z &= X - 9T \\ u &= X - 73T \end{aligned}\right\} \qquad (12)$$

©IJRASET (UGC Approved Journal): All Rights are Reserved

Substituting (11) or (12) in (3), we get

$$v^2 = 64(X^2 - 657T^2) \quad (13)$$

Write $v = 8V$ (14)

Substituting (14) in (13), we get

$$V^2 = X^2 - 657T^2$$

$$X^2 = 657T^2 + V^2 \quad (15)$$

This is in the standard form $x^2 = Dy^2 + z^2$

The corresponding solutions to (15) are

$$\left.\begin{aligned} T &= 2ab \\ V &= 657a^2 - b^2 \\ X &= 657a^2 + b^2 \end{aligned}\right\} \quad (16)$$

Substituting (16) in (11) and (14), we get

$$\left.\begin{aligned} z &= 657a^2 + b^2 + 18ab \\ u &= 657a^2 + b^2 + 146ab \\ v &= 5256a^2 - 8b^2 \end{aligned}\right\} \quad (17)$$

Substituting (17) in (2), we get

$$\left.\begin{aligned} x &= x(a,b) = 5913a^2 - 7b^2 + 146ab \\ y &= y(a,b) = -4599a^2 + 9b^2 + 146ab \\ z &= z(a,b) = 657a^2 + b^2 + 18ab \end{aligned}\right\} \quad (18)$$

Thus (18) represent non-zero distinct integer solution to (1) in two parameters.

D. Observations

1) $x(a,a) - 6052T_{4,a} = 0$

2) $y(a,a) + 4444T_{4,a} = 0$

3) $z(a,a)$ is a perfect square.

4) $x(a,a+1) - y(a,a+1) \equiv 16 \pmod{32}$

5) $x(a,a+1) - y(a,a+1) - T_{68,a} \equiv -16 \pmod{10463}$

6) $y(1,B+3) = 100Gno_B + 9T_{4,B} - 3980$

E. Pattern 3

The Ternary quadratic equation (3) can be written as

$$9u^2 - 9z^2 = 64z^2 - v^2 \quad (19)$$

Factorizing (19) we have

$$(3u + 3z)(3u - 3z) = (8z + v)(8z - v)$$

$$\frac{3u + 3z}{8z + v} = \frac{8z - v}{3u - 3z} = \frac{A}{B}, B \neq 0 \quad (20)$$

This is equivalent to the following two equations.

©IJRASET (UGC Approved Journal): All Rights are Reserved

$$Av - 3Bu + z(8A - 3B) = 0 \quad (21)$$

$$Bv - 3uA - z(3A + 8B) = 0 \quad (22)$$

Applying the method of cross multiplication, we get

$$z = z(A,B) = 3A^2 + 3B^2 \quad (23)$$

$$\left.\begin{aligned} u = u(A,B) &= 3A^2 - 3B^2 + 16AB \\ v = v(A,B) &= -24A^2 + 24B^2 + 18AB \end{aligned}\right\} \quad (24)$$

Substituting (22) in (2), we get

$$\left.\begin{aligned} x = x(A,B) &= -21A^2 + 21B^2 + 34AB \\ y = y(A,B) &= 27A^2 - 27B^2 - 2AB \end{aligned}\right\} \quad (25)$$

Thus (23) and (25) represent non-zero distinct integer solution to (1) in two parameters.

F. Observations

1) $x(A,A) - y(A,A) = (6A)^2$, *a* perfect square.

2) $z(A,A)$ is a nasty number.

3) $x(A,A+1) + y(A,A+1) + T_{44,A} \equiv -6 (\mathrm{mod}\, 53)$

4) $z(A, A(A+1)) = 3T_{4,A}^2 + 12P_A^5$

5) $x(A,1) + Gno_{13n} + 8 = y(A,1) + star_A$

6) $x(A, A(4A-3)) + y(A, A(4A-3)) + 96T_{4,A}^2 = T_{4,A}\left[Gno_{136A} - 143\right]$

G. Pattern 4

(15) can be written as

$$X^2 - V^2 = 657T^2$$

$$(X+V)(X-V) = (657T)T$$

Equating the positive and negative factors we get

$$X + V = 657T \quad (26)$$

$$X - V = T \quad (27)$$

Solving (26) and (27), we get

$$\left.\begin{aligned} X &= 329T \\ V &= 328T \end{aligned}\right\} \quad (28)$$

Substitute (28) in (14) we get

$$v = 2624T$$

$$For\ T = A,\ v = 2624A$$

Substitute the value of v in (11), we get

$$u = 402A$$

Substituting u and v in (2)

©IJRASET (UGC Approved Journal): All Rights are Reserved

$$\left.\begin{aligned} x &= 3026A \\ y &= -2222A \\ z &= 338A \end{aligned}\right\} \qquad (29)$$

Thus (29) represent non-zero distinct integer solution to (1) in one parameter.

H. Observations

1) $z(A^2) + z(A) - 338\,\mathrm{Pr}_A = 0$

2) $2x(A^2) + 2y(A^2) - 4z(A^2) = (16A)^2$, is a perfect square.

3) $y(A^3) + y(A^2) + 4444P_A^5 = 0$


4) $3x(A^2) + 3y(A^2) + z(A^2) - 59T_{4,A} = (53A)^2$, is a perfect square.

5) $x(A^2) + x(A) - 6052P_A^2 = 0$

REFERENCE

[1] Dickson.L.E., "History of the Theory of Numbers", Chelsea Publishing Company, New York, Vol II, 1952.
[2] M.A.Gopalan, Manju Somanath and N. Vanitha, "one Quadratic Diophantine equation with four variables", Bulletin of pure applied Sciences, Vol.24E, No.2, pp.389-391 (2005).
[3] M.A. Gopalan, Manju Somanath and N. Vanitha, "Integral Solutions of $ax^2 + by^2 = w^2 - z^2$", Advances in Theoretical and Applied Mathematics, Vol.l, No.3.pp.223-226(2006).
[4] M.A. Gopalan and S. Vidyalakshmi, "Integral solutions of $x^2+y^2 = w^2+Dz^2$", Advances in Theoretical and Applied Mathematics, Vol.l, No.2,pp.115-118(2006).
[5] M.A. Gopalan and S. Vidyalakshmi, "Integral solutions of $kxy - w(x+y)=z^2$",Advances in Theoretical and Applied Mathematics, Vol.1No.2,pp.167-172(2006)
[6] Mordell.L.J., Diophantine equation, Academic Press, New York (1969)
[7] M.A. Gopalan and S. Vidyalakshmi, "On the Diophantine equation $kxy +yz+zx=w^2$" Pure and Applied Mathematika Sciences, Vol. LXVIII, No. 1-2, September 2008.
[8] M.A. Gopalan, Manju Somanath and N. Vanitha, "On a Quadratic Diophantine equation with four variables", Antartica J.math., 4(1), 41-45, (2007).
[9] M.A. Gopalan, Manju Somanath and N. Vanitha, "Homogeneous Quadratic equation with four unknowns", Acta ciencia Indica, Vol.XXXIIIM, No.3, 915(2007).
[10] M.A. Gopalan, Manju Somanath and N. Vanitha, "On Space Pythagorean equation $x^2+y^2+z^2 = w^2$", International journal of Mathematics, Computer Sciences and Information Technology, Vol.1,No.1, January-June, pp.129-1332008.
[11] M.A. Gopalan and S.Vidyalakshmi, "Quadratic Diophantine equation with four variables $x^2+y^2+xy+y=u^2+v^2uv+u-v$". Impact J.Sci. Tech: Vol 2(3), 125-127, 2008.
[12] M.A. Gopalan, Manju Somanath and N. Vanitha, " On Quadratic Diophantine equation with four unknowns $x^2+y^2=z^2+2w^2-2w+1$ Impact J.Sci. Tech: Vol 2(2), 97-103, 2008.
[13] G.Janaki and S. Vidyalakshmi, "Integral solutions of $xy+x+y+1 = z^2 - w^2$", Antartica J.math., 7(1), 31-37, (2010).

©IJRASET (UGC Approved Journal): All Rights are Reserved



10.22214/IJRASET

45.98

IMPACT FACTOR: 7.129

IMPACT FACTOR: 7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)