



# **iJRASET**

International Journal For Research in  
Applied Science and Engineering Technology



---

# **INTERNATIONAL JOURNAL FOR RESEARCH**

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume: 2      Issue: XII      Month of publication: December 2014**

**DOI:**

**[www.ijraset.com](http://www.ijraset.com)**

**Call:  08813907089**

**E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)**

# A Review on Nanoparticles in Mosquito Control - A Green Revolution in Future

Priya S<sup>1</sup>, Santhi S<sup>2</sup>

<sup>1</sup>Ph.D. Research Scholar, <sup>2</sup>Assistant Professor,

Department of Zoology, Queen Mary's College (Autonomous), Mylapore, Chennai-600 004, Tamil Nadu, India

**Abstract:** Mosquitoes are the potential vectors of many diseases, including malaria, filariasis, dengue, brain fever, etc. There is an urgent need to check the proliferation of the population of vector mosquitoes in order to reduce vector borne diseases by appropriate control methods. Mosquito control is of serious concern in developing countries like India due to the lack of general awareness, development of resistance, and socioeconomic reasons. Nanotechnology, a promising field of research opens up in the present decade and is expected to give major impulses to technical innovations in a variety of industrial sectors in the future. Over the past few decade, nanoparticles of noble metals such as silver exhibited significantly distinct physical, chemical and biological properties from their bulk counterparts. Nano-size particles of less than 100 nm in diameter are currently attracting increasing attention for the wide range of new applications in various fields of industry. Presently, there is a need for increasing the efforts to develop newer and effective methods to control mosquito vectors. The existing chemical and biological methods are not as effective as in earlier period owing to different technical and operational reasons. In particular, this present paper focused on potential role of nanoparticles in mosquito control.

**Keywords -** Larvicides, Plant extracts, Fungi, Bacteria, Silver nanoparticles

## I. INTRODUCTION

Vector control is an essential requirement in control of epidemic diseases such as malaria, filariasis, dengue that are transmitted by different species of mosquitoes. Emergence of insecticide resistance and their harmful effects on non-target organisms and environment has necessitated an urgent search for development of new and improved mosquito control methods that are economical and effective as well as safe for non-target organisms and the environment. Insecticides synthesized from natural products, such as silver, gold or silicon nanoparticles of herbal origin have become a priority in this search.

Nanoparticles are defined as particulate dispersions or solid particles with a size of 10-1000 nm. The word “nano” is derived from a Greek word meaning “dwarf”. In technical terms, the word “nano” means  $10^{-9}$ , or one billionth of a meter. Naturally, the word nanotechnology evolved due to use of nanometer size particles. Targeted nanoparticles exhibit many novel characteristic features, such as extra ordinary strength, more chemical reactivity, magnetic properties and or high electrical conductivity. “Nanotechnology” deals with application of such particles in biological, physical, chemical, environmental, agricultural, industrial or pharmaceutical science. Depending upon the method of preparation, nanoparticles, nanospheres or nanocapsules can be obtained. Although physical and chemical methods are more popular and widely used for synthesis of nanoparticles, the related environmental toxicity and non-biodegradable nature of the products limited their applications. So, the “green” route for synthesis of nanoparticles from herbal origin is of great interest due to eco-friendliness, economic prospects, feasibility and wide range of applications (Salam *et al.*, 2012).

## II. NANOPARTICLES IN MOSQUITO CONTROL

Applications of nanotechnology have been extended in the field of mosquito control by the synthesis of silver/gold nanoparticles from environmentally acceptable plant extract. Synthesized silver or gold nanoparticles help to produce new insecticides and insect repellants. The characterization and the structure determination of these nanoparticles have become possible through the application of modern scientific instruments such as UV-VIS spectroscopy, Fourier Transform Infrared Spectroscopy, X-ray Diffraction, Scanning and Transmission Electron Microscopy (Adhikari *et al.*, 2013). The use of “green” processes for the synthesis of nanoparticles is a new and rapidly developing branch of nanotechnology. However, knowledge of the bioactivity of nanoparticles against mosquitoes is very limited. Accordingly, this review presents green synthesis of nanoparticles and their potential in

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

mosquito control.

### III. NANO PARTICLES OF HERBAL ORIGIN: A RECENT ECO-FRIENDLY TREND IN MOSQUITO CONTROL

The silver nanoparticles (Ag NPs) which are less likely to cause ecological damage have been identified as potential replacement of synthetic chemical insecticides. Hence the need to use green synthesized Ag NPs for the control of mosquitoes causing many diseases. The results described below were based on plant mediated Ag NPs and have been tested against the larvae of mosquito (Table 1).

Marimuthu *et al.*, (2011) synthesized Ag NPs utilizing aqueous leaf extract of *Mimosa pudica* Gaertn (Mimosaceae) and its antiparasitic activities against the larvae of malaria vector, *An. subpictus* Grassi, filariasis vector *Culex quinquefasciatus* Say (Diptera: Culicidae), and *Rhipicephalus (Boophilus) microplus* Canestrini (Acari: Ixodidae). Santhoshkumar *et al.*, (2011) observed the maximum efficacy in crude methanol, aqueous and synthesized Ag NPs using leaf extract of *Nelumbo nucifera* against the larvae of *An. subpictus* (LC<sub>50</sub>=8.89, 11.82, and 0.69 ppm respectively) and against the larvae of *Cx. quinquefasciatus* (LC<sub>50</sub>=9.51, 13.65, and 1.10 ppm respectively).

Rajakumar *et al.*, (2011) studied the larvicidal activity of synthesized Ag NPs using aqueous extract from *Eclipta prostrata* and observed the maximum efficacy in crude aqueous and synthesized Ag NPs against fourth instar larvae of *Cx. quinquefasciatus* (LC<sub>50</sub>=27.49 and 4.56 mg/L; LC<sub>90</sub>=70.38 and 13.14 mg/L), and against *An. subpictus* (LC<sub>50</sub>=27.85 and 5.14 mg/L; LC<sub>90</sub>=71.45 and 25.68 mg/L). Gnanadesigan *et al.*, (2011) synthesized Ag NPs with *Rhizophora mucronata* leaf extract to control the larvae of *Aedes aegypti* and *Cx. quinquefasciatus*. The LC<sub>50</sub> and LC<sub>90</sub> values of the synthesized Ag NP were identified as 0.585 mg/L and 2.615 mg/L for *i* and 0.891 mg/L and 6.291 mg/L for *Cx. quinquefasciatus*.

Jayaseelan *et al.*, (2011) observed the maximum larvicidal activity in the synthesized Ag NPs by leaf aqueous extract of *Tinospora cordifolia* against fourth instar larvae of *An. subpictus* and *Cx. quinquefasciatus* (LC<sub>50</sub>=6.43 and 6.96 mg/L respectively). Arjunan *et al.*, (2012) synthesized the median lethal concentration (LC<sub>50</sub>) of stable Ag NPs using *A. squamosa* leaf broth that killed fourth instar larvae of *Ae. aegypti*, *Cx. quinquefasciatus* and *Anopheles stephensi* (LC<sub>50</sub> = 0.30, 0.41, and 2.12 ppm respectively).

Priyadarshini *et al.*, (2012) synthesized Ag NPs utilizing *Euphorbia hirta* leaf extract against malarial vector *An. stephensi* and found the highest larval mortality in the first to fourth instar larvae (LC<sub>50</sub> = 10.14, 16.82, 21.51, and 27.89 ppm, respectively and LC<sub>90</sub> = 31.98, 50.38, 60.09, and 69.94 ppm, respectively). The LC<sub>50</sub> and LC<sub>90</sub> values of pupae were 34.52 and 79.76 ppm. Sareen *et al.*, (2012) reported that the larvicidal efficacy of Ag NPs synthesized from aqueous leaf extract of *Hibiscus rosasinensis* to control the larvae of *Aedes albopictus*. Patil *et al.*, (2012a) synthesized Ag NPs using *P. daemia* plant latex against *Ae. aegypti* and *An. stephensi*.

Table 1: Green synthesis of silver nanoparticles by using aqueous plant extracts and their potential mosquito host.

| Plant Species               | Common name      | Vernacular name (Tamil) | Plant parts used | Test NPs | Test Mosquito Species                              | City / State | Source                             |
|-----------------------------|------------------|-------------------------|------------------|----------|--|--------------|------------------------------------|
| <i>Eclipta prostrata</i>    | False Daisy      | Karisalaankani          | Leaf             | Ag       | <i>An. subpictus</i> , <i>Cx. quinquefasciatus</i> | Vellore-TN   | Rajakumar <i>et al.</i> , 2011     |
| <i>Mimosa pudica</i>        | Touch Me Not     | Thotta siningi          | Leaf             | Ag       | <i>An. subpictus</i> , <i>Cx. quinquefasciatus</i> | Vellore-TN   | Marimuthu <i>et al.</i> , 2011     |
| <i>Nelumbo nucifera</i>     | Indian Lotus     | Thammarai               | Leaf             | Ag       | <i>An. subpictus</i> , <i>Cx. quinquefasciatus</i> | Vellore-TN   | Santhoshkumar <i>et al.</i> , 2011 |
| <i>Rhizophora mucronata</i> | Asiatic Mangrove | Kandal                  | Leaf             | Ag       | <i>Ae. aegypti</i> , <i>Cx. quinquefasciatus</i>   | Madurai-TN   | Gnanadesigan <i>et al.</i> , 2011  |
| <i>Tinospora cordifolia</i> | Guduchi          | Shindilakodi            | Leaf             | Ag       | <i>An. subpictus</i> , <i>Cx. quinquefasciatus</i> | Vellore-TN   | Jayaseelan <i>et al.</i> , 2011    |

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

|   |   |                                       |             |    |   |                     |                                       |
|---|---|---------------------------------------|-------------|----|---|---------------------|---------------------------------------|
| <i>Annona squamosa</i>  | Custard Apple                                       | Sitapalam                             | Leaf        | Ag | <i>Ae. aegypti</i> , <i>An. stephensi</i> , <i>Cx. quinquefasciatus</i> | Coimbatore-TN       | Arjunan <i>et al.</i> , 2012          |
| <i>Euphorbia hirta</i>  | Asthma Weed   | Amman Pachirisi                       | Leaf        | Ag | <i>An. stephensi</i>  | Coimbatore-TN       | Priyadarshini <i>et al.</i> , 2012    |
| <i>Hibiscus rosasinensis</i>  | China Rose  | Cembaruthi                            | Leaf        | Ag | <i>Ae. albopictus</i>   | Kochi-Kerala        | Sareen <i>et al.</i> , 2012           |
| <i>Nerium oleander</i>  | Oleander  | Arali                                 | Leaf        | Ag | <i>An. stephensi</i>  | Coimbatore-TN       | Roni <i>et al.</i> , 2012             |
| <i>Pedilanthus tithymaloides</i>  | Devil's Backbone                                    | Peru neranji                          | Stem        | Ag | <i>Ae. aegypti</i>  | Coimbatore-TN       | Sundaravadivelan <i>et al.</i> , 2012 |
| <i>Pergularia daemia</i>  | Trellis-Vine  | Veli paruthi                          | Plant latex | Ag | <i>Ae. aegypti</i> , <i>An. stephensi</i>                               | Jalgaon-Maharashtra | Patil <i>et al.</i> , 2012a           |
| <i>Plumeria rubra</i>   | Frangipani  | Sampangi                              | Plant latex | Ag | <i>Ae. aegypti</i> , <i>An. stephensi</i>                               | Jalgaon-Maharashtra | Patil <i>et al.</i> , 2012b           |
| <i>Vinca rosea</i>  | Peri Wimkle   | Nithya kalyani                        | Leaf        | Ag | <i>An. stephensi</i> , <i>Cx. quinquefasciatus</i>                      | Chidambaram - TN    | Subarani <i>et al.</i> , 2012         |
| <i>Ammannia baccifera</i>   | Blistering Ammannia                                 | Kal-luruvi                            | Aerial      | Ag | <i>An. subpictus</i> , <i>Cx. quinquefasciatus</i>                      | Chennai-TN          | Suman <i>et al.</i> , 2013            |
| <i>Anthocephalus cadamba</i> ,<br><i>Cymbopogon citratus</i>  | Kadam,<br>Lemon Grass                               | Vellai kadambu,<br>Karpapurpul        | Leaf        | Au | <i>Cx. quinquefasciatus</i>   | Coimbatore-TN       | Arjunan <i>et al.</i> , 2013          |
| <i>Cadaba indica</i>  | Dabi  | Manatukkurntu                         | Leaf        | Ag | <i>An. stephensi</i> , <i>Cx. quinquefasciatus</i>                      | Coimbatore-TN       | Kalimuthu <i>et al.</i> , 2013        |
| <i>Cocos nucifera</i>   | Coconut   | Thaengaai                             | Mesocarp    | Ag | <i>An. stephensi</i> , <i>Cx. quinquefasciatus</i>                      | Vellore-TN          | Roopan <i>et al.</i> , 2013           |
| <i>Drypetes roxburghii</i>  | Putranjva   | Irukolli                              | Fruit       | Ag | <i>An. stephensi</i> , <i>Cx. quinquefasciatus</i>                      | Burdwan-WB          | Haldar <i>et al.</i> , 2013           |
| <i>Jatropha gossypifolia</i> ,<br><i>Euphorbia tirucalli</i> ,<br><i>Pedilanthus tithymaloides</i> ,<br><i>Alstonia macrophylla</i> | Bellyache Bush, Milk Bush, Devil's Backbone, Batino | Kattamanakku, Tirucalli, Peru neranji | Leaf        | Ag | <i>Ae. aegypti</i> , <i>An. stephensi</i>                               | Jalgaon-Maharashtra | Borase <i>et al.</i> , 2013           |
| <i>Ficus racemosa</i>   | Indian Fig  | Aththi                                | Bark        | Ag | <i>Cx. quinquefasciatus</i> ,<br><i>Cx. Gelidus</i>                     | Vellore-TN          | Velayutham <i>et al.</i> , 2013       |
| <i>Sida acuta</i>   | Common Wireweed                                     | Palambasi                             | Leaf        | Ag | <i>Ae. aegypti</i> , <i>An. stephensi</i> , <i>Cx. quinquefasciatus</i> | Chidambaram - TN    | Veerakumar <i>et al.</i> , 2013       |

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

|                                 |                    |                 |   |    |   |                 |                                  |
|---------------------------------|--------------------|-----------------|---|----|---|-----------------|----------------------------------|
| <i>Solanum nigrum</i>           | Black Nightshade   | Manathakali     | Fresh leaves, dry leaves, green berries | Ag | <i>An. stephensi</i> , <i>Cx. quinquefasciatus</i>                      | Burdwan-WB      | Rawani <i>et al.</i> , 2013      |
| <i>Delphinium denudatum</i>     | Jadwar             | Nirbasi         | Root                                    | Ag | <i>Ae. aegypti</i>  | Kanchipuram -TN | Suresh <i>et al.</i> , 2014      |
| <i>Feronia elephantum</i>       | Wood apple         | Vila            | Leaf                                    | Ag | <i>Ae. aegypti</i> , <i>An. stephensi</i> , <i>Cx. quinquefasciatus</i> | Chidambaram -TN | Veerakumar <i>et al.</i> , 2014b |
| <i>Heliotropium indicum</i>     | Indian Heliotrope  | Thel kodukku    | Leaf                                    | Ag | <i>Ae. aegypti</i> , <i>An. stephensi</i> , <i>Cx. quinquefasciatus</i> | Chidambaram -TN | Veerakumar <i>et al.</i> , 2014a |
| <i>Leucas aspera</i>            | Common Leucas      | Thumbai         | Leaf                                    | Ag | <i>Ae. aegypti</i>  | Salem-TN        | Suganya <i>et al.</i> , 2014     |
| <i>Parthenium hysterophorus</i> | Congress Grass     |                 | Root                                    | Ag | <i>Cx. quinquefasciatus</i>   | Burdwan-WB      | Mondal <i>et al.</i> , 2014      |
| <i>Pongamia pinnata</i>         | Pongam             | Pungai          | Leaf                                    | Ag | <i>Ae. albopictus</i>   | Hyderabad-AP    | Naik <i>et al.</i> , 2014        |
| <i>Sterculia foetida</i>        | Wild Indian Almond | Kutiraippitukku | Seed                                    | Ag | <i>Ae. aegypti</i> , <i>An. stephensi</i> , <i>Cx. quinquefasciatus</i> | Hyderabad-AP    | Rajasekharreddy and Rani, 2014   |
| <i>Calotropis gigantean</i>     | Crown Flower       | Erukku          | Leaf                                    | Ag | <i>Ae. aegypti</i> , <i>An. stephensi</i>                               | Coimbatore-TN   | Priya <i>et al.</i> , 2014       |
| <i>Cleistanthus collinus</i>    | Garari             | Odaichi         | Leaf                                    | Ag | <i>Ae. aegypti</i>  | Musiri-TN       | Ramar <i>et al.</i> , 2014       |

Patil *et al.*, (2012b) synthesized Ag NPs from *Plumeria rubra* latex and observed the LC<sub>50</sub> values for second and fourth larval instars after 24 hr exposure were 1.49, 1.82 ppm against *Ae. aegypti* and 1.10, 1.74 ppm against *An. stephensi*, respectively. The LC<sub>50</sub> values of crude aqueous latex of *P. rubrum* were 181.67, 287.49 ppm and 143.69, 170.58 ppm against 2nd and 4th larval instars of *Ae. aegypti* and *An. stephensi*, respectively.

Roni *et al.*, (2012) determined the larvicidal activity of synthesized Ag NPs through leaf extract of *Nerium oleander* (Apocynaceae) against the first to fourth instar larvae and pupae of malarial vector, *An. stephensi* (Diptera: Culicidae). Subarani *et al.*, (2012) evaluated the larvicidal activities to determine the efficacies of synthesized Ag NPs using aqueous leaf extract of *V. rosea* against the larvae of malarial vector *An. stephensi* Liston and filariasis vector *Cx. quinquefasciatus* Say (Diptera: Culicidae). Sundaravadivelan *et al.*, (2012) tested the biolarvicidal effect of phyto-synthesized Ag NPs using *Pedilanthus tithymaloides* (L.) Poit stem extract against the dengue vector *Ae. aegypti*.

Suman *et al.*, (2013) synthesized Ag NP from aqueous aerial extract of *Ammannia baccifera* and found that it can effectively inhibit the activity of *An. subpictus* and *Cx. quinquefasciatus* larvae. Arjunan *et al.*, (2013) biosynthesized gold nanoparticles using *Cymbopogon citratus* and *Anthocephalus cadamba* and experimented on the larvicidal effect on the filarial vector, *Cx. quinquefasciatus* and observed that the lethal concentrations LC<sub>50</sub> and LC<sub>90</sub> values of 1.08 and 2.76 ppm for gold nanoparticles and 21.82 and 79.52 ppm for the third instar of *Cx. quinquefasciatus*.

Borase *et al.*, (2013) synthesized Ag NPs from aqueous leaves extracts of four plant species (*Jatropha gossypifolia*, *Euphorbia tirucalli*, *Pedilanthus tithymaloides* and *Alstonia macrophylla*) and evaluated their effects on II and IV instars larvae of *Ae. aegypti* and *An. stephensi*. Results revealed that the larvicidal activity of Ag NPs with LC<sub>50</sub> values of 3.50 to 7.01 ppm against II instar and

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

4.44 to 8.74 ppm against IV instar larvae of *Ae. aegypti* and 5.90 to 8.04 ppm for II instar and 4.90 to 9.55 ppm against IV instar of *An. stephensi*. Velayutham *et al.*, (2013) explored the larvicidal activity of green synthesized Ag NPs using aqueous bark extract of *F. racemosa* to control *Cx. quinquefasciatus* and *Cx. gelidus*. The maximum efficacy was observed in crude aqueous extract of *F. racemosa* ( $LC_{50}$ =67.72 and 63.70 mg/L) against the larvae of *Cx. quinquefasciatus* and *Cx. gelidus* and the synthesized Ag NPs ( $LC_{50}$ =12.00 and 11.21 mg/L).

Kalimuthu *et al.*, (2013) synthesized Ag NPs using *Cadaba indica* lam plant against *An. stephensi* and *Cx. quinquefasciatus*. Haldar *et al.*, (2013) synthesized highly stable nanoparticles of metallic silver by a simple, cost-effective, reproducible and previously unexploited biogenic source viz. dried green fruits of *Drypetes roxburghii* (Wall) and reported its mosquito larvicidal activity against *An. stephensi* and *Cx. quinquefasciatus*. Veerakumar *et al.*, (2013) synthesized Ag NPs by using *Sida acuta* plant leaf extract and determined their larvicidal activity against the late third instar larvae of *Cx. quinquefasciatus*, *An. stephensi*, and *Ae. aegypti*. The larvicidal activities of Ag NPs synthesized from fresh leaves, dry leaves and green berries of *S. nigrum* against larvae of *Cx. quinquefasciatus* and *An. stephensi* were tested by Rawani *et al.*, (2013). Roopan *et al.*, (2013) synthesised Ag NPs from the mesocarp layer extract of *Cocos nucifera* and assessed the anti-larvicidal agents against *An. stephensi* and *Cx. quinquefasciatus*.

Veerakumar *et al.*, (2014a) synthesized Ag NPs using *Helitropium indicum* plant leaves against late third instar larvae of *Ae. aegypti*, *An. stephensi* and *Cx. quinquefasciatus*. The  $LC_{50}$  and  $LC_{90}$  values of *H. indicum* aqueous leaf extract appeared to be effective against *An. stephensi* ( $LC_{50}$  - 68.73  $\mu$ g/mL;  $LC_{90}$  - 121.07  $\mu$ g/mL) followed by *Ae. aegypti* ( $LC_{50}$  -72.72  $\mu$ g/mL;  $LC_{90}$  - 126.86  $\mu$ g/mL) and *Cx. quinquefasciatus* ( $LC_{50}$  - 78.74  $\mu$ g/mL;  $LC_{90}$  - 134.39  $\mu$ g/mL). The  $LC_{50}$  and  $LC_{90}$  values of synthesized Ag NPs for *An. stephensi* were 18.40 and 32.45  $\mu$ g/mL, *Ae. aegypti* were 20.10 and 35.97  $\mu$ g/mL, and *Cx. quinquefasciatus* were 21.84 and 38.10  $\mu$ g/mL. Veerakumar *et al.*, (2014b) prepared Ag NPs utilizing *Feronia elephantum* plant leaf extract against late third-instar larvae of *An. stephensi*, *Ae. aegypti*, and *Cx. quinquefasciatus*. The  $LC_{50}$  and  $LC_{90}$  values of synthesized Ag NPs were 11.56 and 20.56  $\mu$ g mL<sup>-1</sup>, 13.13 and 23.12  $\mu$ g mL<sup>-1</sup> and 14.19 and 24.30  $\mu$ g mL<sup>-1</sup> for *An. stephensi*, *Ae. aegypti*, and *Cx. quinquefasciatus* respectively.

Naik *et al.*, (2014) tested the leaf mediated Ag NPs with *Pongamia pinnata* for mosquito control and found that plant extracts showed moderate larvicidal effects but the synthesized Ag NPs was found to be toxic to larvae at  $LC_{50}$  = 0.25 ppm and  $LC_{90}$  =1 ppm. Mondal *et al.*, (2014) investigated the bioactive components present in the root extract of *Parthenium hysterophorus* plant used for the biosynthesis of Ag NPs and analyzed the larvicidal effects of the extract as well as Ag NPs on *Cx. quinquefasciatus*. Suganya *et al.*, (2014) examined the larvicidal potential of solvent leaf extracts of *Leucas aspera* and synthesized Ag NPs using it against fourth instar larvae of *Ae. aegypti*. Rajasekharreddy and Rani (2014) synthesized silver-(protein-lipid) nanoparticles (Ag-PL NPs) (core-shell) using the seed extract from wild Indian Almond tree, *Sterculia foetida* (L.) (Sterculiaceae) and showed potential mosquito larvicidal activity against *Ae. aegypti* (L.), *An. stephensi* Liston and *Cx. quinquefasciatus* Say.

Priya *et al.*, (2014) evaluated the effect of plant synthesized Ag NPs through aqueous leaf extract of *Calotropis gigantea* to control dengue vector *Ae. aegypti* and malarial vector *An. stephensi*. Suresh *et al.*, (2014) synthesized Ag NPs using aqueous root extract of *Delphinium denudatum* (Dd) and investigated potent larvicidal activity against second instar larvae of dengue vector *Ae. aegypti* with a  $LC_{50}$  value of 9.6 ppm. Ramar *et al.*, (2014) determined the larvicidal activity of Ag NPs synthesized from aqueous leaf extract of *Cleistanthus collinus* against the larvae of *Ae. aegypti*. According to Dubey *et al.*, (2010) the potential of plants as biological materials for the synthesis of nanoparticles are yet to be fully explored.

#### IV. FUNGUS GENERATED NOVEL NANOPARTICLES: A NEW PROSPECTIVE FOR MOSQUITO CONTROL

Currently, fungi are being utilized in nanotechnology for the production of nanoparticles; synthesis using fungi has shown that this environmentally benign and renewable source can be used as an effective reducing agent for synthesis of Ag NPs and Au NPs (Table 2). Salunkhe *et al.*, (2011) synthesized Ag NPs by filamentous fungus *Cochliobolus lunatus* and tested it against second, third, and fourth instar larvae of *Ae. aegypti* ( $LC_{50}$ =1.29, 1.48, and 1.58;  $LC_{90}$ =3.08, 3.33, and 3.41 ppm respectively) and *An. stephensi* ( $LC_{50}$ =1.17, 1.30, and 1.41;  $LC_{90}$ =2.99, 3.13, and 3.29 ppm respectively).

Soni and Prakash, (2012a) synthesized fungus mediated Ag NPs using *Chrysosporium tropicum* as a larvicide against the second instar larvae of *Ae. aegypti*. Soni and Prakash, (2012b) described the larvicidal effect of extracellularly synthesised Au NPs with

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

*Aspergillus niger*. Soni and Prakash, (2012d) reported that the efficacy of *Chrysosporium tropicum*, a pathogenic fungus mediated Ag and Au NPs against *Cx. quinquefasciatus* and *An. stephensi*. Soni and Prakash, (2012c) tested the adulticidal efficacies of *C. keratinophilum*, *F. oxysporum f.sp. pisi* and *V. lecanii* against the adults of *Cx. quinquefasciatus* using the fungus mediated Ag NPs.

Table: 2 Mosquito control studies by using nanoparticles synthesized from fungus.

| Fungal Species  | Test NPs | Test Mosquito Species  | City / State        | Author / Year                          |
|---|----------|--|---------------------|--|
| <i>Cochliobolus lunatus</i>   | Ag       | <i>Ae. aegypti</i> , <i>An. stephensi</i>                                  | Jalgaon-Maharashtra | Salunkhe <i>et al.</i> , 2011          |
| <i>Chrysosporium tropicum</i>   | Au, Ag   | <i>Ae. aegypti</i>   | Agra-Delhi          | Soni and Prakash, 2012a                |
| <i>Aspergillus niger</i>  | Au       | <i>Ae. aegypti</i> , <i>An. stephensi</i> , <i>Cx. quinquefasciatus</i>    | Agra-Delhi          | Soni and Prakash, 2012b                |
| <i>Chrysosporium keratinophilum</i> ,<br><i>Fusarium oxysporum</i> ,<br><i>Verticillium lecanii</i> | Ag       | <i>Cx. quinquefasciatus</i>  | Agra-Delhi          | Soni and Prakash, 2012c                |
| <i>Chrysosporium tropicum</i>   | Au, Ag   | <i>Cx. quinquefasciatus</i> , <i>An. stephensi</i>                         | Agra-Delhi          | Soni and Prakash, 2012d                |
| <i>Chrysosporium keratinophilum</i> ,<br><i>Verticillium lecanii</i>                                | Au, Ag   | <i>Ae. aegypti</i> , <i>An. stephensi</i> ,<br><i>Cx. quinquefasciatus</i> | Agra-Delhi          | Soni and Prakash, 2013a                |
| <i>Fusarium oxysporum</i>   | Au, Ag   | <i>Ae. aegypti</i> , <i>An. stephensi</i> ,<br><i>Cx. quinquefasciatus</i> | Agra-Delhi          | Soni and Prakash, 2013b                |
| <i>Aspergillus niger</i>  | Ag       | <i>An. stephensi</i> , <i>Cx. quinquefasciatus</i> , <i>Ae. aegypti</i>    | Agra-Delhi          | Soni and Prakash, 2013c                |
| <i>Beauveria bassiana</i>   | Ag       | <i>Ae. aegypti</i>   | Madurai-TN          | Banu and Balasubramanian, 2014         |
| <i>Trichoderma harzianum</i>  | Ag       | <i>Ae. aegypti</i>   | Coimbatore- TN      | Sundaravadivelan and Padmanabhan, 2014 |
| <i>Penicillium verrucosum</i>   | Ag       | <i>Cx. quinquefasciatus</i>  | Trichy-TN           | Kamalakkannan <i>et al.</i> , 2014     |

Soni and Prakash, (2013a) proposed a green process for the extracellular production of Ag and Au NPs using the soil fungi *Chrysosporium keratinophilum* and *Verticillium lecanii* and investigated the effect of synthesized Ag NPs and Au NPs against the larvae and pupae of *An. stephensi*, *Cx. quinquefasciatus* and *Ae. aegypti*. Soni and Prakash, (2013b) synthesized Ag and Au NPs by using the cell free extract of *Fusarium oxysporum* fungus and tested them to be larvicides and pupicides against the larvae and pupae of *Cx. quinquefasciatus*, *An. stephensi* and *Ae. aegypti*. Soni and Prakash, (2013c) synthesized the Ag NPs by using the soil fungus *Aspergillus niger* and tested against the larvae and pupae of *An. stephensi*, *Cx. quinquefasciatus* and *Ae. aegypti*.

Banu and Balasubramanian, (2014) synthesized silver biolarvicide with the help of entomopathogenic fungi, *Beauveria bassiana*, and assessed it against the different larval instars of dengue vector, *Ae. aegypti*. Sundaravadivelan and Padmanabhan, (2014) investigated the larvicidal and pupicidal effect of mycosynthesized Ag NPs using an entomopathogenic fungi *Trichoderma harzianum* against developmental stages of the dengue vector *Ae. aegypti*. Kamalakkannan *et al.*, (2014) attempted on laboratory evaluation of mycosynthesized Ag NPs from fungus *Penicillium verrucosum* and tested against larvae of filarial vector, *Cx. quinquefasciatus*.

### V. ECO-FRIENDLY BACTERIAL ROUTE TO SYNTHESIZE NPS AND ITS APPLICATION IN MOSQUITO CONTROL

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

Two insecticidal bacteria have been used as larvicides to control larvae of nuisance and vector mosquitoes in many countries, *Bacillus thuringiensis ssp. israelensis* and *Bacillus sphaericus* (Wirth *et al.*, 2010).

Marimuthu *et al.*, (2013) investigated the larvicidal activities of synthesized cobalt nanoparticles (Co NPs) using bio control agent, *Bacillus thuringiensis* against malaria vector, *An. subpictus* and dengue vector, *Ae. aegypti* (Diptera: Culicidae). Dhanasekaran and Thangaraj (2013) evaluated the larvicidal activity of biogenic nanoparticles of *Agaricus bisporus*, *E. coli*, *Pencillium sp.* and *Vibrio sp.* against filariasis causing, *Culex* mosquito vector.

The mortality rate of *Culex* larvae, using *Agaricus bisporus* biogenic nanoparticles were 100% (5 mg/L), 81% (2.5 mg/L), 62% (1.25 mg/L), 28% (0.625 mg/L) and 11% (0.312 mg/L). The results suggested that synthesized biogenic Ag NPs have the potential to be used as an ideal eco-friendly approach for controlling *Culex* species. Banu *et al.*, (2014) revealed that the larvicidal activity of Ag NPs synthesized using *Bacillus thuringiensis* (Bt) against *Ae. aegypti* would control many diseases of public health importance (Table 3). According to Owolade *et al.*, (2008) nanoparticles help to produce new pesticides, insecticides and insect repellants. Over the next two decades, the Green Revolution would be accelerated by means of nanotechnology.

Table: 3 Mosquito control studies by using nanoparticles synthesized from bacteria.

| Bacterial Species                     | Test NPs | Test Species                              | City / State | Author / Year                     |
|---------------------------------------|----------|---|--------------|-----------------------------------|
| <i>Bacillus thuringiensis</i>         | Co       | <i>An. subpictus</i> , <i>Ae. aegypti</i> | Vellore-TN   | Marimuthu <i>et al.</i> , (2013)  |
| <i>E. coli</i> , <i>Pencillium sp</i> | Ag       | <i>Culex sp.</i>                          | Trichy-TN    | Dhanasekaran and Thangaraj (2013) |
| <i>Bacillus thuringiensis</i>         | Ag       | <i>Ae. aegypti</i>                        | Madurai-TN   | Banu <i>et al.</i> , (2014)       |

### VI. TREND OF NANOPARTICLES IN MOSQUITO CONTROL

Green synthesis of nanoparticles in mosquito control using plant extracts, fungi and bacteria collected here can be sorted out as 31 on plants, 11 on fungi and 3 on bacteria in terms of subjects (Fig. 1A). Plant extracts were the prototype organisms among them. And they can be sorted as 20 on leaf, 2 on plant latex, 2 on root, 1 each on stem, fruit, seed, bark, berry, mesocarp and aerial parts respectively. Ag NPs were used in majority of the studies followed by Au NPs. In particular, *Ae. aegypti*, *An. stephensi* and *Cx. quinquefasciatus* were favourites for many researchers. Out of the 45 studies performed, 6 were conducted in 2011, 12 in 2012, 14 in 2013 and 13 in 2014 so far. Almost all the plant extract studies used Ag NPs, while fungal studies used both Ag and Au NPs in mosquito control. In terms of nationality, all the researches were conducted in India. About 62 % of the researches were conducted in Tamil Nadu followed by Delhi 16%, Maharashtra 9%, West Bengal 7%, Andhra Pradesh 4% and Kerala 2% (Fig. 1B). Researchers used only Ag, Au and Co NPs for mosquito control till now. Most of the fungal studies were carried out by Soni and Prakash and the studies on bacteria for NPs synthesis are very meager.




Fig. 1. Studies of nanoparticles in mosquito control as related with (A) green synthesis of nanoparticles and (B) states.



## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

In almost all the studies, the characterization and the structure determination of these nanoparticles were performed using the application of modern scientific instruments such as UV-VIS spectrophotometry, Transmission Electron Microscopy, High-Resolution Transmission Electron Microscopy (HR-TEM), Atomic Force Microscopy (AFM), Field Emission Scanning Electron Microscopy (FE-SEM), X-ray Diffraction (XRD), Fourier Transform Infrared Spectroscopy (FT-IR) and Inductively Coupled Plasma Mass Spectrometry (ICP-MS).

### VII. CONCLUSION

These results revealed that the green, biological synthesis of silver/gold nano particles have the potential to be utilized as a good, rapid, eco-friendly approach for the control of mosquito population. It is totally a new pathway but, can be effectively utilized for the efficient killing of mosquitoes. Therefore, biological control can thus provide an effective and environmental friendly approach, which can be used as an alternative to minimize the mosquito population. To understand the current research trends of nanoparticles in mosquito control, research papers on NPs synthesised using biological organisms such as plant extracts, fungi and bacteria were thoroughly analyzed and discussed in terms of the type of nanoparticles, test species, exposure medium and suitable concentration. The researches demonstrated a wide range of results even when using the same nanoparticles. This was because, the particle size, surface coating, and the test medium supposedly made difference in the results. Therefore, in future, researches can be conducted by considering the above factors also.

### VIII. ACKNOWLEDGEMENT

The authors are grateful to Dr. G. B. Brindha Devi, Assistant Professor, Department of Zoology, Queen Mary's College and Ms. S. P. Preethi, MBBS, Government Sivagangai Medical College, Tamil Nadu, India for their assistance in reviewing the paper.

### REFERENCES

- [1]. Salam HA, Kamaraj RPM, Jagadeeswaran P, Gunalan S, Sivaraj R, "Plants: green route for nanoparticle synthesis," *Int Res J Biol Sci.*, vol. 1 (5), pp. 85-90, 2012.
- [2]. Adhikari U, Ghosh A, Chandra G, "Nano particles of herbal origin: A recent eco-friend trend in mosquito control," *Asian Pac J Trop Dis.*, vol. 3 (2), pp. 167-168, 2013.
- [3]. Marimuthu S, Rahuman AA, Rajakumar G, Santhoshkumar T, Kirthi AV, Jayaseelan C, Bagavan A, Zahir AA, Elango G, Kamaraj C, "Evaluation of green synthesized silver nanoparticles against parasites," *Parasitology Research*, vol. 108 (6), pp. 1541-1549, 2011.
- [4]. Santhoshkumar T, Rahuman AA, Rajakumar G, Marimuthu S, Bagavan A, Jayaseelan C, "Synthesis of silver nanoparticles using *Nelumbo nucifera* leaf extract and its larvicidal activity against malaria and filariasis vectors," *Parasitol Res.*, vol.108, pp. 693-702, 2011.
- [5]. Rajakumar G, Abdul Rahuman A, "Larvicidal activity of synthesized silver nanoparticles using *Eclipta prostrata* leaf extract against filariasis and malaria vectors," *Acta Trop.*, vol. 118, pp. 196-203, 2011.
- [6]. Gnanadesigan M, Anand M, Ravikumar S, Maruthupandy M, Vijayakumar V, Selvam S, Dhineshkumar M, Kumaraguru AK, "Biosynthesis of silver nanoparticles by using mangrove plant extract and their potential mosquito larvicidal property," *Asian Pacific Journal of Tropical Medicine*, pp. 799-803, 2011.
- [7]. Jayaseelan C, Rahuman AA, Rajakumar G, Kirthi VA, Santhoshkumar T, Marimuthu S, "Synthesis of pediculocidal and larvicidal silver nanoparticles by leaf extract from heartleaf moonseed plant, *Tinospora cordifolia* Miers," *Parasitol Res.*, vol. 109, pp. 185-194, 2011.
- [8]. Arjunan NK, Murugan K, Rejeeth C, Madhiyazhagan P, Barnard DR, "Green Synthesis of Silver Nanoparticles for the Control of Mosquito Vectors of Malaria, Filariasis, and Dengue," *Vector-Borne and Zoonotic Diseases*, vol. 12 (3), pp. 262-268, 2012.
- [9]. Priyadarshini KA, Murugan K, Panneerselvam C, Ponarulselvam S, Hwang JS, Nicoletti M, 2012, "Biolarvicidal and pupicidal potential of silver nanoparticles synthesized using *Euphorbia hirta* against *Anopheles stephensi* Liston (Diptera: Culicidae)," *Parasitology Research*, vol. 111 (3), pp. 997-1006.
- [10]. Sareen SJ, Pillai RK, Chandramohanakumar N, Balagopalan M, "Larvicidal potential of biologically synthesised silver nanoparticles against *Aedes Albopictus*," *Res J Rec Sci.*, vol. 1, pp. 52-56, 2012.
- [11]. Patil CD, Borase HP, Patil SV, Salunkhe RB, Salunke BK, "Larvicidal activity of silver nanoparticles synthesized using *Pergularia daemia* plant latex against *Aedes aegypti* and *Anopheles stephensi* and nontarget fish *Poecilia reticulata*," *Parasitology Research*, vol. 111 (2), pp. 555-562, 2012a.
- [12]. Patil CD, Patil SV, Borase HP, Salunke BK, Salunkhe RB, 2012b, "Larvicidal activity of silver nanoparticles synthesized using *Plumeria rubra* plant latex against *Aedes aegypti* and *Anopheles stephensi*," *Parasitology Research*, vol.110 (5), pp. 1815-1822, 2012b.
- [13]. Roni M, Murugan K, Panneerselvam C, Subramaniam J, Hwang JS, "Evaluation of leaf aqueous extract and synthesized silver nanoparticles using *Nerium oleander* against *Anopheles stephensi* (Diptera: Culicidae)," *Parasitol Res.*, DOI 10.1007/s00436-012-3220-3, 2012.
- [14]. Subarani S, Sabhanayakam S, Kamaraj C, "Studies on the impact of biosynthesized silver nanoparticles (AgNPs) in relation to malaria and filariasis vector control against *Anopheles stephensi* Liston and *Culex quinquefasciatus* Say (Diptera: Culicidae)," *Parasitol Res.*, DOI 10.1007/s00436-012-3158-5, 2012.
- [15]. Sundaravadivelan C, Nalini M, "Biolarvicidal effect of phyto-synthesized silver nanoparticles using *Pedilanthus tithymaloides* (L.) Poit stem extract against the dengue vector *Aedes aegypti* L. (Diptera: Culicidae)," *Asian Pac J Trop Biomed.*, pp. 1-8, 2012.
- [16]. Suman TY, Elumali D, Kaleena PK, Rajasree SRR, "GCMS analysis of bioactive components and synthesis of silver nanoparticle using *Ammannia baccifera*

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

- aerial extract and its larvicidal activity against malaria and filariasis vectors," *Ind Crop Prod.*, vol. 47, pp. 39-245, 2013.
- [17]. Arjunan NK, Jeyalalitha T, Murugan K, Madhiyazhagan P, "Bioefficacy of plant-mediated gold nanoparticles and *Anthocephalus cadamba* on filarial vector, *Culex quinquefasciatus* (Insecta: Diptera: Culicidae)," *Parasitology Research.*, vol. 112 (3), pp. 1053-1063, 2013.
- [18]. Borase HP, Patil CD, Salunkhe RB, Narkhede CP, Salunke BK, Patil SV, "Phyto-Synthesized Silver Nanoparticles: A Potent Mosquito Biolarvicidal Agent," *J Nanomedicine Biotherapeutic Discov.*, vol. 3 (1), <http://dx.doi.org/10.4172/2155-983X.1000111>, 2013.
- [19]. Velayutham K, Rahuman AA, Rajakumar G, Roopan SM, Elango G, Kamaraj C, Marimuthu S, Santhoshkumar T, Iyappan M, Siva C, "Larvicidal activity of green synthesized silver nanoparticles using bark aqueous extract of *Ficus racemosa* against *Culex quinquefasciatus* and *Culex gelidus*," *Asian Pacific Journal of Tropical Medicine*, vol. 6 (2), pp. 95-101, 2013.
- [20]. Kalimuthu K, Panneerselvam C, Murugan K, Hwang JS, "Green synthesis of silver nanoparticles using *Cadaba indica* lam leaf extract and its larvicidal and pupicidal activity against *Anopheles stephensi* and *Culex quinquefasciatus*," *Journal of Entomological and Acarological Research.*, vol. 45 (2), 2013.
- [21]. Haldar KM, Haldar B, Chandra G, "Fabrication, characterization and mosquito larvicidal bioassay of silver nanoparticles synthesized from aqueous fruit extract of putranjiva, *Drypetes roxburghii* (Wall)," *Parasitol Res.*, doi: 10.1007/s00436-013-3288-4, 2013.
- [22]. Veerakumar K, Govindarajan M, Rajeswary M, "Green synthesis of silver nanoparticles using *Sida acuta* (Malvaceae) leaf extract against *Culex quinquefasciatus*, *Anopheles stephensi*, and *Aedes aegypti* (Diptera: Culicidae)," *Parasitology Research.*, vol. 112 (12), pp. 4073-4085, 2013.
- [23]. Rawani A, Ghosh A, Goutam Chandra, "Mosquito larvicidal and antimicrobial activity of synthesized nano-crystalline silver particles using leaves and green berry extract of *Solanum nigrum* L. (Solanaceae: Solanales)," *Acta Tropica*, vol. 128 (3), pp. 613-622, 2013.
- [24]. Roopan SM, Rohit, Madhumitha G, Rahuman A, Kamaraj C, Bharathi A, Surendra TV, "Low-cost and eco-friendly phyto-synthesis of silver nanoparticles using *Cocos nucifera* coir extract and its larvicidal activity," *Industrial Crops and Products.*, vol. 43, pp. 631-635, 2013.
- [25]. Veerakumar K, Govindarajan M, Rajeswary M, Muthukumaran U, "Mosquito larvicidal properties of silver nanoparticles synthesized using *Heliotropium indicum* (Boraginaceae) against *Aedes aegypti*, *Anopheles stephensi*, and *Culex quinquefasciatus* (Diptera: Culicidae)," *Parasitology Research.*, vol. 113 (6), pp. 2363-2373, 2014a.
- [26]. Veerakumar K, Govindarajan M, Rajeswary M, Muthukumaran U, "Low-cost and eco-friendly green synthesis of silver nanoparticles using *Feronia elephantum* (Rutaceae) against *Culex quinquefasciatus*, *Anopheles stephensi*, and *Aedes aegypti* (Diptera: Culicidae)," *Parasitology Research*, vol. 113 (5), pp. 1775-1785, 2014b.
- [27]. Naik BR, Gowreeswari GS, Singh Y, Satyavathi R, Daravath SS, Reddy PR, "Bio-Synthesis of Silver Nanoparticles from Leaf Extract of *Pongamia pinnata* as an Effective Larvicide on Dengue Vector *Aedes albopictus* (Skuse) (Diptera: Culicidae)," *Advances in Entomology*, vol. 2 (2) Article ID: 45433, 9 pages, DOI:10.4236/ae.2014.22016, 2014.
- [28]. Mondal NK, Chowdhury A, Dey U, Mukhopadhyaya P, Chatterjee S, Das K, Datta JK, "Green synthesis of silver nanoparticles and its application for mosquito control," *Asian Pac J Trop Dis.*, vol. 4, pp. S204-S210, 2014.
- [29]. Suganya G, Karthi S, Shivakumar MS, 2014, "Larvicidal potential of silver nanoparticles synthesized from *Leucas aspera* leaf extracts against dengue vector *Aedes aegypti*," *Parasitology Research*, vol. 113 (3), pp. 875-880, 2014.
- [30]. Rajasekharreddy P, Rani PU, "Biofabrication of Ag nanoparticles using *Sterculia foetida* L. seed extract and their toxic potential against mosquito vectors and HeLa cancer cells," *Materials Science and Engineering: C*, vol. 39 (1), pp. 203-212, 2014.
- [31]. Priya S, Murugan K, Priya A, Dinesh D, Panneerselvam C, Devi GD, Chandramohan B, Kumar PM, Barnard DR, Rui-De Xue, Jiang-Shiou Hwang, Nicoletti M, Chandrasekar R, Amsath A, Bhagooli R, Hui Wei, "Green synthesis of silver nanoparticles using *calotropis gigantea* and their potential mosquito larvicidal property," *Int. J. Pure Appl. Zool.*, vol. 2 (2), pp. 128-137, 2014.
- [32]. Suresh G, Gunasekar PH, Kokila D, Prabhu D, Dinesh D, Ravichandran N, Ramesh B, Koodalingam A, Siva GV, "Green synthesis of silver nanoparticles using *Delphinium denudatum* root extract exhibits antibacterial and mosquito larvicidal activities," *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, vol. 127 (5), pp. 61-66, 2014.
- [33]. Ramar G, Suman T, Elangomathavan R, Jeyasankar A, "Larvicidal activity of biologically synthesised silver nanoparticles against dengue vector *Aedes aegypti* (Culicidae)," *Discovery*, vol. 9 (23), pp. 65-68, 2014.
- [34]. Dubey SP, Lahtinen M, Sillanpaa M, "Tansy fruit mediated greener synthesis of silver and gold nanoparticles process," *Biochem.*, vol. 45, pp. 1065-1071, 2010.
- [35]. Salunkhe RB, Patil SV, Patil CD, Salunke BK, "Larvicidal potential of silver nanoparticles synthesized using fungus *Cochliobolus lunatus* against *Aedes aegypti* (Linnaeus, 1762) and *Anopheles stephensi* Liston (Diptera: Culicidae)," *Parasitol Res.*, vol. 109, pp. 823-831, 2011.
- [36]. Soni N, Prakash S, "Efficacy of fungus mediated silver and gold nanoparticles against *Aedes aegypti* larvae," *Parasitol Res.*, vol. 110, pp. 175-84, 2012a.
- [37]. Soni N, Prakash S, "Synthesis of gold nanoparticles by the fungus *Aspergillus niger* and its efficacy against mosquito larvae," *Reports in Parasitology*, vol. 2, pp. 1-7.
- [38]. Soni N, Prakash S, "Entomopathogenic fungus generated nanoparticles for enhancement of efficacy in *Culex quinquefasciatus* and *Anopheles stephensi*," *Asian Pac J Trop Dis.* Vol. 2 (2), pp. S356-S361, 2012d.
- [39]. Soni N, Prakash S, "Fungal-mediated nano silver: an effective adulticide against mosquito," *Parasitol Res.*, vol. 111, pp. 2091-2098, 2012c.
- [40]. Soni N, Prakash S, "Microbial synthesis of spherical nanosilver and nanogold for mosquito control," *Annals of Microbiology*, 2013a.
- [41]. Soni N, Prakash S, "Fungus generated novel nanoparticles: a new prospective for mosquito control," *International Journal of Recent Scientific Research*, vol. 4 (10), pp. 481-1487, 2013b.
- [42]. Soni N, Prakash S. Possible Mosquito Control by Silver Nanoparticles Synthesized by Soil Fungus (*Aspergillus niger* 2587). *Advances in Nanoparticles*, 2013c; 2:125-132.
- [43]. Banu AN, Balasubramanian C, "Mycosynthesis of silver nanoparticles using *Beauveria bassiana* against dengue vector, *Aedes aegypti* (Diptera: Culicidae)," *Parasitology Research*, 2014.
- [44]. Sundaravadivelan C, Padmanabhan MN, "Effect of mycosynthesized silver nanoparticles from filtrate of *Trichoderma harzianum* against larvae and pupa of dengue vector *Aedes aegypti* L.," *Environmental Science and Pollution Research*, vol. 21 (6), pp. 4624-4633, 2014.
- [45]. Kamalakannan S, Gobinath C, Ananth S, "Synthesis and Characterization of Fungus Mediated Silver Nanoparticle for Toxicity on Filarial Vector, *Culex*

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

- quinquefasciatus*," *Int. J. Pharm. Sci. Rev. Res.*, vol. 24 (2), pp. 124-132, 2014.
- [46]. Wirth MC, Walton WE, Federici BA, "Evaluation of Resistance to the *Bacillus sphaericus* Bin Toxin is Phenotypically Masked by Combination with the Mosquitocidal Proteins of *Bacillus thuringiensis* Subspecies *israelensis*," *Environmental Microbiology*, vol. 12 (5), pp. 1154-1160, 2010.
- [47]. Marimuthu S, Rahuman AA, Kirthi AV, Santhoshkumar T, Jayaseelan C, Rajakumar G, "Eco-friendly microbial route to synthesize cobalt nanoparticles using *Bacillus thuringiensis* against malaria and dengue vectors," *Parasitology Research*, vol. 112 (12), pp. 4105-4112, 2013.
- [48]. Dhanasekaran D, Thangaraj R, "Evaluation of larvicidal activity of biogenic nanoparticles against filariasis causing *Culex* mosquito vector," *Asian Pac J Trop Dis.*, vol. 3 (3), pp. 174-179, 2013.
- [49]. Banu AN, Balasubramanian C, Moorthi PV, "Biosynthesis of silver nanoparticles using *Bacillus thuringiensis* against dengue vector, *Aedes aegypti* (Diptera: Culicidae)," *Parasitology Research*, vol. 113 (1), pp. 311-316, 2014.
- [50]. Owolade OF, Ogunleti DO, Adenekan MO, "Titanium Dioxide affects disease development and yield of edible cowpea," *EJEA Chem.*, vol. 7 (50), pp. 2942-2947.



10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)