



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 6 Issue: III Month of publication: March 2018 DOI: http://doi.org/10.22214/ijraset.2018.3352

www.ijraset.com

Call: 🕥 08813907089 🔰 E-mail ID: ijraset@gmail.com



Special Features of Some Dinuclear Copper (II) Complexes

Dr. Gopal Chandra Giri Bhairab Ganguly College, Belgharia, Kolkata-700056

Abstract: This article describes synthesis, characterization and properties of some dinucler copper complexes.

I. INTRODUCTION

The dinuclear copper(II) complexes have received a great deal of attention because of the role of dicopper centers in biological processes¹⁻⁸ and magnetic exchange interactions.^{9,10} Copper has been known as an essential bio-element for some time but its biological role(s) has been known only in the last decades due to the rapid development of bioinorganic chemistry.^{4,11-15} Copper-containing enzymes are involved in various processes in living systems. Hemocyanin (O₂ transport), tyrosinase (hydroxylation of monophenols) and catechol oxidase (oxidation of catechols) are classified as type 3 copper proteins (Fig.1) and have magnetically coupled dicopper(II) centers at their active sites. As a result, the biomimetic dinuclear copper(II) complexes have been extensively studied using several analytical techniques such as electronic absorption



L = bridging ligand

Fig. 1 General coordination modes in the type 3 copper proteins.

spectroscopy, EPR, X-ray crystallography and magnetic susceptibility measurements. ¹H NMR spectroscopy has also been emerged as a window for the investigation of structural and magnetic properties of dinuclear copper(II) complexes.

II. RESULTS AND DISCUSSION

Holz *et. al.*¹⁶ have reported one and two dimensional ¹H NMR studies on a series of magnetically diverse μ -alkoxo and μ -phenexo bridged dinuclear copper(II) complexes. These complexes $[Cu_2(L^{1a})(OAc)]^{2+}(1)$, $[Cu_2(L^{2a})(OH)]^{2+}(2)$, $[Cu_2(Me-L^{3a})(OH)]^{2-}(3)$ and $[Cu_2(L^{4a})(OH)]^{2+}(4)$ (where, **HL**^{1a} = *N*,*N*,*N*,*N*'/v'-tetrakis[(2-benzimidazolyl)-methyl]-2-hydroxy-1,3-diaminopropane; **HL**^{2a} = 2,6-bis[{bis(2-pyridylmethyl)amino}methyl]-4-methylphenol; **H**₅**Me-L**^{3a} = *N*,*N'*-(2-Hydroxy-5-methyl-1,3-xylylene)bis(N-carboxymethylglycine) and **HL**^{4a} = 2,6-bis[{bis(2-pyridylethyl)amino}methyl]phenol) (Fig.2) have been characterized by single crystal X-ray crystallography as well as several spectroscopic techniques.



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 6.887 Volume 6 Issue III, March 2018- Available at www.ijraset.com





Fig. 2 Structures of $[Cu_2(L^{1a})(OAc)]^{2+}(1)$, $[Cu_2(L^{2a})(OH)]^{2+}(2)$, $[Cu_2(Me-L^{3a})(OH)]^{2-}(3)$ and $[Cu_2(L^{4a})(OH)]^{2+}(4)$.

In 2004, Powell *et. al.*¹⁷ have reported some dinuclear copper(II) complexes (Fig.3) with carboxylate rich dinucleating ligands $(H_5Me-L^{3a} \text{ and } H_5MeO-L^{3a})$ and studied their magnetic properties at low temperature. In these complexes, phenoxo oxygen atom of the ligand acts as a spacer-cum-bridging group to bind the two copper(II) centers. Low temperature magnetic susceptibility measurements have revealed that the copper(II) centers are antiferromagnetic ally coupled.



Fig. 3 Structures of $[Cu_2(Me-L^{3a})(H_2O)_2]^-$ (5) and $[Cu_2(H_2MeO-L^{3a})(\mu-Cl)(CH_3OH)]$ (6).

Striegler *et. al.*¹⁸ have reported a sugar discriminating alkoxo bridged dinuclear copper(II) complex, $[Cu_2(L^{5a})(OAc)](CIO_4)_2$ (7) (where, $HL^{5a} = 1,3$ -bis[(pyridin-2-ylmethyl)amino]propan-2-ol (Fig. I.4). The reaction of complex 7 with different sugar substrates at pH~12.40 gives 1:1 complex/substrate bound product, resulting in two-state systems. This complex shows a remarkable selectivity during coordination to the underivatized monosaccharides under strong binding conditions in alkaline solution. Recently, they have also studied the different dinuclear copper(II) complexes $[Cu_2(L^{5a})(OAc)](CIO_4)_2$ (7), its unsymmetrical analogue complex (8) and $[Cu_2(L^{6a})(OAc)]$ (9) (where, $H_3L^{6a} = N,N'$ -{1,3-bis[{2-hydroxy-4-[2-(2-methoxyethoxy)-ethoxy]-ethoxy]-ethoxy}benzylideneamino]propan-2-ol} (Fig.4) as new catalysts for the cleavage of glycosidic bonds to open a new direction towards the development of selective artificial glycosidase mimics.¹⁹





International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 6.887

Volume 6 Issue III, March 2018- Available at www.ijraset.com



Fig. 4 Structures of $[Cu_2(L^{5a})(OAc)]^{2+}(7)$, its unsymmetrical analogue (8) and $[Cu_2(L^{6a})(OAc)]$ (9).

Recently, a series of three water-soluble dicopper(II) complexes of benzoic acid derivatives are reported.²⁰ All the three complexes induce oxidative DNA cleavage in the presence of H_2O_2 and noticeable DNA cleavage activity independent from the functional groups (F, Cl, NO₂) attached to the benzene ring. These complexes also display the selective cytotoxic effect against HepG2 cancer cells (2-fold higher) compared to that against the normal cells.

Buchanan and co-workers²¹ have reported the dinuclear copper(II) complexes $[Cu_2(L^{7a})(pz)](ClO_4)_2 \cdot 2CH_3CN$ (10) and $[Cu_2(L^{7a})(OAc)](ClO_4)_2 \cdot CH_3CN$ (11) where, $HL^{7a} = N, N, N', N'$ -tetrakis[(1-methylimidazol-2-yl)-methyl]-1,3-diaminopropan-2-ol (Fig.5) and pz is the pyrazolate anion. The single crystal X-ray structures reveal that both complexes 10 and 11 contain bridging alkoxide ligand including bridging pyrazolate and acetate ions, respectively. The magnetochemical characteristics of these complexes provide the evidences for "magnetic orbital switching". Complex 10 shows strong antiferromagnetic coupling (J = -130 cm⁻¹) between the two copper(II) centers, while complex 11 exhibits the ferromagnetic coupling (J = +16.4 cm⁻¹) between the two copper(II) centers. They have rationalized the differences in the magnetic behavior of these two complexes using the "ligand orbital complementary" concept. The ground state magnetic orbital involved in spin coupling in complex 10 (d_x²-y²) is different from that in complex 11 (d_z²).



Fig. 5 Structure of *N*,*N*,*N*',*N*'-tetrakis[(1-methylimidazol-2-yl)-methyl]-1,3-diaminopropan-2-ol (HL^{7a}).

One of the well-characterized Cu/O₂ systems is based on the dinucleating ligand HL^{8a} (Fig.6).²² Oxygenation of its dicopper(I) complex (12) affords a species (12a)



Fig. 6 Ligands derived from HL^{8a}.



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 6.887 Volume 6 Issue III, March 2018- Available at www.ijraset.com

with intense absorption bands at 505 nm ($\varepsilon = 6300 \text{ M}^{-1}\text{cm}^{-1}$) and 610 nm ($\varepsilon = 2400 \text{ M}^{-1}\text{cm}^{-1}$)^{sh}, attributed to peroxide to copper(II) charge transfer transitions. Oxygenation of the dicopper(I) complex (**13**) of the protonated ligand **HL**^{8a} produces a hydroperoxide species (**13a**) (Fig.7) exhibiting a charge transfer transition at 395 nm ($\varepsilon = 8000 \text{ M}^{-1}\text{cm}^{-1}$) with a shoulder at 450 nm.²³ The Cu---Cu separation of ~ 3.0 Å measured by EXAFS is most consistent with that between two copper(II) ions in a μ -1,1 hydroperoxide bridged dicopper(II) complex. This species can also be generated by protonating the peroxide complex derived from the ligand (**L**^{8a})⁻ (Fig. I.7, (**12a**) \rightarrow (**13a**)). Similar peroxo and/or hydroperoxo species can also be formed when the dinucleating ligands bearing pyrazoles or mixed pyrazole/pyridine subunits are used (Fig.I.6). The corresponding η^{1} -peroxo complexes have characteristic charge transfer features in the regions of ~ 510 and ~ 620 nm, whereas the μ -1,1-hydroperox species has the charge transfer band at ~ 400 nm.²⁴



Fig. 7 Reactivity of copper(I) complexes of HL^{8a} and (L^{8a})⁻ ligands with O₂.

III. CONCLUSIONS

Properly designed di- and polynuclear metal complexes would provide new reactivity patterns and physical properties that could not be achieved with similar mononuclear complexes. Polynuclear complex systems are expected to have greater oxidizing and reducing power and the neighboring metal centers are expected to cooperate in promoting reactions and new electronic interactions leading to the distinct physical properties. The findings of this study will contribute towards designing the new polynuclear transition metal complexes.

IV. ACKNOWLEDGEMENTS

The author acknowledges Dr. Manindranath Bera, University of Kalyani for his nonstop encouragement and continuous support.

REFERENCES

- Karlin, K. D. Science 1993, 261, 701-708; (b) Karlin, K. D.; Tyeklar, Z. Eds.; Bioinorganic Chemistry of Copper; Chapman and Hall: New York, London, 1993.
- (a) Solomon, E. I.; Lowery, M. D.; Lacroix, L. B.; Root, D. E. Methods Enzymol. 1993, 226, 1-33; (b) Solomon, E. I.; Tuezek, F.; Root, D. E.; Brown, C. A. Chem. Rev. 1994, 94, 827; (c) Solomon, E. I.; Sundaram, U. M.; Mackonkin, T. E. Chem. Rev. 1996, 96, 2563-2606
- [3] Magnus, K. A.; Ton-That, H.; Carpenter, J. E. Chem. Rev. 1994, 94, 727-735
- [4] Kitajima, N.; Moro-Oka, Y. Chem. Rev. 1994, 94, 737-757.
- [5] Ryan, S.; Adams, H.; Fenton, Becker, D. E. M.; Schindler, S. Inorg. Chem. 1998, 37, 2134.
- [6] Holland, P. L.; Tolman, W. B. Coord. Chem. Rev. 1999, 190-192, 855-869.
- [7] Decker, H.; Dillinger, R.; Tuczek, F. Angew. Chem., Int. Ed. 2000, 39, 1591.
- [8] Gerdemann, C.; Eicken, C.; Krebs, B. Acc. Chem. Res. 2002, 35, 183-191.
- [9] Reger, D. L.; Pascui, A. E.; Smith, M. D.; Jezierska, J.; Ozarowski, A. Inorg. Chem. 2015, 54, 1487-1500.
- [10] (a) Gusev, A. N.; Shulgin, V. F.; Zamnius, E. A.; Ricci, M.; Minin, V. V.; Aleksandrov, G. G.; Eremenko, I. L.; Linert, W. Inorg. Chim. Acta 2015, 430, 120-124; (b) Massoud, S. S.; Spell, M.; Ledet, C. C.; Junk, T.; Herchel, R.; Fischer, R. C.; Trávníček, Z.; Mautner, F. A. Dalton Trans. 2015, 44, 2110-2121
- [11] Linder, M. C.; Goode, C. Biochemistry of Copper, Plenum Press, New York, 1991.
- [12] Kaim, W.; Rall, J. Angew. Chem., Int. Ed. 1996, 35, 43.
- [13] Kitajima, N.; Moro-oka, Y. J. Chem. Soc., Dalton Trans. 1993, 2665.



International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 6.887

Volume 6 Issue III, March 2018- Available at www.ijraset.com

- [14] Jacobson, R. R.; Tyeklar, Z.; Farooq, A.; Karlin, K. D.; Liu, S.; Zubieta, J. J. Am. Chem. Soc. 1988, 110, 3690-3692.
- [15] Tyeklar, Z.; Karlin, K. D. Acc. Chem. Res. 1988, 22, 241.
- [16] Brink, J. M.; Rose, R. A.; Holz, R. C. Inorg. Chem. 1996, 35, 2878.
- [17] Laborda, S.; Clerac, R.; Anson, C. E.; Powell, A. K. Inorg. Chem. 2004, 43, 5931.
- [18] Striegler, S.; Dittel, M. J. Am. Chem. Soc. 2003, 125, 11518-11524.
- [19] Striegler, S.; Dunaway, N. A.; Gichinga, M. G.; Barnett, J. D.; Nelson, A. G. D. Inorg. Chem. 2010, 49, 2639.
- [20] Lim, E. K.; Teoh, S. G.; Goh, S. M.; Ch'ng, C. D.; Ng, C. H.; Fun, H. K.; Rosli, M. M.; Najimudin, N.; Beh, H. K.; Seow, L. J.; Ismail, N.; Ismail, Z.; Asmawi, M. Z.; Cheah, Y. H. Polyhedron 2009, 28, 1320-1330.
- [21] Nie, H.; Aubin, S. M. J.; Mashuta, M. S.; Porter, R. A.; Richardson, J. F.; Hendrickson, D. N.; Buchanan, R. M. Inorg. Chem. 1996, 35, 3325.
- [22] (a) Karlin, K. D.; Kaderli, K.; Zuberbuhler, A. D. Acc. Chem. Res. 1997, 30, 139-147; (b) Karlin, K. D.; Cruse, R. W.; Gultneh, Y.; Farooq, A.; Hayes J. C.; Zubieta, J. J. Am. Chem. Soc. 1987, 109, 2668-2679; (c) Cruse, R. W.; Kaderli, S.; Karlin, K. D.; Zuberbuhler, A. D. J. Am. Chem. Soc. 1988, 110, 6882-6883; (d) Karlin, K. D.; Hays, J. C.; Gultneh, Y.; Cruse, R. W.; Mckown, J. W.; Hutchinson, J. P.; Zubieta, J. J. Am. Chem. Soc. 1984, 106, 2121-2128; (e) Suzuki, M.; Furutachi, H.; Okawa, H. Coord. Chem. Rev. 2000, 200, 105-129.
- [23] (a) Karlin, K. D.; Cruse, R. W.; Gultneh, Y. J. Chem. Soc., Chem. Commun. 1987, 599-600; (b) Karlin, K. D.; Ghosh, P.; Cruse, R. W.; Farooq, A.; Gultneh, Y.; Jacobson, R. R.; Blackburn, N. J.; Strange, R. W.; Zubieta, J. J. Am. Chem. Soc. 1988, 110, 6769-6780.
- [24] Root, D. E.; Mahrooftahir, M.; Karlin, K. D.; Solomon, E. I. Inorg. Chem. 1998, 37, 4838.











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24*7 Support on Whatsapp)