



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: TPAM-2055 onferendelonth of publication: March 2018

www.ijraset.com

Call: 🛇 08813907089 🕴 E-mail ID: ijraset@gmail.com

Study on Complete Fuzzy Graph and Strong Fuzzy Graph

T Mahalakshmi¹

¹ Asst Professor, Dept. Of Mathematics St. Joseph's College for Women, Hosur.

Abstract: In this paper we define some basic definitions and some properties of fuzzy graph. We compare the general concept of complete fuzzy graph and strong fuzzy graph.

Key words: fuzzy graph, complete fuzzy graph, strong fuzzy graph.

I. INTRODUCTION

It is quite well known that graphs are simply models of relations. A graph is a convenient way of representing information involving relation between objects. Here objects are represented by vertices and relations are represented by edges. Azriel Rosenfeld introduced fuzzy graphs in 1975. It has been growing fast and has numerous applications in various fields. The first definition of a fuzzy graph was by Kaufmann in 1973, based on zadeh's fuzzy relations. But it was azriel Rosenfeld who considered fuzzy relation on fuzzy sets and developed the theory of fuzzy graph in 1975. The concept of signed fuzzy graph is used if the relationship between each pair of nodes is symmetric

II. BASIC PRELIMINARIES

A. Definition 2.1

Let V be a non empty set. A fuzzy graph is a pair of function. G: (σ, μ) where σ is a fuzzy subset of V and μ is a symmetric fuzzy relation on σ . i.e σ : V x [0,1] and μ : V x V \rightarrow [0,1] such that $\mu(u,v) \leq \sigma(u) \wedge \sigma(v)$ for all u,v in V. Here \wedge denote minimum.

We denote the underlying (crisp) graph of G: (σ, μ) by G^{*}(σ^*, μ^*) where σ^* is referred to as the (nonempty) set V of nodes and $\mu^*=E \leq V \times V$. Note that the crisp graph (V,E) is a special case of a fuzzy graph with each vertex and edge of (V,E) having degree of membership 1.

B. Definition 2.2

The fuzzy graph H:(T, v) is called a fuzzy sub graph of G: (σ , μ) induced by P if P \leq V, $T(u)=\sigma(u)$ for all $u, v \in P$. Remark σ is a fuzzy sub sets of a non empty set V and μ is symmetric fuzzy relation on σ

C. Definition 2.3

The graph G * = (V,E) is called the underlying crisp graph of the fuzzy graph G = (σ , μ) where V = { u: σ (u) >0) = support of σ and E= {(u,v) ϵ V x V : μ (u,v) > 0} = support of μ

D. Definition 2.4

A fuzzy graph G: (σ, μ) is said to be strong fuzzy graph if $\mu(u,v) = \sigma(u) \wedge \sigma(v)$ if for all $(u,v) \in E$


E. Definition 2.5

A fuzzy graph G: (σ, μ) is said to be complete fuzzy graph if $\mu(u,v) = \sigma(u) \wedge \sigma(v)$ if for all u,v εV

F. Definition 2.6

The degree of a vertex of G is denoted by $d_G(u)=\Sigma_{u\neq v} \mu(u,v)$. The minimum degree of G is $\delta(G)=\min\{d(v)\}$ and the maximum degree of G is $\Delta(G)=\max\{d(v)\}$

Example of fuzzy graph



Fuzzy graph (F1)

Emerging Trends in Pure and Applied Mathematics(ETPAM-2018)- March 2018

Consider G : (V,E) where $V = \{u_1, u_2, u_3, u_4\}$ and $E = \{u_1u_2, u_3u_4, u_2, u_3, u_4u_1\}$

G. Theorem 2.1:

In fuzzy graph $\Sigma d_G \sigma(u) = 2 \Sigma \mu(u,v)$ that is sum of degree of vertices of G is equal to twice the sum of degree of membership of all those edges which are incident on a vertex $\sigma(u)$


 $\begin{array}{ll} \mbox{From figure F1}, \ d(\sigma \ (u_1) \)= \ \mu \ (u_1,u_2) \ + \ \mu \ (u_1,u_4) \ = 0.3 + 0.5 = 0.8 \\ \mbox{Like that} & \ d(\sigma \ (u_2)) \ = 0.5 \ , \ d(\sigma \ (u_3)) \ = 0.6 \ , \ d(\sigma \ (u_4)) \ = 0.9 \\ \mbox{We get} & \ \Sigma \ d_G \sigma \ (u) \ = 0.8 + 0.5 + 0.6 + 0.9 \ = 2.8 \\ \mbox{Now} & \ \Sigma \ \mu \ (u,v) \ = \ \mu \ (u_1,u_2) \ + \ \mu \ (u_2,u_3) \ + \ \mu \ (u_2,u_4) \ + \ \mu \ (u_4,u_1) \\ & = 0.3 + 0.2 + 0.4 + 0.5 \\ & = 1.4 \end{array}$

Then we get $\Sigma d_G \sigma(u) = 2 \Sigma \mu(u,v)$.

H. Theorem2.2

Every complete fuzzy graph is strong fuzzy graph but converse is not true


1) Proof: A fuzzy graph G: (σ, μ) is strong fuzzy graph if if $\mu(u,v) = \sigma(u) \wedge \sigma(v)$ if for all $(u,v) \in E$ and it is complete fuzzy graph if $\mu(u,v) = \sigma(u) \wedge \sigma(v)$ if for all $u,v \in V$



Complete fuzzy graph (fig2)

$$\begin{split} E &= \{ \ u_1 u_2, u_2 u_4 \ , \ u_4, u_3, \ u_1 \ u_3 \} \ , V = \{ \ u_1, u_2, u_3, u_4 \} \\ Here & \mu \ (u_1, u_2) \ = \ \sigma \ (u_1) \ ^{\wedge} \sigma \ (u_2) \ = \ 0.4^{\wedge} 0.6 \ = \ 0.4 \\ & \mu \ (u_2, u_4) \ = \ \sigma \ (u_2) \ ^{\wedge} \sigma \ (u_4) \ = \ 0.6^{\wedge} 0.3 \ = \ 0.3 \\ & \mu \ (u_3, u_4) \ = \ \sigma \ (u_3) \ ^{\wedge} \sigma \ (u_4) \ = \ 0.5^{\wedge} 0.3 \ = \ 0.3 \\ & \mu \ (u_3, u_1) \ = \ \sigma \ (u_3) \ ^{\wedge} \sigma \ (u_1) \ = \ 0.5^{\wedge} 0.4 \ = \ 0.4 \end{split}$$

All edges satisfies the condition of complete graph i.e, $\mu(u,v) = \sigma(u) \wedge \sigma(v)$ if for all $(u,v) \in E$




Emerging Trends in Pure and Applied Mathematics(ETPAM-2018)- March 2018

$$\begin{split} E &= \{ \begin{array}{l} u_1 u_2 \,, u_2 u_3 \,\,, \, u_3 , u_1 \, \} \,\,, V {=} \{ \begin{array}{l} u_1 , u_2 \,, u_3 \, \} \\ Here & \mu \, \left(u_1 , u_2 \right) \,= \, \sigma \, \left(u_1 \right) \,^{\wedge} \sigma \, \left(u_2 \right) = 0.6 \,^{\wedge} 0.4 = 0.4 \\ & \mu \, \left(u_2 , u_3 \right) \,= \, \sigma \, \left(u_2 \right) \,^{\wedge} \sigma \, \left(u_3 \right) = 0.4 \,^{\wedge} 0.3 = 0.3 \\ & \mu \, \left(u_3 , u_1 \right) \,= \, \sigma \, \left(u_3 \right) \,^{\wedge} \sigma \, \left(u_1 \right) = 0.3 \,^{\wedge} 0.6 = 0.3 \end{split}$$

All vertices satisfies the condition of strong graph i.e, $\mu(u,v) = \sigma(u) \wedge \sigma(v)$ if for all $u, v \in V$

Now we compare fig 1 and fig 2, in complete fuzzy graph all elements of E satisfies the condition μ (u,v)= σ (u) $\wedge \sigma$ (v) so clearly edge set E covers all the vertices therefore it automatically become strong fuzzy graph.

Conversely, Some time particular edges covers all vertices in that case given graph is satisfies only strong fuzzy graph but it does not satisfy complete fuzzy graph.





Given fuzzy graph satisfy strong fuzzy graph i.e, $\mu (u_1, u_2) = \sigma (u_1) \wedge \sigma (u_2) = 0.3 \wedge 0.4 = 0.3$ $\mu (u_2, u_4) = \sigma (u_2) \wedge \sigma (u_4) = 0.3 \wedge 0.4 = 0.3$ $\mu (u_1, u_3) = \sigma (u_1) \wedge \sigma (u_3) = 0.3 \wedge 0.6 = 0.3$ but it is not complete fuzzy graph (since $\mu (u_1, u_4) \neq \sigma (u_1) \wedge \sigma (u_4)$)

III. CONCLUSION

We have discussed the complete fuzzy graph and strong fuzzy graph. Finally every complete fuzzy graph is strong fuzzy graph but not conversely.

REFERENCES

- [1] Dr.G.Nirmala and s.Prabavathi "Characteristics of fuzzy Petersen graph withfuzzy rule" international journal of science and research vol3 Nov 201
- [2] Dr.G.Nirmala and s.Prabavathi "Application of fuzzy if-then rule in fuzzy Petersen graph" international journal of science and research vol3 Nov 2
- [3] A.Muneera , Dr.R.V.N.Srinivas rao"Study on fuzzy graphs and applications" Bulletin of mathematics and statistics research Vol 4 S1 2016
- [4] J.A. Bondy and U.S.R. Murty, Graph Theory With Application.











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24*7 Support on Whatsapp)